activity both in the absence and presence of the learned input and does not respond to the input. The other behavior shows chaotic behavior wandering among the memorized patterns in the absence of the input and responds significantly to the input. Remarkably, the network structure for the latter is similar to that in the previous given model. It suggests that wandering spontaneous behavior is helpful to respond to external familiar stimuli.

2PT225 大腸菌の複数の鞭毛モーターの回転方向の協調的な切り換え現象の理論

A theory of the coordinated switching in the rotational direction of multiple flagellar motors in a single bacterium

Toshinori Namba^{1,2}, Tatsuo Shibata² (¹Department of Mathematical and Life Sciences, Hiroshima University, ²RIKEN CDB)

Many living organisms have systems to respond to the change in external environments, such as signal transduction systems. Such systems responsible for many important processes consist of complex interaction of molecular components. In these systems, molecular signals need to propagate in the cytoplasm. Terasawa et al. have recently reported a coordinated switching in the rotational direction of the multiple flagellar motors in an *E. coli* [1], suggesting that signaling molecules could propagate in cytoplasm to coordinately regulate the motors. The bacteria regulate the swimming behavior by controlling the rotational direction of flagellar motors to respond to extracellular stimuli. Thus, this coordination in the motor rotation may be important for the efficient regulation of the swimming behavior. This coordinated motor behavior can be explained by a spatial propagation of response regulator molecule. In this presentation, we propose a mathematical model of spatial regulation of response regulator molecules. We show possible behaviors of the signal molecules in the cytoplasm, such as wavelike propagation. Based on such a theoretical results, we will discuss the condition of the coordinated motor behaviors.

[1] S. Terasawa et al. (2011) Biophys. J. 100, 2193-2200.

2PT226 走化性真核細胞の変形と勾配認識

Gradient sensing of deformed eukaryotic chemotactic cells

Akinori Baba, Tetsuya Hiraiwa, Tatsuo Shibata (CDB, Riken)

Eukaryotic chemotactic cells sense the gradient of ligand molecules from the noisy input of the ligand binding/unbinding of the receptors distributed around the surface. For example, Dictyostelium Discoideum cells are possible to sense a shallow (about 1-2% of) gradient. To understand the eukaryotic chemotaxis, a theoretical model has been studied in terms of the maximum likelihood estimation of the chemical gradient from the noisy input. If the gradient is sufficiently shallow and/or the ligand concentration is sufficiently dilute, it is hard for a cell to find the correct direction from the ligand binding/unbinding of the receptors. In this case, the estimated direction of the gradient is strongly affected by the input noise. We consider the cell deformation to an elliptical shape in this model. For the shallow limit of the gradient, the estimated direction for an elliptical cell is anisotropic due to the anisotropic distribution of the receptors. We will discuss the comparison between the two scenarios of the gradient estimation: the simultaneous estimation of the steepness and the direction, and the estimation of only the direction with knowing the correct steepness.

2PT227 アメーバ運動による3次元経路探索の理論的考察

A theoretical study for three-dimensional path finding by amoeboid migration

Shin I. Nishimura (Hiroshima University)

Amoeboid cells such as lymphocytes migrate within collagen web matrix and pass through narrow channels between cells. It is known that T lymphocytes inhabiting lymph nodes migrate with searching antigen presenting cells (APC) that present acceptable antigens. Neutrophils appearing from vessels should find and catch external microbes within the body. Amoeboid cells in our body should migrate within complex structure. It is not well known how those cells arrive at appropriate positions through complex maze-like substrata. Even if external signal guides them, amoebae sometimes turn their direction to avoid obstacles. How they keep persistent migration even in such obstacle-rich space? To answer those questions, I construct a model for a three-dimensional amoeba and obstacles. The model cell has membrane, actin and control factor (CF) that prevent actin polymerization. CF is assumed to accumulate by retraction but become diluted by protrusion. This mechanism amplifies fluctuation of CF distribution and leads to directional migration of the cell. The simulated cell shows amoeboid-like migration not only on plane surface but also in a three-

dimensional jungle gym that imitates complex structure within organs. This result implies that the strategy of amoeboid migration is versatile for any environment.

2PT228 一細胞分布解析を用いた大腸菌運動性の非遺伝的な影響評価

Single cell distribution analysis of motility reveals the heterogeneity of Escherichia coli

Daisuke Takagi, Kazumi Hakamada, Jun Miyake (Osaka university)

Genotype does not always agree with the phenotype, though cells can maintain their function. Such inconsistency leads us the idea that the system is made robust. In recent study, many reports try to elucidate their robustness and stable of the system using single cell analysis. Taniguchi et al. reported the distribution of gene expression follow gamma distribution, and Ozbudak et al. reported the multi-stability of bacterial system. These suggest it is difficult to elucidate the systems without understanding their distribution. In this study, four deletion mutants of Escherichia coli ($\Delta flhC$, $\Delta yhjH$, $\Delta yfiN$, and $\Delta ycgR$) and their parental strain BW2513 were selected and distributions of motility of single cells were observed. The peaks of their logarithmic distributions were observed 3.7 (BW25113), 3.2 ($\Delta flhC$), 3.5 ($\Delta yhjH$), 4.3 ($\Delta yfiN$), and 5.5 ($\Delta ycgR$), respectively and they had statistical significance ($p < 0.001$). The peaks of order corresponds to the function of genes. In the case of using $\Delta ycgR$, the bimodal distribution of was given and it means this system has multi-stability. On the contrary, unimodal distribution was given in the case of using BW25113. This result suggests $ycgR$ plays important role of giving stability of the system of flagella. It is necessary to construct mathematical model to elucidate the system.

2PT229 時計タンパク質 KaiB-KaiC 相互作用間の ATP による制御

The regulation of cyanobacterial circadian clock proteins KaiB-KaiC interaction by ATP

Risa Mutoh¹, Atsuhito Nishimura², So Yasui², Kiyoshi Onai², Masahiro Ishiura² (¹Institute for Protein Research, ²Center for Gene Research)

The cyanobacterial circadian clock oscillator is composed of three clock proteins called KaiA, KaiB and KaiC. The oscillator generates circadian oscillations in the phosphorylation level, ATPase activity of KaiC and complex formation among the three Kai proteins in the presence of ATP. KaiC is a homohexameric ATP-binding protein with autokinase activity and very weak temperature-independent ATPase activity. The KaiC subunit has a duplicated structure, which is composed of N-terminal and C-terminal domains, and each domain has a series of ATPase motifs (a Walker's motif A, a Walker's motif B and a catalytic glutamate (CatE)).

In the absence of ATP, KaiC was monomeric and formed a complex with KaiB. In the presence of Mg^{2+} and ATP (Mg-ATP), KaiC became a homohexameric ATPase with bound Mg-ATP and formed a complex with KaiB, whereas the KaiCs hexamerized by unhydrolyzable substrates such as ATP and Mg-ATP analogs did not. KaiC N-terminal domain protein formed a complex with KaiB, whereas its C-terminal one did not, indicating that KaiC associates with KaiB via its N-terminal domain. A hexameric KaiC mutant lacking N-terminal ATPase activity did not form a complex with KaiB, whereas that lacking C-terminal ATPase activity formed the complex. Thus, the hydrolysis or release of KaiC-bound ATP by N-terminal ATPase motifs is required for the complex formation.

2PT230 時計タンパク質 KaiC の ATP 放出

ATP release from cyanobacterial circadian clock protein KaiC

Keita Iwata^{1,2}, Risa Mutoh³, Kiyoshi Onai¹, Masahiro Ishiura¹ (¹Center for Gene Research, Nagoya Univ., ²Sch. of Sci., Nagoya Univ., ³Institute for Protein Research)

Cyanobacteria are the simplest organisms that exhibit circadian rhythms. The cyanobacterial circadian clock oscillator is composed of three clock proteins called KaiA, KaiB, and KaiC. KaiC has autokinase and ATPase activities. KaiC subunit has a duplicated structure, which is composed of N-terminal and C-terminal domains, and each domain has a series of ATPase motifs (a Walker's motif A, a Walker's motif B, and a catalytic glutamate (CatE)). KaiC subunit has two phosphorylation sites, Ser431 and Thr432, on its C-terminal domain. In the presence of ATP, ATP binds to the ATPase motifs of KaiC and induces the hexamerization of KaiC.

Here, we investigated ATP release from KaiC by the real time monitoring of bioluminescence using firefly luciferase. We measured the amount of ATP released from KaiC, using wild-type KaiC and mutant KaiCs with a mutation in the phosphorylation sites and ATPase motifs. We observed different ATP