3PT130 陰溶媒膜モデルとレプリカ交換シミュレーションによる α -ヘリカル型膜タンパク質の構造予測**Prediction of α -helical membrane protein structures by replica-exchange simulations with implicit membrane model**

Ryo Urano¹, Yuko Okamoto^{1,2,3} (¹Dept. of Phys. Grad. Sch. of Sci. Nagoya Univ., ²Bio. Research Center Grad. Sch. of Sci. Nagoya Univ., ³Center for Comput. Sci. Grad. Sch. of Eng. Nagoya Univ.)

Membrane proteins are essential for life. Their structures are related to their functions. Their 3-dimensional structures are typically in α -helical structures. Although the determination of the structures is usually done by x-ray diffraction, solid-state NMR, or electron microscope experiments, it takes a long time to prepare the samples. The purpose of the present work is to predict the 3-dimensional structure of α -helical membrane proteins by computer simulation. The procedure is as follows. At first, we extract trans-membrane amino acid residues from membrane proteins by prediction tools based on bioinformatics. Next, the ideal helix structure is formed for the trans-membrane helix regions. During simulations, the replica-exchange method and a simple implicit membrane model to represent restrictions of membrane proteins from membrane environment are applied. The Monte Carlo move sets include backbone torsion angles for the deformation of helix structures from ideal helix formation. This implicit membrane model represents the restriction of trans-membrane helices within the membrane regions. A principal component analysis is used to examine the simulation trajectory and divide structures into local- minimum free energy states. We applied our method to the prediction of the structures of small membrane proteins including bacteriorhodopsin. In this poster presentation, we show the results of these structure predictions.

3PT131 Homology modeling and molecular dynamics study of proton permeation pathway in a channel protein

Susumu Chiba¹, Kota Kasahara¹, Matsuyuki Shiota^{1,2}, Kengo Kinoshita^{1,2,3} (¹Grad. Sch. of Information Sci., Tohoku Univ., ²Tohoku Medical Megabank Organization, Tohoku Univ., ³IDAC, Tohoku Univ.)

Hv1, voltage sensitive proton channel is known as voltage sensor only protein (VSOP). In contrast to other voltage sensitive ion channels, VSOP doesn't have a pore domain but has only a voltage sensor domain (VSD), through which proton permeate. Recent simulation studies based on homology models of VSOP showed the possibility that proton permeation is caused by proton "hopping" (Grotthuss mechanism), from water to water in the crevice of transmembrane region, but details of permeation mechanism is still unclear. To analyze proton pathway, molecular dynamics (MD) simulation is powerful method. Since the proton pathway is formed by the interaction between water and the charged residues in transmembrane region, precise modeling of membrane region is important to simulate and estimate characteristic of proton permeation. VSD of voltage-gated potassium channel (Kv channel) was used for template of homology modeling in the preceding studies. On the other hand, voltage-gated sodium channel (Nav channel) NavAb, which was recently entered in PDB database, share higher sequence identity with VSOP than Kv channel. Therefore we made VSOP-NavAb alignment considering sequence of transmembrane regions, and subsequently we modeled structure based on the alignment. We performed the MD simulation of the predicted model embedded in POPE membrane for 200 ns. It was observed that a continuous pathway of water molecules exist in the crevice of the channel. Finally we discuss the behavior of waters and charged residues, and proton permeation mechanism.

3PT132 大腸菌機械受容チャネル MscL の開口過程におけるタンパク質と水との協調過程の役割の分子動力学的解析**Molecular Dynamics Analysis of the Role of Protein-Water Interplay in the Opening Process of E-coli Mechanosensitive Channel MscL**

Yasuyuki Sawada¹, Masahiro Sokabe^{1,2} (¹Dept. Physiol. Nagoya Univ. Grad. Sch. Med., ²FIRST Res. Ctr. for Innovative Nanobiodevice, Nagoya Univ.)

The bacterial mechanosensitive channel MscL is constituted of homopentamer of a subunit with two transmembrane inner (TM1) and outer (TM2) α -helices. Neighboring TM1s cross and interact with each other near the cytoplasmic side through hydrophobic interaction, forming the most constricted hydrophobic part of the pore. The crossings stabilize the closed state of MscL with hydrophobic interactions, thus called gate. The major issue of MscL is to understand the gating (opening) mechanism driven by tension in the membrane. We performed

molecular dynamics (MD) simulations of tension-driven opening of MscL, particularly focusing on the conformational change of MscL associated with water. Upon membrane stretch, Phe78 in TM2 helix, which is suggested to be a major tension sensor in our previous work, was dragged by lipids and TM1s were gradually tilted accompanied by outward sliding of the crossings between TM1 helices. While TM1s were tilted, a breakage of the helical structure around the gate was shown and an exposure of oxygen atoms of the backbone to the inner surface of the gate occurred, followed by formation of hydrogen bonds between water and the exposed oxygen. This allowed further access of water to the gate and weakened the hydrophobic interaction at the crossings, leading to a further opening of the gate. It is suggested that the combination of conformational changes of MscL by membrane stretch and following hydrogen bond formation forms a positive feedback cycle, which seems to be critical to induce rapid channel opening.

3PT133 ATP-Binding induced conformational change in CFTR as studied by MD simulation

Tomoka Hagiya¹, Tadaomi Furuta¹, Shuntaro Chiba¹, Yoshiro Sohma², Minoru Sakurai¹ (¹Grad. Sch. of Biosci. & Biotech., Tokyo Tech., ²Dept of Pharmacol., Sch. Med., Keio Univ.)

The Cystic fibrosis transmembrane conductance regulator (CFTR, ABCC7) is a chloride channel and belongs to the ATP-binding cassette (ABC) protein superfamily that binds and hydrolyzes ATP to drive substrate transport across membranes. For the gating mechanism, the following hypothesis has been proposed: after phosphorylation of the regulatory domain, ATP molecules bind to the two nucleotide-binding domains (NBDs), which in turn induces the dimerization of the NBDs and the concomitant opening of the channel gate. To clarify its mechanism, we first built a homology model of CFTR (with an inward-facing conformation) by using several X-ray crystal structures of ABC transporters as templates. Subsequently we explored its stable conformations and dynamics using molecular dynamics (MD) simulations, which were performed for Mg and ATP-bound (MgATP) state and nucleotide-free (Free) state. Interestingly, in the MgATP state, NBD1 and NBD2 were dimerized so that the ATP molecules were sandwiched between Walker A and the signature motif. Additionally we found the closing of the TMDs internal cavity in the intracellular loop regions. On the other hand, the NBDs remained monomers in the Free state. This study suggests that ATP-binding not only causes the NBD dimerization, but also exerts a significant influence on the TMD structure.

**3PT134 リボソームと機能性ペプチドを用いた酵素活性検出法の開発
Development of enzyme activity detection system using liposome and functional antimicrobial peptide**

Masayoshi Kashibe¹, Shin Mizukami^{1,2}, Kazuya Kikuchi^{1,2} (¹Graduate School of Engineering, Osaka University, ²Immunology Frontier Research Center, Osaka Univ.)

The development of enzyme activity detection system has attracted wide attention in the field of biology, pharmacology, and medicine. We developed a new system for enzyme activity assay by using anionic liposome and the derivatives of an antimicrobial peptide, temporin L (TL). It was reported that TL has membrane-damaging activity against anionic membranes of micro-organisms. Previously, we reported that modification of the Lys amino group of TL by a photosensitive protective group suppressed its membrane-damaging activity. Thus, we hypothesized the membrane-damaging activity of TL derivatives can be modulated also by modification of other amino acid residues. We designed several TL derivatives modified with a phosphate group at the various amino acid residues. The membrane-damaging activities in some of the derivatives were suppressed and liposome disruption does not occur. When the peptides were treated with a phosphatase, these activities were recovered. This conversion induced the inclusion release from liposome, in which fluorophores were quenched, and the inclusion release induced the fluorescence enhancement. This result indicates that this system can be used for the detection of phosphatase activity.

Since various modifications of peptides for many types of enzyme assay is possible, it could lead to detection system for various enzyme activities. Moreover, since liposomes can include various compounds such as proteins, and drugs, this system would be applied not only to diagnosis but also to therapeutics.