1P049 独立成分分析 tICA を用いたタンパク質主鎖の遅い運動の解析

Slow dynamics of protein backbone in molecular dynamics simulation revealed by time-structure based independent component analysis

Sotaro Fuchigami (*Grad. Sch. of Medical Life Science, Yokohama City Univ.*)

Protein motions occur over a wide range of time scales, but not all are important for protein functions, which time scales are generally longer. Thus, it would be reasonable to consider that slower motions of proteins are more relevant to their functions. To identify such slow protein dynamics from simulation results, we have proposed the time-structure based independent component analysis (tICA). In the present study, we selected lysine-, arginine-, ornithine-binding protein (LAO) as a target protein, and performed one-microsecond molecular dynamics simulation in explicit water. By applying the tICA to the simulation results, several motions in the LAO were identified as slow modes, and were confirmed with additional analyses to be actually occurred.

1P050 トリプリオンタンパク質に対する抗体 G2 の複数の抗原を特異的に認識する性質の特徴付け

Characterization of multispecific monoclonal antibody G2 directed against chicken prion protein

Yuji Kamatari¹, Masayuki Oda², Takahiro Maruno³, Yuji Kobayashi³, Naotaka Ishiguro⁴ (¹*Life Sci. Res. Center, Gifu Univ.*, ²*Grad. Sch. Life Env. Sci., Kyoto Pref. Univ.*, ³*Graduate School of Engineering, Osaka University*, ⁴*Fac. Applied Biol. Sci., Gifu Univ.*)

We generated a monoclonal antibody, designated G2, with residues 174-247 of the chicken prion protein (ChPrPC) and found that G2 could react with at least three proteins other than ChPrPC. Surprisingly, there were no regions of amino acid sequence similarity between ChPrPC and those proteins. The amino acid sequence of the G2 epitope within one of the protein was not related to the G2 epitope within ChPrPC. ELISA, SPR, and ITC experiments indicated that these two epitope peptides have similar binding affinity for G2. However, we found that these two peptides differed substantially in several binding characteristics in SPR and ITC experiments. These differences may be explained by G2 adopting different binding conformations and undergoing different binding pathways.

1P051 リソスタシンのカルサイト結合部位の同定

Identification of calcite-binding site of lithostathine

Seiya Togashi^{1,2}, Yuichi Hanada^{1,2}, Maho Nara^{2,3}, Sakae Tsuda^{1,2} (¹*Grad. Sch. Sci., Hokkaido Univ.*, ²*BPRI, AIST*, ³*Hokkaido High-Tech. Bio.*)

Lithostathine (LIT) binds to surfaces of calcite (CaCO₃) crystal to modify its morphology. LIT is proteolyzed into two fragments (LIT133 and LIT11) by trypsin. To identify the calcite-binding site of LIT, we compared binding efficiency of intact LIT, LIT133 and LIT11 by examining their ability to modify the calcite morphology. Intact LIT modified calcite into a prismatic shape, implying that the protein binds to the prism planes ((110) and (1-10)). LIT133 modified the prism planes less efficiently than intact LIT. No morphological change of calcite was detected for LIT11 even at high concentrations. These results suggest that calcite-binding sites are located in LIT133 portion, where acidic residues aligned in its molecular surface.

1P052 オクタリピート領域をもつプリオンペプチドにおける金属イオンとの競合結合性

Competitive binding of metal ions to octarepeat region of prion protein

Masahiro Yagi, Kazuya Iwama, Haruto Onda, Wakako Hiraoka (*Graduate School of Science and Technology, Meiji University*)

Metal ions binding to prion protein (PrP) is related to cause and progress prion disease. Cu²⁺ binding to octarepeat region in PrP is considered as an important process for misfolding of PrP. In this report, competitive binding of metal ions (Cu²⁺, Co²⁺, Ni²⁺, etc) to the octarepeat peptide (PHGGGWGQ) was investigated with spectroscopic studies. Visible circular dichroism (CD) and visible absorption spectra showed the binding of each metal ion to the peptide. UV-CD spectra indicated the conformational change of secondary structure in the metal binding peptide. To determine the affinity of each metal ion to the peptide, the metal binding peptide was titrated with competitive metal ion, using visible CD and visible absorption spectroscopy.

1P053 QCM によるグルカゴンと酸性膜との相互作用解析

Analysis of interaction between glucagon and acidic lipid membrane by QCM

Takamichi Horie, Ayano Momose, Izumi Yamane, Hideki Fujita, Eri Yoshimoto, Izuru Kawamura, Akira Naito (*Grad. Sch. Eng., Yokohama Natl Univ.*)

Glucagon is a peptide hormone consisting of 29 amino acid residues, and forms amyloid-like fibrils in aqueous solutions. Previously, kinetics of fibrillation in the presence of neutral (DMPC) and acidic (DMPC + DMPG) lipid membrane was investigated under acetic acid solution by solid state NMR. It suggested that fibril nucleation rate in the presence of acidic lipids was faster than that in the neutral lipids. In this study, association constants (K_a) between glucagon and lipids were determined by QCM. As a result, K_a values significantly rose upon increasing acidity of lipids. Membrane disruption was also observed in the lipids including 25% of DMPG. It turned out that glucagon-acidic lipids interaction accelerates the fibril nucleation rate.

1P054 ジンジバインプロテアーゼがもつ Ig-like domain の役割

Function of the Ig-like domain of gingipain proteinase

Keiko Sato¹, Hideharu Yukitake¹, Daisuke Nakane², Satoshi Shibata¹, Yuka Narita¹, Koji Nakayama¹ (¹*Nagasaki Univ.*, ²*Gakushuin Univ.*)

Porphyromonas gingivalis possesses cysteine proteinases, named gingipains, which are important virulence factors of the periodontal pathogen. A proprotein of Lys-gingipain (Kgp) consists of signal peptide, propeptide, proteinase domain, Ig-like domain, adhesin domains and C-terminal domain. Ig-like domains, which are found in a variety of proteins, are the widespread structural motif that has a sandwich-like structure formed by two sheets of antiparallel beta strands. Kgp lacking the Ig-like domain was degraded by the HtrA protease that plays a role in protein quality control in the periplasm of Gram-negative bacteria, suggesting that the Ig-like domain functions as an intra-molecular chaperone.