1P218 細胞侵入ペプチドであるトランスポータン 10 の脂質膜透過はポア形成の前に起こる**Permeation of Cell-Penetrating Peptide Transportan 10 through Lipid Membranes before Pore Formation**

Md. Zahidul Islam¹, Hirotaka Ariyama¹, Md. Jahangir Alam², Masahito Yamazaki^{1,2} (¹*Grad. Sch. Sci. & Tech., Shizuoka Univ.*, ²*Res. Inst. Electronics, Shizuoka Univ.*)

The elementary process and the mechanism of permeation of cell-penetrating peptide transportan 10 (TP10) through lipid membranes are not well known. Here we investigated the permeation of carboxyfluorescein(CF)-labeled TP10 through lipid membranes and the leakage of water-soluble fluorescein probe Alexafluor (AF) in 20%DOPG/80%DOPC-GUVs simultaneously using the single GUV method with confocal microscopy. We found that CF-TP10 entered into the inside of GUV from the outside before starting leakage of AF, i.e., the pore formation. The fraction of entry of CF-TP10 among all the observed GUVs increased with an increase in CF-TP10 concentration. Through the TP10-induced pores, larger molecules such as dextran 10K and lipid vesicles passed through the membranes.

1P219 Interaction Of Warm-Sensing Chemical Capsaicin with the Biomimetic Membranes

Neha Sharma, Pooja Gusain, Tsuyoshi Yoda, Masahiro Takagi (*Japan Advanced Institute of Science and Technology*)

Capsaicin, one of main components of chilli peppers produces warming sensation. It is used in pain relievers, topical ointments. It specifically interacts with TRPV1 membrane receptor and depletes a substance which is transmitter of pain and heat. However, the non-specific interaction of Capsaicin and lipid bilayers are not negligible.

Here, cell-sized liposome was used to investigate thermosensitive response of Capsaicin-containing membranes. It was shown that capsaicin-containing membranes are more thermosensitive than control at high temperature range. This could be that the hydrophobic interaction between the non-polar tail groups of lipid membrane and Capsaicin. These findings are useful to understand mechanism behind the effect of capsaicin on biological membranes.

1P220 Dynamic Response of Menthol on Thermo-Induced Cell Membrane: More than Receptors

Pooja Gusain, Neha Sharma, Tsuyoshi Yoda, Masahiro Takagi (*Japan Advanced Institute of Science and Technology*)

It is important to understand the physicochemical mechanism that is responsible for the morphological changes in the cell membrane in the presence of various stimuli and sensory compounds. Menthol, popularly known for its cooling effect, activates TRPM8- a cold activated thermoTRP ion channel. We are interested in direct interaction of menthol with bio-membrane.

In this study, we demonstrate how menthol affects the dynamical movement of thermo-induced membrane and reveal unclear mechanism behind cooling sensation of menthol. Hence we used cell-sized liposomes to study the cell dynamics. We found that menthol containing liposomes are more thermo-sensitive. This can be explained by hydrophilic interaction between menthol with head group of lipid molecules.

1P221 急速緩衝液交換法による時間分解全反射赤外分光法の開発 Development of a Rapid Buffer-Exchange System for Time-Resolved ATR-FTIR Spectroscopy with the Step-Scan Mode

Yuji Furutani^{1,2}, Tetsunari Kimura^{1,2}, Kido Okamoto³ (¹*Inst. Mol. Sci.*, ²*SOKENDAI*, ³*UNISOKU*)

Attenuated total reflectance (ATR)-FTIR spectroscopy has been widely used to probe protein structural changes under various stimuli. Here we developed a novel, rapid buffer-exchange system for time-resolved ATR-FTIR spectroscopy to monitor the ligand- or ion-binding reaction of a protein. By using the step-scan mode (time resolution; 2.5 ms), we confirmed the completion of the buffer-exchange reaction within ~25 ms; the process was monitored by the infrared absorption change of a nitrate band at 1350 cm⁻¹. We also demonstrated the anion-binding reaction of a membrane protein, *Naatronomonas pharaonis* halorhodopsin (pHR). The formation of chloride- or nitrate-bound pHR was confirmed by an increase of the retinal absorption band at 1528 cm⁻¹.

1P222 哺乳類 two-pore 型カリウムチャネル TWIK-1 の全反射赤外分光解析**ATR-FTIR spectroscopic analyses on a mammalian two-pore domain potassium channel, TWIK-1**

Hisao Tsukamoto¹, Koichi Nakajo², Yoshihiro Kubo², Yuji Furutani¹ (¹*Institute for Molecular Science*, ²*National Institute for Physiological Sciences*)

A mammalian two-pore domain potassium channel TWIK-1 (or KCNK1, K2P1) contributes to generate a "leak" K⁺ conductance that is essential for the resting membrane potential. TWIK-1 and other two-pore domain channels possess two selectivity filter (pore) domains per subunit and function as a dimer, although conventional potassium channels have one pore domain per subunit and function as a tetramer. Interestingly, TWIK-1 can permeate Na⁺ under some conditions whereas other mammalian potassium channels are highly K⁺-selective. The mechanism underlying the unique property is unclear whereas crystal structure of TWIK-1 was solved. In this study, we carried out ATR-FTIR spectroscopic analyses on TWIK-1 with an aim to reveal how the channel accomplishes the unique function.

1P223 有効電場中におけるグラミシジン A を含んだ脂質二重層膜の静電ポテンシャルと圧力特性**Electrostatic potential and lateral pressure profile of lipid bilayer containing gramicidin A in effective electrostatic field**

Hiroaki Saito, Kazutomo Kawaguchi, Hidemi Nagao (*Kanazawa University*)

The linear peptide gramicidin A (GA) forms an ion channel specific for monovalent cation and has been extensively used to study the organization, dynamics and function of GA channels. On the other hand, the electric field and lateral pressure of the membrane influences the permeability of polar small molecules and ions across the membrane. In this study, we carry out a molecular dynamics (MD) simulation of a GA in the DMPC lipid bilayers to investigate the structure of GA/DMPC membrane and the ion selectivity of the GA channel. We provide the electrostatic potential maps of the GA/DMPC membrane in the effective electrostatic field. The lateral pressure profile across the membrane is estimated, and the mechanism of ion permeation of the GA in the membrane is discussed.