2P037 蛋白質のミスフォールド状態から生じる蛋白質の異常凝集
Misfolding triggers a pathogenic conversion of protein conformations

Soichiro Kitazawa¹, Ryo Kitahara¹, Makoto Urushitani² (¹*Pharmaceutical science, Ritsumeikan Univ.*, ²*Molecular Neuroscience Research Center, Shiga University of Medical Science.*)

Although conformational fluctuations of a protein among different states have been recognized to be important for protein function, such fluctuations may trigger a pathogenic conversion of protein conformations. Aggregation of a nuclear protein TDP-43 in abnormal cytoplasmic inclusions in motor neurons is a pathological signature of sporadic ALS. Here, we demonstrate that pressure perturbation accelerates an irreversible misfolding of RNA recognition motif-1 (RRM1) of TDP-43, which triggers aggregation of TDP-43. NMR spectroscopy revealed structure and dynamic characteristics of the misfolded conformation of RRM1.

2P038 べん毛輸送装置構成蛋白質 FliP ペリプラズミックスループの結晶化と遺伝学的解析
Crystallization and genetic analyses of a periplasmic loop of FliP, a component of the flagellar protein export apparatus

Takuma Fukumura¹, Yumiko Saijo-Hamano¹, Yukio Furukawa¹, Tatsuya Kawaguchi², Katsumi Imada², Keiichi Namba¹, Tohru Minamino¹ (¹*Grad. Sch, Frontier Biosci., Osaka Univ.*, ²*Grad. Sch. Sci. Osaka Univ.*)

Most of flagellar proteins are transported by a specific export apparatus from the cytoplasm to the growing distal end of the flagellum where they self-assemble. The export apparatus consists of a water-soluble ATPase complex and a proton-driven export gate made of six membrane proteins. FliP is one of the component of the gate complex. FliP is predicted to have four transmembrane helices (TM) and has a relatively large periplasmic loop (FliPp) between TM-2 and TM-3. Genetic analysis showed that the C-terminal region of FliPp, especially Glu-178, is important for the export function. We overexpressed and purified *Thermotoga maritima* FliPp and found that FliPp forms homo tetramer. We also succeeded in crystallizing FliPp, and the crystal diffracted to 2.2 Å resolution.

2P039 Structural defects in fibrillin associated with Marfan syndrome

Yaxin Lu¹, Richmond Jeremy², Murat Kekic¹, Jianlin Yin², Brett Hambly¹ (¹*Pathology Discipline and Bosch Institute, Sydney Medical School, University of Sydney*, ²*Central Clinical School, Sydney Medical School, University of Sydney*)

Fibrillin (FBN) is a 350kDa glycoprotein vital for formation of elastic and non-elastic fibres in connective tissue. A FBN protein complex also controls TGF-β bioavailability. FBN mutations cause Marfan syndrome (MFS), leading to both defective microfibrils and dysregulation of TGF-β signalling. FBN1 consists of repeating epidermal growth factor (EGF) and TGF-β-binding protein (TB) domains. Paired EGF domains bind Ca²⁺ for structural rigidity. 600 MFS mutations in FBN have been classified in terms of their structural and functional consequences. A large majority of missense mutations alter EGF tandem repeat structural rigidity, by either altering disulphide formation, Ca²⁺ binding or domain interaction. Some mutations in TB domains alter storage of latent TGF-β.

2P040 Microtubule-associated protein 4-mediated bundle formation of microtubules and actin filaments

Shoma Saito¹, Ayumu Kuramoto¹, Hikari Makihara¹, Tsuyoshi Yamazaki², Taro Q.P. Noguchi², Susumu Kotani³, Kiyotaka Tokuraku¹ (¹*Grad. Sch. Appl. sci., Muroan Inst.*, ²*Miyakonojo Nation. Col. Tech.*, ³*Kanagawa University*)

Microtubule-associated protein (MAP) 4 is a non-neural MAP that plays important roles in the intracellular organization of microtubules. Recent study revealed that MAP4 also binds to actin filaments at the C-terminal part of the Proline-rich region in the microtubule-binding domain (Matsushima et al., J. Biochem., 2012). In this study, we examined MAP4 functions when both actin filaments and microtubules coexisted in a sample as well as inside cells by fluorescence microscopic observation and coprecipitation assay. The results showed that MAP4 bound to microtubules lost actin filament-binding ability. On the other hand, MAP4 bound to actin filaments retained microtubule-binding and microtubule assembly-promoting abilities.

2P041 タンパク質の圧電効果とアロステリック制御
Piezoelectric effect in a protein and its involvement in allosteric regulation

Jun Ohnuki¹, Takato Sato¹, Koji Umezawa¹, Taro Q.P. Uyeda², Mitsunori Takano¹ (¹*Grad. Sch. of Adv. Sci. & Eng., Waseda Univ.*, ²*Biomedical Res. Inst., AIST*)

Intermolecular interaction of proteins is often regulated by mechanical strain. For instance, the affinity between myosin and actin filament is affected by the internal strain of myosin due to lever-arm swing or by the strain imposed on the actin filament. However, its molecular mechanism is not well-established. By conducting molecular dynamics simulation, we first investigated the response within the myosin molecule that accompanies forced lever-arm swing, and discovered an electric charge response in the actin-binding region via large-scale hydrogen-bond rearrangements: myosin behaves like a piezoelectric actuator. We will examine if this piezoelectricity actually affects the myosin-actin interactions, and if it also occurs in actin filaments.

2P042 MDで観測されたGアクチンのヌクレオチド依存構造状態とFアクチン安定性との関連
Nucleotide-dependent structural states of G-actin observed by MD simulation and its implication for F-actin stability

Jun Ohnuki, Mitsunori Takano (Dept of Phys & Appl Phys, Waseda Univ)

Actin treadmill is fundamental to cellular motility, which is based on ATP-consuming transformation between monomeric (G) and filamentous (F) states. It is obvious that the ATP energy drives treadmill, but how it is used is unclear, even though the structural change upon ATP hydrolysis, which is accelerated in F-state, is likely to hold the key. We thus conducted MD simulation of G-actin to investigate intrinsic structural difference between the ATP and ADP-bound states. Extensive MD simulation allowed us to obtain statistically-definite results: ADP-bound actin is more rigid and less conforming to F-state, indicating how F-actin is destabilized upon ATP hydrolysis. By analyzing hydrogen-bond network we further clarify how this difference arises.