2P067 β_2 -ミクログロブリンのアミロイド形成における様々な脂肪酸の効果

Effects of various fatty acids on the amyloid fibrillation of β_2 -microglobulin

Akira Ishii¹, Masatomo So¹, Hisashi Yagi¹, Hironobu Naiki², Yuji Goto¹ (¹*Inst. Protein Res., Osaka Univ.*, ²*Fac. Med. Sci., Univ. Fukui*)

Amyloid fibrillation is dominated by the environmental factors. Although various additives, such as SDS, phospholipids and RNA aptamers, are known to accelerate and/or inhibit fibrillation, little is known about the mechanism how these additives effect fibrillation. Clarifying the mechanisms of acceleration and inhibition of fibrillation by additives is useful not only to treat amyloid-related diseases but also to understand physical properties of amyloid fibrils. In this study, we investigated the dependence of fibrillation on the concentration of various fatty acids and found that there is an optimal concentration for each compound to accelerate fibrillation and that fibrillation was inhibited under the high concentrations.

2P068 β_2 ミクログロブリンのアミロイド前駆状態の残余構造の特性化

The properties of the residual structure of amyloid precursor state of β_2 -microglobulin

Kazumasa Sakurai^{1,2}, Akihiro Maeno¹, Hironobu Naiki³, Yuji Goto², Kazuyuki Akasaka¹ (¹*HPPRC, Kinki Univ.*, ²*Inst. Protein Res., Osaka Univ.*, ³*Fac. Med. Sci., Univ. Fukui*)

β_2 -Microglobulin (β_2 m), associated with dialysis-related amyloidosis, is known to form amyloid fibrils at low pH via a partially structured state. Based on the results of the various NMR experiments, we previously suggested the partially structured state to be “activated” to become extensively unfolded, in which state the hydrophobic residues are exposed and associate with the seed. In addition, under high pressure conditions, increase in HSQC signal intensity over the sequence and decrease in transverse relaxation rate on the 60’s aromatic-rich region were observed, suggesting that β_2 m is experiencing inter- and intramolecular association processes simultaneously. We will further discuss the thermodynamic nature of the interactions.

2P069 超音波によるアミロイド線維形成促進のメカニズム The mechanism of ultrasonication-induced amyloid fibril formation

Masatomo So¹, Yuichi Yoshimura¹, Hisashi Yagi¹, Hirotugu Ogi², Kentaro Uesugi², Hironobu Naiki³, Yuji Goto¹ (¹*Institute for Protein Research, Osaka University*, ²*Graduate School of Engineering Science, Osaka University*, ³*Faculty of Medical Sciences, University of Fukui*)

Amyloid fibril formation takes a long time because of a high free-energy barrier of nucleation. Although ultrasonication has been used for breaking the mature fibrils, it found to be one of the most powerful methods for accelerating spontaneous fibril formation. However, the mechanism is still unclear. To understand the effects of ultrasonication on fibril formation, we measured ultrasonic wave using a piezoelectric probe. The lag time of fibril formation was significantly correlated with ultrasonic amplitude. The results indicate that ultrasonic amplitude is a dominant factor determining the fibril formation. We will discuss the effects of ultrasonication on fibril formation and fibril morphologies.

2P070 Solubility and Supersaturation-Dependent Protein Misfolding Revealed by Ultrasonication

Yuxi Lin, Young-Ho Lee, Yuichi Yoshimura, Hisashi Yagi, Yuji Goto (*Institute for Protein research, Osaka University*)

We studied the alcohol-induced fibrillation of lysozyme at various concentrations of ethanol, trifluoroethanol (TFE) and hexafluoropropanol (HFIP). Under the conditions where the alcohol-denatured lysozyme retained metastability, ultrasonication effectively triggered fibrillation. The optimal alcohol concentration depended on the alcohol species. HFIP showed a maximum at 12-16%. For TFE, a broad maximum at 40-80% was observed. Ethanol showed an increase of fibrillation above 60%. These profiles were opposite to the equilibrium solubility of lysozyme in water/alcohol mixtures. The results indicate that, although fibrillation is determined by solubility, supersaturation prevents conformational transition and that ultrasonication is effective in breaking supersaturation.

2P071 熱測定によるアミロイド線維形成バーストに関する研究 Direct observation of burst of amyloid fibril formation by calorimetry

Tatsuya Ikenoue¹, Young-Ho Lee¹, Jozsef Kardos², Yuji Goto¹ (¹*Inst. Pro. Res., Osaka Univ.*, ²*Inst. Bio., Eotvos Lorand Univ.*)

Although amyloid fibrils are related to various pathologies, energetic natures of fibrillation remain largely unclear. We show bursting formation of amyloid fibrils of β_2 -microglobulin (β_2 m) using isothermal titration calorimetry. β_2 m amyloid burst occurred with emitting large heat after a lag phase for nucleation. Enthalpy changes (ΔH) of β_2 m misfolding were smaller than β_2 m folding, and displayed a negative heat capacity change (ΔC_p) related to buried area. By contrast, amorphous aggregation was followed by smaller ΔH than amyloid burst without a lag phase. Interestingly, (mis)folding and aggregation of acid-denatured β_2 m showed similar ΔC_p values, implying similar buried area following structural conversions with distinct level of ordered molecular contacts.

2P072 Efficient Lookup Table using a Linear Function of Inverse Distance Squared

Jaewoon Jung¹, Takaharu Mori^{2,3}, Yuji Sugita^{1,2,3} (¹*AICS, Riken*, ²*Riken*, ³*QBiC, Riken*)

The major bottleneck in molecular dynamics (MD) simulations of biomolecules exists in the calculation of short-range non-bonded interactions. Non-bonded interaction includes time-consuming inverse square roots and complementary error functions in particle mesh Ewald (PME). To avoid such time-consuming operations while keeping accuracy, we propose a new lookup table for short-range interaction by defining energy and gradient as a linear function of inverse distance squared. In our lookup table approach, densities of table points are inversely proportional to squared pair distances, enabling accurate evaluation of energy and gradient at small pair distances. The new lookup table scheme allows fast pairwise non-bonded calculations owing to efficient usage of cache memory.