2P121 Local structural similarity between interphase chromatin and mitotic chromosomes in living mammalian cells

Tadasu Nozaki^{1,2}, Tomomi Tani³, Sachiko Tamura¹, Takeharu Nagai⁴, Kazuhiro Maeshima¹ (¹Natl. Inst. Genet., ²Inst. Adv. Biosci., Keio Univ., ³Marine Biological Laboratory, ⁴ISIR, Osaka Univ.)

The long string of genomic DNA must be organized three dimensionally in the cell to utilize genome information. Recently we demonstrated that the nucleosome fiber (10-nm fiber), in which DNA is wrapped around core histones, is irregularly folded in the cell without a 30-nm chromatin fiber. Our super-resolution nucleosome 2-D mapping identified numerous compact chromatin domains in both interphase chromatin and mitotic chromosomes in living cells. This finding provides us a novel view on the mitotic chromosome assembly: in the interphase nuclei, numerous compact chromatin domains are already formed. During mitosis, these chromatin domains are assembled together, presumably by condensins and other protein factors to create a rod-like chromosomal shape.

2P122 レドックス DNA の電子移動反応に及ぼす二本鎖内架橋の影響
Effect of intreststrand cross-linking of redox-labeled DNA duplex on its electron transfer reaction

Yasuhiro Mie, Keiko Kowata, Yasuo Komatsu (*Bioproduction Res. Inst., AIST*)

The electrochemical investigation of interstrand cross-linked DNA (ICL-DNA) would help us to understand the biological DNA electron transfer and to construct efficient bioelectronics as well. In this study, the first insight into this matter is presented. We prepared 17-mer DNA duplexes incorporating Nile blue (NB) at one end as a redox marker and a disulfide tether at the other end for immobilization onto an electrode. The duplexes were covalently cross-linked by bifunctional cross-linkers that utilize either a propyl or naphthalene residue for replacing a base pair. Interestingly, the ICL-DNA with naphthalene residues exhibited DNA-mediated electron transfer, while the standard DNA and ICL-DNA with propyl-linker showed direct electron transfer pathway.

2P123 Bacterial ribosomal RNA as a target for sequence-specific inhibition

Joanna Trylska¹, Sapna G. Thoduka¹, Zofia Dabrowska¹, Anna Gorska¹, Maciej Jasinski^{1,2}, Tomasz Witula¹ (¹University of Warsaw, Centre of New Technologies, ²University of Warsaw, MISMAP College)

Many antibiotics block protein synthesis in pathogenic bacteria by interacting with bacterial ribosomal RNA (rRNA) and inhibiting the ribosome. However, bacteria have developed resistance against known antibiotics so there is a need to look for new antibacterials. We design short modified oligonucleotides to target in a sequence-specific manner rRNA. We test different DNA analogues such as 2'O-methyl-RNA, peptide nucleic acid (PNA), and locked nucleic acid (LNA) which hybridize well with RNA and are resistant to cellular enzymes. PNA is also known to have good strand invasion properties. Next, we check the ability of these oligonucleotides to inhibit protein synthesis in vitro in a cell-free transcription/translation system.

2P124 DNA の粗視化モデルによる Ars インスレーターの運動性と機能性の解析

Analysis of the fluctuation and functionality of Ars-insulator by coarse-grained model of DNA

Keisuke Yamamoto, Sayuri Tatemoto, Naoaki Sakamoto, Akinori Awazu (*Department of Mathematical and Life Sciences, Hiroshima University*)

Appropriate gene expressions are achieved by proper actions of enhancers. Since transcriptional activation by enhancer is independent of orientation and distance from promoter, interference of regulatory actions between neighboring genes may cause inappropriate gene expressions. In order to block inappropriate enhancer-promoter communications, insulator sequences are inserted between such inappropriate pair of the enhancer promoter. In this study, we focus on the structure and dynamics of Ars-insulator in sea urchin to uncover the mechanism of such insulator activity. For this purpose, we construct a coarse-grained model of DNA involving Ars-insulator region and some mutants and perform the normal mode analysis to compare the molecular fluctuation and activity.

2P125 四重鎖形成可能な相補鎖 DNA を導入することによる四重鎖リボザイムのカリウムイオン濃度依存的な活性スイッチングの高効率化

Enhancement of a Quadruplex-ribozyme activity in response to K⁺: a quadruplex-forming complementary DNA enables accurate switching

Yudai Yamaoki^{1,2,3}, Tsukasa Mashima¹, Takashi Nagata^{1,2}, Masato Katahira^{1,2} (¹Institute of Advanced Energy, Kyoto University, ²Graduate School of Energy Science, Kyoto University, ³JSPS Research Fellow)

An RNA R11, r(GGA)₃GG, forms a quadruplex under high K⁺ conditions. We replaced a part of a hammerhead ribozyme by R11. Although an activity of a designed ribozyme was enhanced in response to K⁺ via quadruplex formation in R11, our ribozyme showed non-negligible residual activity even in the absence of K⁺. In this study, in order to repress this residual activity, we introduced a complementary DNA that also can form a quadruplex in the presence of K⁺, to our ribozyme. It drastically repressed the residual activity. As a result, an activity ratio with/without K⁺ has reached 16-fold. Because K⁺ concentration is low outside the cell and high inside the cell our ribozyme may have a potential as a therapeutic drug and as an advanced drug delivery system.

2P126 抗プリオンアプタマーの構造学的基盤とその活性
Structural basis of anti-prion aptamer and its activity

Tsukasa Mashima¹, Fumiko Nishikawa², Yuji O. Kamatari³, Masayuki Saimura¹, Takashi Nagata^{1,4}, Satoshi Nishikawa², Kazuo Kuwata⁵, Masato Katahira^{1,4} (¹Inst. of Adv. Energy, Kyoto Univ., ²AIST, ³Life Sci. Res. Center, Gifu Univ., ⁴Grad. Sch. of Energy Sci., Kyoto Univ., ⁵Unit. Grad. Sch. of Drug Disc. and Med. Info. Sci., Gifu Univ.)

We have obtained an RNA aptamer as to prion protein (PrP), r(GGAGGAGGAGGA) (R12). R12 tightly binds to PrP with the dissociation constant of 10⁻⁸ M. We identified two separated lysine rich clusters, designated as P1 and P16, within bovine PrP as binding sites for R12. Here, we have revealed that in the R12:PrP complex, one protomer of the R12 dimer interacts with P1 and the other with P16, simultaneously. This simultaneous double binding renders the high affinity to R12 against PrP.

We also demonstrate the anti-prion activity of R12 using mouse neuronal cells persistently infected with a human TSE agent, which constitutively expressed PrP^{Sc}. After incubation with R12, the accumulation of PrP^{Sc} in the cell was explicitly reduced.