3P128 蛋白質およびリガンドの水和熱力学量計算に向けた形態計測法的アプローチ**A Morphometric Approach for the Accurate Solvation Thermodynamics of Proteins and Ligands**

Yuichi Harano¹, Roland Roth², Shuntaro Chiba³ (¹*IPR, Osaka Univ.*, ²*Inst. Theo. Phys. Tuebingen Univ.*, ³*Bio Center, TITECH*)

We have developed a versatile method for calculating solvation thermodynamic quantities for molecules, starting from their atomic coordinates. The contribution of each atom to the thermodynamic quantities is estimated as a linear combination of four fundamental geometric measures of the atomic species, which are defined by Hadwiger's theorem, and the coefficients reflecting their solvation properties. This treatment enables us to calculate the solvation free energy with high accuracy despite of the limited computational load. The method can readily be applied to macromolecules in an all-atom molecular model, allowing the stability of these molecules' structures in solution to be evaluated.

3P129 溶質-溶媒間のレナードジョンスポテンシャルパラメタが溶媒とダイナミクスに及ぼす影響**Effects of Lennard-Jones potential parameters between the solute and solvent on the solvation dynamics**

Yoshito Kondo, Tetsuro Nagai, Takuya Takahashi (*College of Life Sciences, Ritsumeikan Univ*)

It has been reported that mobility-increased water molecules exist around structure-breaker solutes experimentally. They have larger translational self-diffusion coefficients than a water molecule in bulk has.

In this study, we seek for a set of suitable Lennard-Jones potential parameters of interactions between solvent and solute based on TIP5P water model to more quantitatively reproduce the dynamical behavior of water molecules. In addition, we analyze the radial distribution functions and the number of hydrogen bonds with a number of combinations of Lennard-Jones parameter values. We then investigate their correlations to the dynamical behavior to study how the water dynamical behavior is governed.

3P130 MD および QM 計算による水和水のダイナミクスと電荷計算**MD and QM calculations of dynamics and charges of hydration water**

Takuya Takahashi (*College of Life Sciences, Ritsumeikan University*)

Classical MD simulations of water molecules around several types of solutes (poly-phosphates, AMP, and monovalent ions) were done with modified TIP5P water model. The hydration water molecules around mono-phosphate and di-phosphate molecules with a moderate negative charge showed increased mobility compared with bulk water and the results were partly consistent with the experimental results. Then, QM calculations were done with Gaussian software. In case of the tri-phosphate molecules, the calculated absolute charge of water oxygen atom was smaller as the solute charge became negative and the tendency was not changed in several calculation conditions such as different system size, basis function and chemical model.

3P131 カタウレイボヤ未受精卵に一過的低張刺激を与えると、幼生での左右非対称性が乱れる**Brief hypo-osmotic treatment on eggs disrupts the left-right asymmetry of the larvae in *Ciona intestinalis***

Shimpei Katsumoto, Kohei Hatta, **Masashi Nakagawa** (*Grad. Sch. Sci., Univ. Hyogo*)

Ascidian *Ciona intestinalis* tadpole larvae exhibit left-right asymmetry. Neurula embryos transiently rotate within the chorion. In tailbud embryos, the *Ci-pitx* gene is expressed in the left-side epidermis. The photoreceptors are situated only on the right side. We found that a brief hypo-osmotic treatment before, but not after, the neurula stage resulted in bilateral expression of *Ci-pitx* in tailbud embryos and the randomization of the location of photoreceptors in larvae. This treatment impaired the transient counterclockwise rotation within the chorion at the neurula stage. These results suggest that dead test cells blocked the neural rotation and impaired left-right asymmetry.

3P132 How To Achieve Sequential Local Folding of Epithelial Tube in Epididymis Development: Experimental and Mathematical Study

Tsuyoshi Hirashima, Ryoichiro Kageyama (*Inst. for Virus Research, Kyoto Univ.*)

In the murine epididymis development an epithelial tube of epididymis shows sequential folding in its head region locally although it remains unclear what is required for the morphogenesis of epididymal tube in the cellular level. Our mathematical model, in which the tube is represented by a bead-spring chain, predicts two sufficient conditions for the sequential local folding: local mitosis in the head region and high viscosity of a tissue surrounding in the tube. We confirmed experimentally that there are more mitotic cells in the head region than in the tail region and myofibroblasts that can produce the viscosity surround in a periphery of the tube. Furthermore, some perturbation experiments such as a transfection or an inhibitor assay verify the model prediction.

3P133 アフリカツメガエル卵母細胞における全 ATP 量の測定と ATP ライブイメージング**ATP quantification and live-imaging in *Xenopus laevis* oocyte**

Takashi W. Ijiri¹, Jun-ichi Kishikawa¹, Hiromi Imamura², Maho Sakiie³, Shuichi Ueno³, Yasuhiro Iwao³, Ken Yokoyama¹, Ken-ichi Sato¹ (¹*Fac. Life Sci., Kyoto Sangyo Univ.*, ²*Hakubi Center, Kyoto Univ.*, ³*Grad. Sch. Med., Yamaguchi Univ.*)

ATP plays various physiological roles; however, ATP metabolism in *Xenopus* oocyte has not been studied fully yet. First, we quantified global ATP in an oocyte using luciferase reaction-based reagent. The result did not show significant difference between immature and mature oocytes. Next, we set up an ATP live-imaging system using FRET-based indicator, ATeam. To observe its fluorescence, we prepared translucent immature oocytes that were injected with ATeam protein. Under fluorescence microscopy, the oocytes displayed strong FRET without background. Importantly, the FRET consistently increased in response to the addition of ATP, suggesting that ATeam works in the translucent oocytes. Local ATP distribution during oocyte maturation is currently under investigation.