3P164 F1-ATPase β サブユニットの全原子溶媒和自由エネルギー解析

All-atom hydration analysis of the β subunit in F1-ATPase

Toru Ekimoto¹, Mitsunori Ikeguchi¹, Nobuyuki Matubayashi² (¹*Yokohama City University*, ²*Kyoto University*)

F1-ATPase is a protein complex to hydrolyze adenosine tri-phosphate (ATP). Within the complex, the part called β subunit carries the catalytic activity. In this study, we focus on the ATP binding of the β subunit and treat the β subunit isolated in solvent water. To separately treat the water and ATP effects on the binding, we conceptually decompose the binding process into two steps. The first step is the conformational change of the β subunit from the open to closed states. The second is the ATP binding to the closed state. The effect of water is taken into account at all-atom level using the molecular dynamics simulation coupled with the method of energy representation. We will show that the solvent water promotes the first step and inhibits the second.

3P165 腸球菌由来 V_1 ATPase の軸強制回転シミュレーションによる回転機構の解明

Rotation mechanism of V_1 -ATPase studied by steered MD simulations

Yuta Isaka¹, Ichiro Yamato², Takeshi Murata^{3,4}, Mitsunori Ikeguchi¹ (¹*Grad. Sch. of Med. Life Sci., Yokohama City Univ.*, ²*Dept. Biol. Sci. Tech., Tokyo Univ. of Science*, ³*Fac. of Sci., Chiba Univ.*, ⁴*JST, PRESTO*)

V-ATPase is a rotary molecular motor functioning as a proton pump in cell membranes. A soluble part of V-ATPase is called V_1 , mainly composed of the A3B3 hexamer ring and the central stalk, D and F subunits. Recently, the high-resolution X-ray crystal structure of *Enterococcus hirae* V_1 ATPase was determined. Because the crystal structure is a snapshot of the dynamic processes in its functions, molecular dynamics (MD) simulations were required for elucidating the rotation mechanism. In this study, we performed the steered MD simulations, in which the central stalk was forced to be rotated. Conformational changes in the A3B3 hexamer ring in response to the stalk rotation were observed during the MD simulations.

3P166 Free energy simulations for the conformational change of the $\alpha\beta$ subunits in F1-ATPase after the ATP hydrolysis

Yuko Ito, Mitsunori Ikeguchi (*Yokohama City Univ.*)

F1-ATPase is a rotary motor enzyme. A single molecule experiment has identified that after the ATP hydrolysis, one of the β subunits in the F1-ATPase changes the conformation from the closed form to the half closed form. We performed ensemble sampling simulations to elucidate this structural change mechanism. The simulations were carried out with different nucleotide states. The obtained energy profiles agree well with the facts observed in single molecule experiments. Also, in the structural change from the open to closed form, we could obtain the half closed form (β_{HC}) in a minimum of the free energy surface.

3P167 回転モーター F1-ATPase の化学状態ごとのポテンシャルエネルギー

The potential energy of the rotary motor F1-ATPase for given chemical states

Kengo Adachi¹, Taisaku Ogawa¹, Kazuhiro Oiwa², Masasuke Yoshida³, Kazuhiko Kinoshita, Jr.¹ (¹*Waseda Univ.*, ²*Adv. ICT Res. Inst., NICT*, ³*Kyoto Sangyo Univ.*)

F1-ATPase is an ATP-driven rotary molecular motor that synthesizes ATP when forcibly rotated in reverse. The remaining challenge is to determine the potential energy that governs the mechanical conformational change in the motor. To measure the potential energy for rotation for given chemical states in the three catalytic sites, we controlled the rotation with magnetic tweezers by attaching a magnetic bead(s) to the central rotor, and simultaneously imaged the binding of fluorescently (Cy3) labeled ADPs and ATPs with TIRF microscopy. The torque was measured by the bead(s) displacement against magnets for the binding events of Cy3-nucleotides. We will report the individual potential which is estimated from integrating the torque for each chemical state.

3P168 F_0F_1 -ATP 合成酵素による ATP 駆動のプロトンポンプ活性の定量測定

Quantitative assay of ATP-driven proton-pump activity of F_0F_1

Ken Tasaki¹, Yuzo Kasuya¹, Naoki Soga¹, Toshiharu Suzuki², Masasuke Yoshida², Kazuhiko Kinoshita Jr.¹ (¹*Dept. Phys., Fac. Sci. Eng., Waseda Univ.*, ²*Dept. Mol. Bio., Fac. Life Sci., Kyoto Sangyo Univ.*)

The F_0F_1 -ATP synthase (F_0F_1) synthesizes ATP when proton motive force drives protons through F_0 . As a reverse reaction, F_0F_1 can pump protons in the opposite direction by ATP-hydrolysis. Although ATPase/synthesis functions have been deeply investigated, the proton-pump activity has been poorly characterized, qualitative even in bulk assays. Here, we report quantitative estimation of the proton-pump activity of F_0F_1 of thermophilic origin reconstituted into liposomes. The activity was monitored with a pH sensitive fluorescent probe pH-rodo conjugated to phospholipid, and the buffering capacity of the system was carefully estimated in several conditions. We also measured the ATP-hydrolysis activity under identical conditions.

3P169 ドメイン交換による V_0V_1 の MgADP 阻害機構の解明 Analysis of the MgADP-inhibition mechanism of V_0V_1 by domain swapping approach

Jun-ichi Kishikawa¹, Atsuko Nakanishi¹, Shou Furuie², Ken Yokoyama¹ (¹*Life Sci., Kyoto Sangyo Univ.*, ²*Dept. Phys., Osaka Med. College*)

V-type ATPase (V_0V_1) of *Thermus thermophilus* shows sensitivity to MgADP-inhibition, favorable for ATP synthesis reaction. Contrary, V_0V_1 from *Enterococcus hirae* and eukaryotes show insensitivity to MgADP-inhibition, favorable for continuous ATP hydrolysis. To clarify how these V_0V_1 show the different sensitivity to MgADP-inhibition, we produced domain swapping V_1 consisting of both *T. thermophilus* and *E. hirae* enzymes. Our results indicate that the domain-domain interaction in catalytic A subunit defines the binding affinity of V_1 for phosphate, which is critical for the sensitivity to MgADP-inhibition in V_1 . Based on recent reports of crystal structures of both V_1 , we discuss the molecular basis of MgADP-inhibition mechanism of V_0V_1 .