3P164 F₁-ATPase β サブユニットの全原子溶媒和自由エネルギー解析

All-atom hydration analysis of the β subunit in F₁-ATPase

Toru Ekimoto¹, Mitsunori Ikeguchi¹, Nobuyuki Matubayashi² (¹*Yokohama City University*, ²*Kyoto University*)

F₁-ATPase is a protein complex to hydrolyze adenosine tri-phosphate (ATP). Within the complex, the part called β subunit carries the catalytic activity. In this study, we focus on the ATP binding of the β subunit and treat the β subunit isolated in solvent water. To separately treat the water and ATP effects on the binding, we conceptually decompose the binding process into two steps. The first step is the conformational change of the β subunit from the open to closed states. The second is the ATP binding to the closed state. The effect of water is taken into account at all-atom level using the molecular dynamics simulation coupled with the method of energy representation. We will show that the solvent water promotes the first step and inhibits the second.

3P165 腸球菌由来 V₁ATPase の軸強制回転シミュレーションによる回転機構の解明

Rotation mechanism of V₁-ATPase studied by steered MD simulations

Yuta Isaka¹, Ichiro Yamato², Takeshi Murata^{3,4}, Mitsunori Ikeguchi¹ (¹*Grad. Sch. of Med. Life Sci., Yokohama City Univ.*, ²*Dept. Biol. Sci. Tech., Tokyo Univ. of Science*, ³*Fac. of Sci., Chiba Univ.*, ⁴*JST, PRESTO*)

V-ATPase is a rotary molecular motor functioning as a proton pump in cell membranes. A soluble part of V-ATPase is called V₁, mainly composed of the A3B3 hexamer ring and the central stalk, D and F subunits. Recently, the high-resolution X-ray crystal structure of *Enterococcus hirae* V₁ ATPase was determined. Because the crystal structure is a snapshot of the dynamic processes in its functions, molecular dynamics (MD) simulations were required for elucidating the rotation mechanism. In this study, we performed the steered MD simulations, in which the central stalk was forced to be rotated. Conformational changes in the A3B3 hexamer ring in response to the stalk rotation were observed during the MD simulations.

3P166 Free energy simulations for the conformational change of the $\alpha\beta$ subunits in F₁-ATPase after the ATP hydrolysis

Yuko Ito, Mitsunori Ikeguchi (*Yokohama City Univ.*)

F₁-ATPase is a rotary motor enzyme. A single molecule experiment has identified that after the ATP hydrolysis, one of the β subunits in the F₁-ATPase changes the conformation from the closed form to the half closed form. We performed ensemble sampling simulations to elucidate this structural change mechanism. The simulations were carried out with different nucleotide states. The obtained energy profiles agree well with the facts observed in single molecule experiments. Also, in the structural change from the open to closed form, we could obtain the half closed form (β_{HC}) in a minimum of the free energy surface.

3P167 回転モーター F₁-ATPase の化学状態ごとのポテンシャルエネルギー

The potential energy of the rotary motor F₁-ATPase for given chemical states

Kengo Adachi¹, Taisaku Ogawa¹, Kazuhiro Oiwa², Masasuke Yoshida³, Kazuhiko Kinoshita, Jr.¹ (¹*Waseda Univ.*, ²*Adv. ICT Res. Inst., NICT*, ³*Kyoto Sangyo Univ.*)

F₁-ATPase is an ATP-driven rotary molecular motor that synthesizes ATP when forcibly rotated in reverse. The remaining challenge is to determine the potential energy that governs the mechanical conformational change in the motor. To measure the potential energy for rotation for given chemical states in the three catalytic sites, we controlled the rotation with magnetic tweezers by attaching a magnetic bead(s) to the central rotor, and simultaneously imaged the binding of fluorescently (Cy3) labeled ADPs and ATPs with TIRF microscopy. The torque was measured by the bead(s) displacement against magnets for the binding events of Cy3-nucleotides. We will report the individual potential which is estimated from integrating the torque for each chemical state.

3P168 F₀F₁-ATP 合成酵素による ATP 駆動のプロトンポンプ活性の定量測定

Quantitative assay of ATP-driven proton-pump activity of F₀F₁

Ken Tasaki¹, Yuzo Kasuya¹, Naoki Soga¹, Toshiharu Suzuki², Masasuke Yoshida², Kazuhiko Kinoshita Jr.¹ (¹*Dept. Phys., Fac. Sci. Eng., Waseda Univ.*, ²*Dept. Mol. Bio., Fac. Life Sci., Kyoto Sangyo Univ.*)

The F₀F₁-ATP synthase (F₀F₁) synthesizes ATP when proton motive force drives protons through F₀. As a reverse reaction, F₀F₁ can pump protons in the opposite direction by ATP-hydrolysis. Although ATPase/synthesis functions have been deeply investigated, the proton-pump activity has been poorly characterized, qualitative even in bulk assays. Here, we report quantitative estimation of the proton-pump activity of F₀F₁ of thermophilic origin reconstituted into liposomes. The activity was monitored with a pH sensitive fluorescent probe pH-rodo conjugated to phospholipid, and the buffering capacity of the system was carefully estimated in several conditions. We also measured the ATP-hydrolysis activity under identical conditions.

3P169 ドメイン交換による V₀V₁ の MgADP 阻害機構の解明 Analysis of the MgADP-inhibition mechanism of V₀V₁ by domain swapping approach

Jun-ichi Kishikawa¹, Atsuko Nakanishi¹, Shou Furuike², Ken Yokoyama¹ (¹*Life Sci., Kyoto Sangyo Univ.*, ²*Dept. Phys., Osaka Med. College*)

V-type ATPase (V₀V₁) of *Thermus thermophilus* shows sensitivity to MgADP-inhibition, favorable for ATP synthesis reaction. Contrary, V₀V₁ from *Enterococcus hirae* and eukaryotes show insensitivity to MgADP-inhibition, favorable for continuous ATP hydrolysis. To clarify how these V₀V₁ show the different sensitivity to MgADP-inhibition, we produced domain swapping V₁ consisting of both *T. thermophilus* and *E. hirae* enzymes. Our results indicate that the domain-domain interaction in catalytic A subunit defines the binding affinity of V₁ for phosphate, which is critical for the sensitivity to MgADP-inhibition in V₁. Based on recent reports of crystal structures of both V₁, we discuss the molecular basis of MgADP-inhibition mechanism of V₀V₁.