1P115 ヒト cytochrome b561 form 3 (hb561-3)タンパク質の機能解析
Analyses of the physiological function of hb561-3: the
cytochrome b561 form 3 protein in human

Yuma Takahashi, Takako Yamazoe, Akikazu Asada, Motonari Tsubaki
 (Grad. Sch. Sci., Univ. Kobe)

Cytochrome b561 is a membrane protein having a role in regeneration of vesicular ascorbate (AsA) by transmembrane electron transfer. However, the physiological function of hb561-3 is not clarified. In the present study, we investigated the intercellular localization of hb561-3 protein in cultured human A549 cells and its interaction in the purified state with AsA. We found that hb561-3 protein is expressed in endoplasmic reticulum (ER), as found previously for its very close homolog, 101F6 protein. Upon interaction with AsA, oxidized heme of hb561-3 protein was reduced, but with a much slower rate than that of 101F6 protein. These results suggested that hb561-3 protein might interact with a distinct substrate other than AsA as the physiological electron donor.

1P116 常磁性緩和促進を利用した転写因子 FMBP-1 の分子ダイナミクスの解析
Molecular dynamics of transcription factor FMBP-1 proved by
paramagnetic relaxation enhancement

Kosuke Yuhara¹, Masakatsu Kamiya¹, Yasuhiro Kumaki¹, Shigeharu Takiya¹, Takashi Kikukawa¹, Keiichi Kawano^{1,2}, Makoto Demura¹, Tomoyasu Aizawa¹ (¹Grad. Sch. Life. Sci., Hokkaido Univ., ²Chitose Inst. Sci. Tech.)

FMBP-1 is a transcription factor of silk protein, fibroin H chain. FMBP-1 has a STPR domain, which is a novel DNA binding motif found in wide spectrum of eukaryotes. CD and other spectroscopic experiments showed that STPR domain is partially folded and has a flexible structure in the DNA-free state and DNA binding induces a conformational change into a more rigid structure. However, the detail of its binding process remains almost unknown due to the lack of information at the atomic-level about structure and dynamics of STPR domain. Here, we used paramagnetic relaxation enhancement (PRE) to characterize the dynamical aspect of DNA binding of FMBP-1. We measured the intramolecular contacts of STPR domain from PRE experiments and estimated the conformational dynamics.

1P117 MBD4 蛋白質のメチル化 CpG 結合ドメインと塩基除去活性ドメインをつなぐリンカー領域の構造と機能
Structural analysis of a linker region between methyl CpG
binding and glycosylase domains in MBD4

Itaru Takeshita (Grad. Sch. Eng., Kyoto Univ.)

Methyl CpG Binding Domain Protein 4 (MBD4) is a DNA repair enzyme that recognizes and excises a mismatch thymine base in methylated CpG sequence. MBD4 has two functional domains, the N-terminal methyl CpG binding domain (MBD) and the C-terminal glycosylase domain (GD), separated by a 265 amino acid linker. The crystal structures of these domains have been already solved. However, it remains unclear how these two domains act together in MBD4 function. In this study, structural properties of full length MBD4 were analyzed using SAXS combined with biochemical techniques. Our data suggest that MBD and GD are spatially coordinated relatively close to each other despite the long linker region. The linker region is implied to facilitate coordinated action of MBD and GD.

1P118 新奇 DNA 結合ドメイン STPR を持つ転写因子 FMBP-1 の in situ コンディションにおける DNA 認識動態の FCS 解析
FCS analysis of DNA recognition movements of transcription
factor FMBP-1 contains a novel DNA binding domain STPR in
situ condition

Motosuke Tsutsumi¹, Hideki Muto¹, Mai Kimoto¹, Masakatsu Kamiya¹, Takashi Kikukawa¹, Shigeharu Takiya¹, Makoto Demura¹, Keiichi Kawano^{1,2}, Masataka Kinjo¹, Tomoyasu Aizawa¹ (¹Grad. Sch. of Life Sci., Hokkaido Univ., ²Chitose Inst. Sci. Tech.)

Fibroin modulator binding protein-1 (FMBP-1) is a transcription factor of the fibroin H chain gene in the silkworm *Bombyx mori*. FMBP-1 has a unique tandem repeat DNA binding domain "the one score and three amino acid peptide repeat (STPR)". However, the functional details of STPR and molecular dynamics in live cell remain unknown. Here, we report an analysis of diffusion properties of FMBP-1 in posterior silk gland cells of *Bombyx mori* by fluorescence correlation spectroscopy (FCS). In our observation, EGFP fusion of FMBP-1 showed clear nuclear localization and the existence of movement with four distinct diffusion properties in a single measurement. This result might indicate co-existence of "facilitated diffusion" with specific binding on chromosomal DNA.

1P119 転写因子はいかにして障害物を迂回するか。分子シミュレーションによるアプローチ
How transcription factor can bypass obstacles? Molecular
simulation approaches

Mami Saito¹, Tsuyoshi Terakawa², Shoji Takada¹ (¹Grad. Sch.Sci., Univ. Kyoto, ²Univ. Columbia)

According to single molecular experiments, transcription factors search their target sites with sliding on DNA and 3D diffusion. These experiments mainly were under the conditions of liner DNA without obstacles or with fixed obstacles by antibodies. But in cells the obstacles are other transcription factors that can detach from DNA or histones wounded by DNA. Although the searching mechanism on molecular level in such conditions is still unclear, the transcription factor behavior in such condition is difficult to investigate with experiments. So we tried to reveal the searching mechanism with coarse-grained molecular dynamics software, Cafemol.

1P120 RecA 蛋白質の生体分子認識能力によるカーボンナノチューブ表面の DNA の生物学的機能の評価
Evaluation of biological function of DNA on the single-walled
carbon nanotubes by using biomolecular recognition ability of
RecA proteins

Shusuke Oura, Masahiro Ito, Yoshikazu Homma, Kazuo Umemura (Grad. Sch. Sci., Tokyo University of Science)

We examined the biological function of single-stranded DNA (ssDNA) molecules on the single-walled carbon nanotubes (SWNTs) by using biomolecular recognition ability of RecA proteins. In atomic force microscopy, the height distribution of observed rod-like hybrids showed that the ssDNA-SWNTs with RecA had larger height than the ssDNA-SWNTs. In agarose gel electrophoresis, the migration speed of the ssDNA-SWNTs was obviously decreased after adding RecA. For comparison, we performed similar tests for carboxymethylcellulose-SWNTs (CMC-SWNTs). Unlike the case of ssDNA-SWNTs, RecA didn't bind to CMC-SWNTs. Our results revealed that ssDNA molecules on the SWNT surfaces maintain their native function because RecA proteins recognized ssDNA on the SWNT surfaces precisely.