2P187 表皮細胞の傷の治癒には伸展によるヘミチャネルからの ATP 放出と TRPC6 を介した Ca^{2+} 流入が効く
Wound healing in keratinocyte is accelerated by mechanosensitive ATP release via hemichannels and Ca^{2+} influx through TRPC6 channels

Kishio Furuya^{1,2}, Hiroya Takada², Masahiro Sokabe^{1,2} (¹*Mechanobiology Labo, Grad Sch Med, Nagoya Univ.*, ²*Dept Physiol, Grad Sch Med, Nagoya Univ.*)

The cutaneous wound healing was accelerated by stretch and impaired in TRPC6-KO mice. HaCaT keratinocyte treated with hyperforin, a traditional herbal medicine and also TRPC6 activator, were cultured in an elastic chamber. At 3 h after making scratching, ATP release and intracellular Ca^{2+} response by stretch were determined by live-imaging. ATP release was observed only from foremost cells and it caused spreading Ca^{2+} waves. The Ca^{2+} response and wound healing were inhibited by apyrase, suramin, CBX and diC8- PIP_2 . In addition, hemichannel permeable dye calcein only entered ATP-releasing cells. These results suggested that stretch-accelerated wound closure was due to ATP release via hemichannels from the foremost cells and subsequent Ca^{2+} waves mediated by TRPC6.

2P188 Trial for detecting the activation and inactivation of chemoreceptor array in a single *E. coli* cell

Hajime Fukuoka¹, Tomoko Horigome², Yuichi Inoue¹, Hiroto Takahashi¹, Akihiko Ishijima¹ (¹*IMRAM, Tohoku Univ.*, ²*Grad. Sch. Life Sci., Tohoku Univ.*)

In chemotaxis system, the intracellular signal protein, CheY, is phosphorylated and dephosphorylated at the chemoreceptor array localized at cellular pole. Recently, we demonstrated multiple flagellar motors coordinately switch the rotational direction and such rotational switching was induced by binding and dissociation of CheY-P. Therefore, we speculated that the receptor array might produce pulses of CheY-P production by spontaneous activation of receptor array in steady-state of an *E. coli*. To investigate this possibility, we tried to simultaneously detect the activity of receptor array from the fluorescently labeled chemotaxis proteins and the rotational switching of a motor in a single *E. coli* cell. We would like to discuss our results in the annual meeting.

2P189 単離ヒドラ神経細胞の神経突起動態
Neurite dynamics of isolated hydra nerve cells

Rui Tanaka¹, Mihar Nagaishi², Mio Ogawa², **Hiroyuki Nakagawa**^{1,2} (¹*Div. Bio., Fac. Sci., Fukuoka Univ.*, ²*Grad. Sch. Sci., Fukuoka Univ.*)

Hydra, or cnidaria in general, is one of the first organisms with nerve cells evolving into the brain of higher bilaterians, such as insect and vertebrate. It is expected to be retained basic nature of nerve cell common to all nervous systems in them. However a few studies have been reported about their cellular characters as nerve cell, such as neurite dynamics. We, thus, isolated nerve cells from fresh water hydra, *Hydra magnipapillata*, and observed the dynamics of their neurites. Most of nerve cells well-attached to glass surface were observed as bipolar or multi-polar shape. During observation, neurites containing actin filament and microtubule were repeated the cycle of elongation and retraction at the rate of about 2 micrometer/sec.

2P190 サルモネラ菌の遊泳行動と生物対流
Chemotactic behavior of *Salmonella* and bioconvection

Takahiro Abe, Shuichi Nakamura, Seishi Kudo (*Grad. Sch. Eng., Tohoku Univ.*)

When cell density is low, chemotactic response of bacteria has been known as a biased-random walk. However, that at high cell density has not been fully understood. We analyzed chemotactic responses of *Salmonella* by a capillary assay under an optical microscope with changing cell density to characterize the chemotaxis in high cell density. Analysis by PIV (Particle Image Velocimetry) showed that large-scale convection flow was generated when bacteria accumulated to attractant. The direction and stability of convection were affected by the direction and effective magnitude of gravitational force. Random swimming of individual cells was observed even during the convection.

2P191 Rotation analysis of the spirochete cell body by 3D dark-field microscopy

Kyosuke Takabe, Md. Shafiqul Islam, Seishi Kudo, Shuichi Nakamura (*Grad. Sch. Engineering., Univ. Tohoku*)

Leptospira are spirochetes and have flagellar motors at both cell ends. When Leptospira swim, the anterior end is spiral-shaped (S) and the posterior end is hook-shaped (H). The S-end is known to rotate counterclockwise (CCW) to propel the cell. In contrast, the H-end has been believed to rotate CCW. Based on a theoretical analysis, we recently proposed a hypothesis that the H-end rotates clockwise (CW). However, direct observations of the H-end rotation to determine its rotational direction have not been performed. In this study, we examined the rotational direction of the H-end by three-dimensional dark-field microscopy. We found that most cells showed the CW rotations in the H-ends, which gives an evidence for the counterrotation of the H-end against the S-end.

2P192 細菌の走化性に起因する リング状パターンの形成過程の解析
Formation process of a ring-like pattern induced by bacterial chemotaxis

Tadahiko Sato, Shuichi Nakamura, Seishi Kudo (*Grad. Sch. Eng., Univ. Tohoku*)

Motile bacteria respond to the concentration gradient of chemicals and migrate toward the place more suitable for growth, which is called chemotaxis. It is known that, when an agar drop including an attractant is placed in a bacterial suspension, a ring-like pattern is formed around the drop. Although the ring formation is caused by positive chemotaxis, the formation mechanism is still unclear. In this study, we quantitatively analyzed the formation process of the ring. We found that the ring-formation speed and the ring size after reaching the steady state were increased with the concentration of attractants. We are now trying to observe the behavior of individual cells during the ring formation.