2P235 プロテオロドプシンのプロトンドナー Glu108 の変異が光化学特性に及ぼす影響について**Effects on the photochemical properties in proteorhodopsin by the mutation of the Glu108 residue**

Jun Tamogami¹, Takashi Kikukawa², Kimi Okubo¹, Makoto Demura², Toshifumi Nara¹, Naoki Kamo² (¹*College Pharm. Sci., Matsuyama Univ.*, ²*Fac. Adv. Life Sci., Hokkaido Univ.*)

Proteorhodopsin (PR) is one of the microbial rhodopsins, which acts as a light-driven proton pump. It has been considered that Glu108 in PR (corresponding to Asp96 in bacteriorhodopsin) functions as an internal proton donor of the Schiff base during its cyclic photochemical reaction (called photocycle). On the other hand, the corresponding residue in a novel PR-like protein found in *Exiguobacterium sibiricum* (ESR) was a lysine, however, ESR had the proton pumping activity over a wide pH range and fast turnover photocycle. This indicates that the existence of a carboxylic residue in the cytoplasmic domain is not necessary for proton pumping. In this study, we investigated the effects of the mutation of Glu108 on the photocycle and accompanying proton transfer in PR.

2P236 *Acetabularia* rhodopsin II (ARII) の Asp81 変異体による一過性光誘起電流**Transient photo-induced current by Asp81 mutants of *Acetabularia* rhodopsin II (ARII) heterologously expressed in *Xenopus laevis* oocytes**

Seiji Miyauchi¹, Kazumi Shiono¹, Takashi Kikukawa², Makoto Demura², Jung Kwang-Hwan³, Naoki Kamo² (¹*Toho Univ., Grad. Sch. Pharm. Sci.*, ²*Hokkaido Univ., Grad. Sch. Life Sci.*, ³*Sogang Univ., Inst. Biol. Interfaces*)

ARII is a light-driven proton pump rhodopsin. Asp81 in ARII corresponding to Asp85 in bacteriorhodopsin is considered to function as a proton acceptor from the protonated Schiff base.

We here determined how this proton acceptor coordinates with the protonated Schiff base and other amino acid residues to pump out a proton. The photo-induced current by D81S mutant exhibited transient; the light illumination induced the transient inward-current, whereas the turn-off of the light illumination induced the outward transient current. These transient currents remained unchanged irrespective of extracellular ions, Na⁺ and Cl⁻, implying that the transient currents might be due to an intracellular circulation of a certain ion.

2P237 Low-temperature FTIR spectroscopy of the Light-driven sodium ion pump: *Krokinobacter eikastus* rhodopsin 2

Faisal Hammad Mekky Koua¹, Rei Abe-Yoshizumi², Hikaru Ono², Shota Ito², Yoshitaka Kato², Keiichi Inoue^{1,2}, Hideki Kandori^{1,2} (¹*OptoBioTech. Res. Cent., Nagoya Inst. Tech.*, ²*Dept. Front. Mat., Nagoya Inst. Tech.*)

Krokinobacter eikastus rhodopsin 2 (KR2) is a newly discovered light-driven compatible sodium ion-proton pump^[1]. KR2 pumps Na⁺ and Li⁺, while turning into proton pump in the presence of K⁺ or larger cations. The Na⁺ ion pumping activity has been shown to be completely suppressed at acidic pH and for D116N, the retinal Schiff base counterion mutant. To understand the mechanism behind this, we applied low-temperature light-induced difference FTIR^[2] and UV-vis spectroscopy at a range of pH for the wild type KR2 and the D116N mutant. We discuss the possible mechanisms of the light-driven Na⁺ ion pumping based on the analysis for these proteins and other mutants.

[1] Inoue, K et al. *Nat. Commun.* 4, 1678-1687 (2013).

[2] Ono, H et al. *J. Phys. Chem. B* 118, 4784-4792 (2014).

2P238 ロドプシンの低い熱活性化頻度の分子メカニズム**Molecular mechanism of the low thermal activation rate of rhodopsin**

Keiichi Kojima¹, Masataka Yanagawa², Takahiro Yamashita¹, Yasushi Imamoto¹, Takesi Matsuyama Hoyos¹, Koji Nakanishi³, Yumiko Yamano⁴, Akimori Wada⁴, Yasushi Sako², Yoshinori Shichida¹ (¹*Grad. Sch. Sci., Kyoto Univ.*, ²*Cell. Info. Lab., Riken*, ³*Columbia Univ.*, ⁴*Kobe Pharm. Univ.*)

Most vertebrate retinas contain two types of photoreceptor cells, rods and cones, which are responsible for scotopic and photopic vision, respectively. Because rods show a low threshold of photon detection and high sensitivity, the low thermal activation rate of rhodopsin, rod visual pigment, is essential for the function of rods. Phylogenetic analysis indicates that rhodopsin has evolved out of cone visual pigment, but it remains unknown how rhodopsin acquired the low thermal activation rate in the molecular evolution. We estimated the thermal activation rates of rhodopsin, cone visual pigments and their mutants by biochemical and spectroscopic measurements. Based on our results, we discuss the molecular mechanism of the low thermal activation rate of rhodopsin.

2P239 ロドプシン構成的活性変異体 M257Y における構造平衡の分子観測**Single-molecule observation of the conformational equilibrium in constitutively active mutant of rhodopsin, M257Y**

Ryo Maeda¹, Michio Hiroshima^{1,3}, Yasushi Imamoto², Takahiro Yamashita², Yasushi Sako¹, Yoshinori Shichida² (¹*Cellular Informatics Lab., RIKEN*, ²*Grad. Sch. Biophys., Kyoto Univ.*, ³*QBiC, RIKEN*)

Rhodopsin, a G-protein coupled receptor, has retinal as a ligand and the ligand-free state opsin scarcely activates G-protein. Previous studies have shown that opsin state of constitutively active mutant (CAM) activates G-protein much effectively than that of wild type but the mechanism has been unclear. We labeled M257Y (a typical CAM) with Alexa594 fluorophore and monitored the conformational changes directly by single-molecule measurements using TIRFM. We compared conformational dynamics in the dark, photoactivated, and opsin states of M257Y and wild type rhodopsin, and found that conformational changes occurred more frequently in M257Y opsin than wild type opsin. The frequent generation of active state would account for the constitutive activity of opsin.

2P240 In situ 光照射固体 NMR による光受容膜タンパク質センサーロドプシン I の光反応過程の解析**Photocycle of sensory rhodopsin I as revealed by in situ photo irradiation solid-state NMR**

Yoshiteru Makino¹, Hiroki Yomoda¹, Yuya Tomonaga¹, Tetsuro Hidaka¹, Izuru Kawamura¹, Takashi Okitsu², Akimori Wada², Yuki Sudo³, Naoki Kamo⁴, Akira Naito¹ (¹*Grad. Sch. Eng, Yokohama Natl Univ.*, ²*Kobe Pharm. Univ.*, ³*Grad. Sch. Pharm, Okayama Univ.*, ⁴*Grad. Sch. Life Sci, Hokkaido Univ.*)

Salinibacter ruber Sensory rhodopsin I (SrSRI) is a photoreceptor membrane protein with a retinal chromophore, and has a dual function as positive and negative phototaxis. To characterize photo-intermediates in the photocycle of SrSRI, configuration change of [20-¹³C] retinal was distinctly observed by *in situ* photo irradiation solid state NMR [1]. During green and light irradiation, the retinal changed from all-*trans* to 13-*cis*, indicating that M-intermediate (attractant) was stationary trapped. M-intermediate changed to P-intermediate (repellent) with protonated Schiff base by illumination with UV-light. In conclusion, the dual function in SrSRI is induced by the color-discriminating isomerization of retinal. [1]Y. Makino et al.(2014) *Angew. Chem. Int. Ed. in press*