3P229 低温・Xe 加圧下における DEPC リポソーム相変化と神経細胞のラマン測定
Raman spectra change at the phase transition of DEPC liposome and cell membrane of neuron under low temperature and xenon pressure

Tsutomu Uchida¹, Masafumi Nagayama², Kazutoshi Gohara¹, Amadeu K. Sum³ (¹*Fac. Eng., Hokkaido Univ.*, ²*Hokkaido Univ. Edu., Asahikawa*, ³*Colorado Sch. Mines, USA*)

The change of spectra of DEPC liposome with temperature and xenon pressure was observed by microscopic Raman spectroscopy to identify the phase transition of liposome. At atmospheric pressure, DEPC liposome transforms from gel to liquid crystal at approximately 120°C and shifts to lower temperature at xenon pressure. The Raman spectra indicated the slight change of C-H stretching modes at the phase transition temperature under xenon pressure. These findings support the previously obtained results by MD simulations, which showed the evidence of xenon dissolved in the lipid bilayer. Based on these observations, the Raman spectra of neurons under low temperature and xenon pressure were observed.

3P230 局所熱パルス法による神経細胞のカルシウム放出の誘導
Ca²⁺-burst in rat hippocampal neurons induced by microscopic heat pulses

Yuki Kawamura¹, Kotaro Oyama¹, Hideki Itoh^{1,2}, Madoka Suzuki^{3,4}, Shin'ichi Ishiwata^{1,3,4} (¹*Sch. Adv. Sci. Eng., Waseda Univ., Tokyo, Japan*, ²*Inst. Med. Biol., A*STAR, Singapore*, ³*Org. Univ. Res. Initiatives, Waseda Univ., Tokyo, Japan*, ⁴*WABIOS, Singapore*)

Temperature is an essential parameter determining physical and chemical processes in cells. We previously reported highly thermosensitive Ca²⁺ dynamics in HeLa cells (V. Tseeb et al, 2009). In a search of novel methods controlling neural activities, we studied the thermoresponsive Ca²⁺ dynamics in neurons. We found that single rat hippocampal neurons responded to microscopic heat pulses induced by an infra-red laser beam focused nearby. The Ca²⁺ change in neurite was larger than that in soma. In addition, we achieved to build a molecular model for the thermoresponsive Ca²⁺ signaling, which is different from HeLa cells in function of ion channels. Our results demonstrate that microscopic heat pulses can potentially be useful for exciting neurons locally and noninvasively.

3P231 ヒト iPS 細胞由来ニューロンとアストロサイト共培養による電気生理学的特徴
Electrophysiological activity of a human induced pluripotent stem cell derived neuron and astrocyte co-culture

Aoi Odawara^{1,3}, Ikuro Suzuki² (¹*Department of Bionics, Tokyo University of Technology*, ²*Department of Electronics and Intelligent Systems, Tohoku Institute of Technology*, ³*Japan Society for the Promotion of Science*)

Human induced pluripotent stem cell (hiPSC)-derived neurons may be effectively used for drug discovery and cell-based therapy. However, the immaturity of cultured human iPSC-derived neurons and the lack of established functional evaluation methods are problematic. We here used a multi-electrode array (MEA) system to investigate the effects of the co-culture of rat astrocytes with hiPSC-derived neurons on the long-term culture, spontaneous firing activity, and drug responsiveness effects. The co-culture facilitated the long-term culture of hiPSC-derived neurons for >3 months and long-term spontaneous firing activity was also observed. As a result, The co-culturing with rat astrocytes allowed hiPSC-derived neurons to maintain long time spontaneous activity.

3P232 導電性高分子含浸ファイバー電極の脳活動測定と刺激への適用
Application of conductive polymer-coated fiber electrodes to neural recording and stimulation in vivo

Satoshi Watanabe, Hideyuki Takahashi, Keiichi Torimitsu (*Dept. Bioeng. Robotics, Grad. Sch. Eng., Tohoku Univ.*)

Electrodes for chronic implantation are desired to be made of soft material that induces minimal stress, and should have low impedance and biocompatibility for stable long-term functioning. We fabricated a fiber-shaped electrode by coating silk fibers with the conductive polymer, poly(3,4-ethylenedioxythiophene) p-toluenesulfonate (PEDOT-pTS), by chemical polymerization. Using this electrode, we recorded field potentials from embryonic chick brain with a signal-to-noise ratio comparable to conventional metal electrodes. The fiber electrode is expected to apply for clinical purposes such as deep brain stimulation in neuropathological diseases.

3P233 運動性シアノバクテリア *Pseudanabaena* sp. ILC 545 の「彗星状コロニー」の形成ダイナミクス
Dynamics of comet-like colony formation in the filamentous cyanobacterium, *Pseudanabaena* sp. ILC 545

Yu Shoji¹, Hiroki Yamamoto¹, Yuki Fukasawa¹, Hideo Iwasaki^{1,2} (¹*Waseda University*, ²*metaPhorest*)

We have analyzed colony-pattern formations in the multicellular (filamentous) cyanobacterium, *Pseudanabaena* sp. ILC 545, which we isolated previously. ILC 545 strain glides their major axis. As the bacterial density increases, we observed various colony patterns, due to collisions and entanglement. These colony patterns can be classified into three groups: “bundle” colonies with nematic alignment, “comet-like” colonies with fast population movement, and “disk” colonies forming vortices.

We focused on the comet-like colony formation. Concretely we analyzed the relationship between colony size and speed, and movement of one filament in the colony. Here, we report a model on the morphogenesis of comet-like colony formation, and some related quantitative analyses.

3P234 Analysis on colony formation in a filamentous cyanobacterium with an extended self-driven particle model with a cellular automaton method

Masato Ishii¹, Yuki Fukasawa¹, Masaya Takiguchi¹, Kain Yanagi¹, Hideo Iwasaki^{1,2} (¹*Waseda University*, ²*metaPhorest*)

Cyanobacteria, a group of eubacteria performing oxygenic photosynthesis, show a variety of patterns depending on environmental conditions such as light and nutrients. The motile cyanobacterium, *Geitlerinema* sp. ILC 546, forms a unique colony pattern, which is characterized by multiple rings on an agar medium with robust trails. We performed some quantitative analysis on the movement of gliding filaments, and employed a cellular automaton method to modify the self-driven particle (Vicsek) model for population dynamics of filamentous units. Some critical conditions which seem important for the robust-trail ring pattern formation will be reported.