Advance Publication

Method and Protocol

GA-guided mD-VcMD: A genetic-algorithm-guided method for multi-dimensional virtual-system coupled molecular dynamics

Junichi Higo¹, Ayumi Kusaka², Kota Kasahara³, Narutoshi Kamiya¹, Itaya Hayato⁴, Xie Qilin⁵, Takuya Takahashi³, Ikuo Fukuda¹, Kentaro Mori^{1,6}, Yutaka Hata¹ and Yoshifumi Fukunishi⁷

¹ Graduate School of Simulation Studies, University of Hyogo, Kobe, Hyogo 650-0047, Japan

² Institute for Protein Research, Osaka University, Suita, Osaka 565-0871, Japan

³ College of Life Sciences, Ritsumeikan University, Kusatsu, Shiga 525-8577, Japan

⁴ Graduate School of Life Sciences, Ritsumeikan University, Kusatsu, Shiga 525-8577, Japan

⁵ College of Pharmaceutical Sciences, Ritsumeikan University, Kusatsu, Shiga 525-8577, Japan

⁶ National Institute of Technology, Maizuru College, Kyoto 625-8511 Japan

⁷ Cellular and Molecular Biotechnology Research Institute, National Institute of Advanced Industrial Science and Technology (AIST), Koto-ku, Tokyo 135-0064, Japan

doi: 10.2142/biophysico.BSJ-2020008

Received: April 16, 2020

Accepted: December 22, 2020

Released online in J-STAGE as advance publication: December 26, 2020

(Edited by Motonori Ota)

Significance

We introduced a generalized ensemble method named “genetic-algorithm guided multi-dimensional virtual-system coupled molecular dynamics (GA-guided mD-VcMD)” to enhance conformational sampling of a biomolecular system. This method is free from fine tuning of a weight function, and a production run is not required. A thermodynamic weight is assigned to each snapshot by a simple procedure. Statistics of poorly sampled conformations are compensated by a selective sampling procedure. This method was applied to a simple system that involves an energy barrier between potential-energy minima, and demonstrated that GA-guided mD-VcMD has a considerably high sampling efficiency.

Category of Article: Methods and Protocols

**GA-guided mD-VcMD: A genetic-algorithm-guided method for
multi-dimensional virtual-system coupled molecular dynamics**

Junichi Higo¹, Ayumi Kusaka², Kota Kasahara³, Narutoshi Kamiya¹, Itaya Hayato⁴, Xie
Qilin⁵, Takuya Takahashi³, Ikuo Fukuda¹, Kentaro Mori^{1,6}, Yutaka Hata¹, and
Yoshifumi Fukunishi⁷

¹ *Graduate School of Simulation Studies, University of Hyogo, 7-1-28 Minatojima
Minami-machi, Chuo-ku, Kobe, Hyogo 650-0047, Japan*

² *Institute for Protein Research, Osaka University, 3-2 Yamada-oka, Suita, Osaka
565-0871, Japan*

³ *College of Life Sciences, Ritsumeikan University, 1-1-1 Noji-higashi, Kusatsu, Shiga
525-8577, Japan*

⁴ *Graduate School of Life Sciences, Ritsumeikan University, 1-1-1 Noji-higashi, Kusatsu,
Shiga 525-8577, Japan*

⁵ *College of Pharmaceutical Sciences, Ritsumeikan University, 1-1-1 Noji-higashi,*

Kusatsu, Shiga 525-8577, Japan

⁶ *National Institute of Technology, Maizuru College, 234 Shiroya, Maizuru, Kyoto,*

625-8511 Japan

⁷ *Cellular and Molecular Biotechnology Research Institute, National Institute of*

Advanced Industrial Science and Technology (AIST), 2-3-26, Aomi, Koto-ku, Tokyo,

135-0064, Japan

Corresponding author: Junichi Higo. Graduate School of Simulation Studies, University

of Hyogo, 7-1-28 Minatojima Minami-machi, Chuo-ku, Kobe, Hyogo 650-0047, Japan

e-mail: higo@protein.osaka-u.ac.jp

Running title: Genetic-algorithm-guided enhanced sampling method

Abstract

We introduced a conformational sampling method in an earlier report. The multi-dimensional virtual-system coupled molecular dynamics (mD-VcMD) enhances conformational sampling of a biomolecular system by computer simulations. Herein, new sampling method, a subzone-based mD-VcMD, is presented as an extension of mD-VcMD. Then, the subzone-based method is extended further using a genetic algorithm (GA) named the GA-guided mD-VcMD. In these methods, iterative simulation runs are performed to increase the sampled region gradually. The new methods have the following benefits: (1) They are free from a production run: i.e., all snapshots from all iterations are useful for analyses. (2) They are free from fine tuning of a weight function (probability distribution function or potential of mean force). (3) A canonical ensemble (i.e., a thermally equilibrated ensemble) is generated from a simple procedure. A thermodynamic weight is assigned to each snapshot. (4) Selective sampling can be performed for particularly addressing a poorly sampled region without breaking the proportion of the canonical ensemble if the poorly sampled conformational region emerges in sampling. By applying the methods to a simple system that involves

an energy barrier between potential-energy minima, we demonstrated that the new methods have considerably higher sampling efficiency than the original mD-VcMD does.

Key words: Canonical ensemble, Computer simulation, Conformational sampling, Enhanced sampling, Generalized ensemble

Introduction

Generalized ensemble methods are useful for effectively searching a wide conformational space of a molecular system [1,2]. These methods enhance sampling along a reaction coordinate (RC) such as energy or a structural parameter (e.g., a root-mean-square difference from a reference conformation). Moreover, they are powerful for the exploration of considerably stable states (major basins) in the conformational space. However, these methods might overlook less-stable states (minor basins): One can assume that a major basin and a minor basin, the positions of which are separated in the full-dimensional conformational space, mutually overlap in the RC axis. As a result, conformational trapping might occur. The minor basins might be overlooked [3,4]. This oversight does not lead to a severe difficulty when the minor basin is out of scope. However, an alternative approach must be used when the minor basin acts as bridges among the major basins or when it has key features for exerting biophysical functions of the system.

Adaptive umbrella sampling [5,6] can avoid such an oversight or trapping problem because one can set the RC such that the major and minor basins are

discriminated along the RC axis. However, this method requires fine tuning of a weight function (potential of mean force (PMF)) along the RC axis. Practically speaking, the difficulty of the fine tuning might result in a very long simulation (or increment of the number of iterative simulations) to sample the conformational space widely.

To avoid these difficulties, we introduced a multi-dimensional virtual-system coupled molecular dynamics (mD-VcMD) simulation [7,8], which is designated herein as the original mD-VcMD (or simply the original method). This method enhances conformational sampling with repetition of iterative simulations in a multi-dimensional RC space. Furthermore, this method is free from fine tuning of the weight function because the weight function does not appear in the proceedings of the iterative simulations. We applied the original mD-VcMD method to a large and complicated system consisting of a ligand (endothelin-1) and its receptor (human endothelin type B receptor; one of the GPCR proteins), where the receptor was wrapped by cholesterol, embedded in an explicit membrane, and surrounded by an explicit solvent [8].

We encountered a difficulty in the original-mD-VcMD study: Once a poorly sampled region emerged in the multi-dimensional RC space in an iterative simulation,

this region might cause conformational trapping in a subsequent iteration. Consequently, a more robust method is necessary to advance the iteration.

For this purpose, in the present study, we introduce two methods with extension of the original mD-VcMD: a subzone-based mD-VcMD method, which is an extension of the original mD-VcMD; and a GA-guided mD-VcMD method, which extends the subzone-based mD-VcMD to expand sampling to non-sampled RC regions using genetic algorithm (GA). As presented in this paper, we explain the GA-guided and original mD-VcMD methods. We apply the methods to a simple model with a potential-energy barrier between potential-energy minima. Because of the simplicity of this model, the difficulty of the sampling task is determined by the effective barrier height between the minima, which is modulated readily by changing the simulation temperature. At a high temperature, both the original and GA-guided mD-VcMD methods showed quick convergence of the numerical distribution to the analytical distribution of the system. By contrast, at a lower temperature, only the GA-guided method exhibited quick convergence.

Methods and Protocols

We introduce many terms and quantities in this paper and its related Supportive Information (SI). Supplementary Table S1 presents positions for which the terms and quantities are defined or explained. Sections 1–4 of SI are preparatory sections for the original mD-VcMD method. Section 5 of SI explains the original mD-VcMD method itself [7,8]. The original method produces the statistical weight $Q_{cano}(\mathbf{L})$ assigned to zones $\zeta_{\mathbf{L}}$ (virtual zones) introduced into the RC space (precise definition of $Q_{cano}(\mathbf{L})$ is given in SI). Integration of $Q_{cano}(\mathbf{L})$ over all zones provides the probability distribution of the system in the entire RC space. The subzone-based and GA-guided mD-VcMD methods are extensions of the original method to compute $Q_{cano}(\mathbf{L})$.

The subzone-based and GA-guided methods are designed to treat a multiple-RC space, just as the original mD-VcMD method. In fact, the original method was applied to a complicated system using a seven-dimensional RC space [8]. However, to simplify the explanation, we use a one-dimensional (1D), two-dimensional (2D), or

three-dimensional (3D) RC space in this paper. One can increase the space dimensionality straightforwardly.

Subzone-based mD-VcMD Method to Compute $Q_{cano}(\mathbf{L})$

Introduction of subzone and subzone's snapshot count (SS count)

Here, we present the subzone-based method to compute $Q_{cano}(\mathbf{L})$. The sections below are based on sections 1--5 of SI. Figure 1A, which is substantially as shown in Supplementary Fig. S2C, is a portion of a 2D RC space, where four linked zones shown by differently colored frames overlap in an intersection (shaded region). The arrows indicate directions to shift the linked zones to remove the inter-zone overlaps. Figure 1B is an extended RC space resulted from the zone shifting, where the virtual states (indices assigned to nine virtual zones) are numbered as $\mathbf{L}_1, \dots, \mathbf{L}_9$ with arbitrary order. Subscripts i for $\mathbf{L}_i$ are not those to specify linked virtual states (section 3 of SI). The linked virtual states are expressed with parentheses as $\mathbf{L}_{(i)}$. The extended RC space is introduced for convenience to make explanation simple. The zone-overlap exists as in Fig. 1A in the actual mD-VcMD simulation.

Here we introduce *subzones*, which overlap perfectly on intersections in the original RC space. The shaded region in Fig. 1A corresponds to four shaded regions in Fig. 1B. The subzones are these shaded regions in Fig. 1B. Similarly, four checked subzones in Fig. 1B overlap on another intersection (not presented in Fig. 1A). Once the subzones are introduced, any zone is constructed by subzones. For instance, in Fig. 1B, each zone $\mathbf{L}_i$ consists of four subzones with indices expressed as $\mathbf{S}_j$ ($j = 1, \dots, 4$). In an m -dimensional RC space, a zone comprises 2^m subzones in general. Numbering of the subzones is arbitrary. Whereas $\mathbf{L}$ specifies the zone position in the whole RC space, $\mathbf{S}$ specifies the relative position of a subzone in each zone.

We express a subzone $\mathbf{S}$ in zone $\mathbf{L}$ as $[\mathbf{S}, \mathbf{L}]$. We denote the number of snapshots detected in the subzone $\mathbf{S}$ during the iteration M by $c^{[M]}(\mathbf{S}, \mathbf{L})$, and normalize it by the number of snapshots detected in the zone $\mathbf{L}$ as

$$\frac{c^{[M]}(\mathbf{S}, \mathbf{L})}{\sum_{\mathbf{S}' \in \mathbf{L}} c^{[M]}(\mathbf{S}', \mathbf{L})} \rightarrow c^{[M]}(\mathbf{S}, \mathbf{L}). \quad (1)$$

This normalization results in $\sum_{\mathbf{S} \in \mathbf{L}} c^{[M]}(\mathbf{S}, \mathbf{L}) = 1$. We refer to this quantity $c^{[M]}(\mathbf{S}, \mathbf{L})$ as *subzone's snapshot count (SS count)* at $[\mathbf{S}, \mathbf{L}]$. The SS count for a non-sampled subzone in iteration M is zero: $c^{[M]}(\mathbf{S}, \mathbf{L}) = 0$.

Here, we characterize zones as three types: This zone is called an *empty zone* (*E zone*) if all the subzones in zone $\zeta_{\mathbf{L}}$ are not sampled ($c^{[M]}(\mathbf{S}, \mathbf{L}) = 0, \forall \mathbf{S}$ in $\mathbf{L}$). This zone is designated as a *completely sampled zone* (*CS zone*) if all subzones in $\zeta_{\mathbf{L}}$ are sampled ($c^{[M]}(\mathbf{S}, \mathbf{L}) > 0, \forall \mathbf{S}$ in $\mathbf{L}$). Finally, if some subzone is empty and the others are sampled in $\zeta_{\mathbf{L}}$, then this zone is called an *incompletely sampled zone* (*IS zone*). Figure 2 presents those zones assuming that the first, second, and third iterations are performed sequentially using a 2D RC space.

Next, we introduce two delta functions δ and D as

$$\delta^{[M]}(\mathbf{S}, \mathbf{L}) = \begin{cases} 1 & \text{(if } c^{[M]}(\mathbf{S}, \mathbf{L}) > 0 \text{ in iteration } M) \\ 0 & \text{(if } c^{[M]}(\mathbf{S}, \mathbf{L}) = 0) \end{cases} \quad (2)$$

and

$$D^{[M]}(\mathbf{L}) = \begin{cases} 1 & \text{(if } \boldsymbol{\zeta}_L \text{ is CS zone in iteration } M) \\ 0 & \text{(otherwise)} \end{cases}. \quad (3)$$

Then, we average the SS count $c^{[I]}(\mathbf{S}, \mathbf{L})$ ($I = 1, \dots, M$) at each $[\mathbf{S}, \mathbf{L}]$ over the first to M -th iterations with the following procedures.

Procedure 1: If $\boldsymbol{\zeta}_L$ is a CS zone in one or more iterations (Fig. 2A), then the average of SS count is defined as

$$c^{[1;M]}(\mathbf{S}, \mathbf{L}) = \frac{\sum_{I=1}^M D^{[I]}(\mathbf{L}) c^{[I]}(\mathbf{S}, \mathbf{L})}{\sum_{I=1}^M D^{[I]}(\mathbf{L})}. \quad (4)$$

In Eq. 4, IS and E zones are eliminated from the averaged SS counts. Superscript $[1;M]$ denotes that the SS counts are computed using an iteration window from the first to M -th iterations.

Procedure 2: If $\boldsymbol{\zeta}_L$ is an IS or E zone in all iterations and a CS zone is not involved, then average is calculated as

$$c^{[1;M]}(\mathbf{S}, \mathbf{L}) = \frac{\sum_{I=1}^M c^{[I]}(\mathbf{S}, \mathbf{L})}{\sum_{I=1}^M \delta^{[I]}(\mathbf{L}, \mathbf{S})}. \quad (5)$$

The resultant SS counts do not vary when $c^{[I]}(\mathbf{S}, \mathbf{L})$ is replaced by $\delta^{[I]}(\mathbf{S}, \mathbf{L})c^{[I]}(\mathbf{S}, \mathbf{L})$ in Eq. 5. A CS zone can result from combining IS zones, as presented in Fig. 2B.

By contrast, as shown in Fig. 2C, some non-sampled subzones (say $[\mathbf{S}', \mathbf{L}]$) might remain after averaging, which results in an IS zone for the zone and $c^{[1;M]}(\mathbf{S}', \mathbf{L}) = 0$. Then, we correct the zero SS counts as follows: First, non-zero SS counts are averaged in ζ_L :

$$\bar{c}(\mathbf{L}) = \frac{\sum_{\mathbf{S} \in L} c^{[1;M]}(\mathbf{S}, \mathbf{L})}{\sum_{\mathbf{S} \in L} \delta^{[1;M]}(\mathbf{S}, \mathbf{L})}, \quad (6)$$

and replace the zero SS count by $\bar{c}(\mathbf{L})$:

$$c^{[1;M]}(\mathbf{S}', \mathbf{L}) = \bar{c}(\mathbf{L}) \quad (\text{for nonsampled subzones}) \quad (7)$$

Figure 2C presents an example for this case. Finally, all subzones in the IS zone have a non-zero value: $c^{[1;M]}(\mathbf{S}, \mathbf{L}) > 0$ ($\forall \mathbf{S}$ in ζ_L).

Procedure 3: All of the SS counts are zero if ζ_L is an E zone in all the iterations. Then

we correct $c^{[1;M]}(\mathbf{S}, \mathbf{L})$ simply as

$$c^{[1;M]}(\mathbf{S}, \mathbf{L}) = 1 \quad (\text{if } \zeta_L \text{ is E zone in all iterations}). \quad (8)$$

Figure 2D shows procedure 3.

Computation of $Q_{cano}(\mathbf{L})$ from SS counts

In this section, we present a method to calculate $Q_{cano}^{[M]}(\mathbf{L})$ by fitting SS counts $c^{[1;M]}(\mathbf{S}, \mathbf{L})$ among neighboring subzones. The idea of subzone fitting is originated from our earlier study [4] (see Eq. 4.2 of Ref. 4). The fitting procedure is shown schematically by the distribution transformation presented in the earlier report (Fig. 5B $\rightarrow$ Fig. 5C of Ref. [4]), where fractions (short colored curves) of a canonical distribution produce the full distribution (black-line curve). Although the fitting method presented in that earlier report is not the SS count-fitting but function-fitting, the logic is the same.

Assuming n_{is} intersections in the entire RC space, and assuming that these intersections are numbered in an arbitrary order, then we designate these subzones as a *subzone pair* if a couple of subzones mutually overlap in the RC space. Here, the number of subzone pairs originated from the j -th intersection ($j = 1, \dots, n_{is}$) is represented by $n_{pair}(j)$: $n_{pair}(j) = n_{link}(n_{link} - 1)/2$; also, n_{link} is a function of j ($n_{link} = n_{link}(j)$) because n_{link} depends on the position of the j -th intersection in the RC space. Figure 3 is a 2D RC space presented in the extension form. Four subzones represented by red circles originate from an intersection. The subzones generate six ($= 4 \times 3/2$) subzone pairs, which are indicated by red lines. Similarly, four subzones shown by blue circles originate from another intersection. Six blue lines specify subzone pairs. Similar arguments can be made for the mD RC space. One can assume a k -th subzone pair with respect to the j -th intersection ($k = 1, \dots, n_{pair}(j)$). We denote the indices of the pairing subzones by $p_{k,j}^{(1)}$ and $p_{k,j}^{(2)}$, where superscripts (1) and (2) show simply that the two subzones are pairing. Similarly, we designate the zone indices (virtual states) regarding the subzone pairs as $q_{k,j}^{(1)}$ and $q_{k,j}^{(2)}$. Then, the pairing subzones

in the entire RC space are expressed as $[\mathbf{S}_{p_{k,j}^{(1)}}, \mathbf{L}_{q_{k,j}^{(1)}}]$ and $[\mathbf{S}_{p_{k,j}^{(2)}}, \mathbf{L}_{q_{k,j}^{(2)}}]$. Any of the pairing subzones in the RC space can be specified with varying k and j .

Next, we define a *modulated SS count* ρ by multiplying a factor $\sigma_{\mathbf{L}}$ to

$c^{[1;M]}(\mathbf{S}, \mathbf{L})$ as

$$\rho^{[M]}(\mathbf{S}, \mathbf{L}; \sigma_{\mathbf{L}}) = \sigma_{\mathbf{L}} c^{[1;M]}(\mathbf{S}, \mathbf{L}), \quad (9)$$

where $\sigma_{\mathbf{L}}$ depends on $\mathbf{L}$. Finally, we define an objective function F as

$$F(\boldsymbol{\sigma}) = \sum_{j=1}^{n_{is}} \sum_{k=1}^{n_{pair}^{(j)}} (f_{j,k} - 1), \quad (10)$$

where

$$f_{j,k} = \frac{\max \left[\rho^{[M]}(\mathbf{S}_{p_{k,j}^{(1)}}, \mathbf{L}_{q_{k,j}^{(1)}}; \sigma_{\mathbf{L}_{q_{k,j}^{(1)}}}), \rho^{[M]}(\mathbf{S}_{p_{k,j}^{(2)}}, \mathbf{L}_{q_{k,j}^{(2)}}; \sigma_{\mathbf{L}_{q_{k,j}^{(2)}}}) \right]}{\min \left[\rho^{[M]}(\mathbf{S}_{p_{k,j}^{(1)}}, \mathbf{L}_{q_{k,j}^{(1)}}; \sigma_{\mathbf{L}_{q_{k,j}^{(1)}}}), \rho^{[M]}(\mathbf{S}_{p_{k,j}^{(2)}}, \mathbf{L}_{q_{k,j}^{(2)}}; \sigma_{\mathbf{L}_{q_{k,j}^{(2)}}}) \right]}. \quad (11)$$

The multiplication factors, $\sigma_{L_1}, \sigma_{L_2}, \dots$, are packed in a vector form: $\boldsymbol{\sigma} = [\sigma_{L_1}, \sigma_{L_2}, \dots]$.

Equation 11 is invariant in exchange of superscripts (1) and (2). Double summations in Eq. 11 move over all subzone pairs in the entire RC space. The “ -1 ” in Eq. 10 is introduced to set F to zero when $\rho^{[M]}(\mathbf{S}_{p_{kj}^{(1)}}, \mathbf{L}_{q_{kj}^{(1)}}; \sigma_{L_{q_{kj}^{(1)}}}) = \rho^{[M]}(\mathbf{S}_{p_{kj}^{(2)}}, \mathbf{L}_{q_{kj}^{(2)}}; \sigma_{L_{q_{kj}^{(2)}}})$. Then, we minimize F by modulating $\boldsymbol{\sigma}$ and refer to the resultant $\boldsymbol{\sigma}$ providing the minimized F as $\boldsymbol{\sigma}^{min} = [\sigma_{L_1}^{min}, \sigma_{L_2}^{min}, \dots]$. Finally, $Q_{cano}^{[M]}(\mathbf{L})$ is given as

$$Q_{cano}^{[M]}(\mathbf{L}) = \sum_{\mathbf{S}_i \in \mathbf{L}} \rho^{[M]}(\mathbf{S}_i, \mathbf{L}; \sigma_{\mathbf{L}}^{min}), \quad (12)$$

where the summation is taken over all subzones in zone $\boldsymbol{\zeta}_L$.

Minimization of F corresponds to fitting the function $c^{[1;M]}(\mathbf{S}, \mathbf{L})$ among neighboring zones by modulating $\boldsymbol{\sigma}$. This fitting is shown schematically in Fig. 4, which is presented in a 1D RC space to make the fitting procedure simple. Figure 4A shows a plot of the SS counts $c^{[1;M]}(\mathbf{S}, \mathbf{L})$ before fitting, where the SS counts of two pairing subzones (i.e., two open circles in a broken-line rectangle) are not mutually similar. Figure 4B presents the modulated SS counts after fitting, where the two

modulated SS counts pairing (circles unified in the rectangle) have mutually equal values.

In the 1D RC space, the objective function F converges analytically to zero by minimization. In a 2D or higher dimensional RC space, however, F might not converge to zero; also, σ^{min} is not determined analytically. Consequently, one might minimize F by a Monte Carlo simulation with modulation of σ . Using the converged σ^{min} , $Q_{cano}^{[M]}(\mathbf{L})$ is calculated (Eq. 12). Also, $J_{\mathbf{L}^{(i)}}$ is set as

$$J_{\mathbf{L}^{(i)}}^{[M]} = \left[Q_{cano}^{[M]}(\mathbf{L}^{(i)}) \sum_{j=1}^{n_{link}} \frac{1}{Q_{cano}^{[M]}(\mathbf{L}^{(j)})} \right]^{-1}, \quad (13)$$

which is equivalent to Eq. S7 in SI. Then the $(M + 1)$ -th iteration is done using $J_{\mathbf{L}^{(i)}}^{[M]}$.

It is noteworthy that $Q_{cano}^{[M]}(\mathbf{L})$ is calculated using $Q_{cano}^{[M-1]}(\mathbf{L})$ in the original method (Eq. S8 of SI). Also, a convergence check ($Q_{entire}^{[M]}(\mathbf{L}) \approx const$; see section 5 of SI) is necessary to judge the quality of the resultant $Q_{cano}^{[M]}(\mathbf{L})$. The subzone-based method requires neither $Q_{cano}^{[M-1]}$ nor a convergence check. The fewer requirements

stand as a benefit of the subzone-based method, and also of the GA method explained later.

Furthermore, we can assess the accuracy of the resultant $Q_{entire}^{[M]}(\mathbf{L})$ locally in each zone: Given a zone $\zeta_{\mathbf{L}}$, we first list subzone pairs relating to the zone. The green lines in Fig. 3 show subzone pairs related to a zone $\zeta_{\mathbf{L}_7}$. We express the i -th subzone pair as $[\mathbf{S}_{p_i^{(1)}}, \mathbf{L}]$ and $[\mathbf{S}_{p_i^{(2)}}, \mathbf{L}_{q_i^{(2)}}]$. In the i -th subzone pair, the superscript (1) is assigned to the subzone $[\mathbf{S}_{p_i^{(1)}}, \mathbf{L}]$ involved in $\zeta_{\mathbf{L}}$. Also, (2) represents subzone $[\mathbf{S}_{p_i^{(2)}}, \mathbf{L}_{q_i^{(2)}}]$, which is not involved in $\zeta_{\mathbf{L}}$ in the subzone pair. Then, a local function, $E_{local}(\mathbf{L})$, is defined as

$$E_{local}(\mathbf{L}) = \frac{1}{N_{pair}(\mathbf{L})} \sum_{i=1}^{N_{pair}(\mathbf{L})} \frac{\max \left[\rho^{[M]}(\mathbf{S}_{p_i^{(1)}}, \mathbf{L}; \sigma_{\mathbf{L}}^{min}), \rho^{[M]}(\mathbf{S}_{p_i^{(2)}}, \mathbf{L}_{q_i^{(2)}}; \sigma_{\mathbf{L}_{q_i^{(2)}}}^{min}) \right]}{\min \left[\rho^{[M]}(\mathbf{S}_{p_i^{(1)}}, \mathbf{L}; \sigma_{\mathbf{L}}^{min}), \rho^{[M]}(\mathbf{S}_{p_i^{(2)}}, \mathbf{L}_{q_i^{(2)}}; \sigma_{\mathbf{L}_{q_i^{(2)}}}^{min}) \right]} - 1, \quad (14)$$

where $N_{pair}(\mathbf{L})$ represents the number of subzone pairs related to the zone $\zeta_{\mathbf{L}}$. The summation is taken over all subzone pairs related to the zone $\zeta_{\mathbf{L}}$. Equation 14 can be regarded as a part of Eq. 10.

If the subzone-based mD-VcMD simulation is sufficiently long, then the modulated SS counts fit well between the pairing subzones in theory, which results in $E_{local}(\mathbf{L}) \rightarrow 0$ ($\forall \mathbf{L}$). In an actual simulation, the sampling accuracy might be uneven: Some RC regions might be sampled poorly. One need not sample this region further if a region is unimportant biophysically. However, if the region is important, then the region should be sampled selectively. A simple method for selective sampling is presented in subsection “Selectively distributed initial conformations (SDIC)”. We can judge that a zone ζ_L is sampled accurately when $E_{local}(\mathbf{L}) \leq E_{sh}$ is satisfied, where E_{sh} is a threshold set by a user. For our actual sampling, we set $E_{sh} = 4$ [9].

GA to Determine $c^{[1;M]}(\mathbf{S}, \mathbf{L})$ for IS and E Zones

Procedures 1–3 in subsection “Introduction of subzone and subzone’s snapshot count (SS count)” proposed the method to fill SS counts $c^{[1;M]}(\mathbf{S}, \mathbf{L})$ for all the CS, IS, and E zones when the M -th iteration is finished. Then, we may proceed to the next iteration (iteration $M + 1$) using the obtained SS counts. However, these procedures do not provide a method to presume the SS counts for IS and E zones. Here, we introduce the GA-guided method to fill $c^{[1;M]}(\mathbf{S}, \mathbf{L})$ in the IS and E zones.

GA database

For preparation, we generate a database (called *GA database*) used for the GA procedure as follows: First, we consider a compact block composed of CS zones:

$$\{\zeta_L: L = [i + \Delta i, j + \Delta j, k + \Delta k, \dots]; \Delta i, \Delta j, \Delta k, \dots = -1, 0, +1\}. \quad (15)$$

This block is called a *GA block*. We refer to the zone at the center of the block ($\Delta i = \Delta j = \Delta k = \dots = 0$) as a *central GA zone*. Figures 5A and 5B respectively portray the blocks in 2D and 3D RC spaces. In the 2D RC space, GA blocks are classified into three types: inside, side, and corner GA blocks, depending on the position of the central GA zone in the RC space. In the 3D RC space, blocks of four types are possible: inside, face, side, and corner GA blocks. If a block is not an inside one, then the variations of $\Delta i, \Delta j, \Delta k, \dots$ in Eq. 15 might range from 0 to +1 or from -1 to 0. In a higher dimensional RC space, we should classify the blocks into more than four types. When the M -th iteration has done, we have SS counts from various iteration windows: $c^{[1;1]}$

$(\mathbf{S}, \mathbf{L})$, $c^{[1;2]}(\mathbf{S}, \mathbf{L})$, ..., $c^{[1;M]}(\mathbf{S}, \mathbf{L})$, as computed using Eqs. 4 and 5. Then, the GA database consists of GA blocks from those various iteration windows.

Figure 6A portrays an example of GA blocks in the 2D RC space. We designate the GA blocks as GA_j ($j = 1, \dots, 6$), where j is set in an arbitrary order. The GA database might involve GA blocks that are taken from the same area of the RC space but from different iteration windows. We regard those GA blocks as different ones in the GA database. Consequently, the number of GA blocks in the GA database increases with progress of the iteration.

Each zone in GA_i is expressed by its relative position to the central GA zone. The relative position is expressed using a relative index $\mathbf{\Lambda}$: $\mathbf{\Lambda} = [\Lambda^{(\alpha)}, \Lambda^{(\beta)}, \Lambda^{(\gamma)}, \dots]$ and $\Lambda^{(h)} = -1, 0, 1$ ($h = \alpha, \beta, \gamma, \dots$) if the block GA_i is an inside block. For a non-inside block, Λ_j ranges as $\Lambda^{(h)} = -1, 0$ or $\Lambda^{(h)} = 0, -1$. It is noteworthy that a subzone $\mathbf{S}$ in a zone $\mathbf{\Lambda}$ is expressed by its relative position in the zone (Fig. 1B). Then, we specify the subzone by $[\mathbf{S}, \mathbf{\Lambda}]$. The SS count for the subzone $[\mathbf{S}, \mathbf{\Lambda}]$ in GA_i is expressed as $c^{GA_i}(\mathbf{S}, \mathbf{\Lambda})$.

Recovery zone

Next, we proceed to estimation of SS counts in IS and E zones without using procedures 2 and 3 in subsection “Introduction of subzone and subzone’s snapshot count (SS count)”. After the M -th iteration, we select an IS or E zone. One can assume that the selected zone is surrounded by N_{surr} zones. For instance, in the 2D RC space, $N_{surr} = 8, 5$ or 3 for an inside, side, or corner zone, respectively (Fig. 5A), and for the 3D RC zone, $N_{surr} = 26, 17, 11$, or 7 for an inside, face, side, or corner zone, respectively (Fig. 5B). If the selected IS/E zone is surrounded by more than $N_{surr}/2$ CS zones, then this selected zone is designated as a *recovery zone (REC zone)*. Figure 6B presents examples of four REC zones, where REC zones 1, 2, and 3 are surrounded respectively by six, six, and five CS zones of eight zones ($N_{surr} = 8$). REC zone 4 is surrounded by four CS zones of five zones ($N_{surr} = 5$). A compact block consisting of the REC zone and zones surrounding the REC zone is called a *recovery block (REC block)* if a selected zone is judged as a REC zone. The i -th REC block is denoted by REC_i . Similarly to GA blocks, REC blocks are classified into some types: inside, side, and corner blocks for a 2D RC space, and inside, face, side, corner blocks for a 3D RC space.

In fact, more REC zones can be taken than those shown in Fig. 6A. Note that the REC zones can overlap in the RC space: REC_2 overlaps REC_1 and REC_3 in Fig. 6A. We also note that the SS counts, $c^{[1;M]}(\mathbf{S}, \mathbf{L})$, in a REC block are computed only from the iteration window $[1;M]$ (Eqs. 4 and 5), in contrast to those in a GA block, which are computed from various iteration windows $[1;1], \dots, [1;M]$.

Resetting REC block

We use indices $[\mathbf{S}, \mathbf{\Lambda}]$ to specify subzones in a REC block, as well as those in a GA block, and refer to an SS count, $c^{[1;M]}(\mathbf{S}, \mathbf{\Lambda})$, in REC_i as $c^{REC_i}(\mathbf{S}, \mathbf{\Lambda})$. One should recall that the REC zone in all REC blocks are IS or E zone (Fig. 6B). Now we reset the SS counts $c^{REC_i}(\mathbf{S}, \mathbf{\Lambda})$ of REC_i as $c^{REC_i}(\mathbf{S}, \mathbf{\Lambda}) \rightarrow 1$ ($\forall \mathbf{S}; \mathbf{\Lambda} = [0,0,0]$). For that reason, the REC zones become E zones (Fig. 6C). The SS counts in this reset REC zone are estimated using the GA procedure described below.

Similarity of spatial patterns of SS counts between of REC_i and GA_j

To assess similarity of the SS counts between blocks REC_i and GA_j , we introduce a correlation coefficient $c_{cor}(REC_i, GA_j)$:

$$c_{cor}(REC_i, GA_j) = \frac{\langle REC_i \times GA_j \rangle - \langle REC_i \rangle \langle GA_j \rangle}{SD_{REC_i} \times SD_{GA_j}}, \quad (16)$$

where

$$\langle REC_i \rangle = \frac{1}{N_{s\Lambda}} \sum_{\Lambda \in REC_i} \sum_{S \in \Lambda} c^{REC_i}(S, \Lambda), \quad (17)$$

$$\langle GA_j \rangle = \frac{1}{N_{s\Lambda}} \sum_{\Lambda \in GA_j} \sum_{S \in \Lambda} \delta_{REC_i}(S, \Lambda) c^{GA_j}(S, \Lambda), \quad (18)$$

$$SD_{REC_i} = \left[\frac{1}{N_{s\Lambda}} \sum_{\Lambda \in REC_i} \sum_{S \in \Lambda} c^{REC_i}(\Lambda, S)^2 - \langle REC_i \rangle^2 \right]^{1/2}, \quad (19)$$

$$SD_{GA_j} = \left[\frac{1}{N_{s\Lambda}} \sum_{\Lambda \in GA_j} \sum_{S \in \Lambda} \delta_{REC_i}(S, \Lambda) c^{GA_j}(S, \Lambda) - \langle GA_j \rangle^2 \right]^{1/2}, \quad (20)$$

$$\langle REC_i \times GA_j \rangle = \frac{1}{N_{s\Lambda}} \sum_{\Lambda \in REC_i} \sum_{S \in \Lambda} c^{REC_i}(\Lambda, S) c^{GA_j}(\Lambda, S), \quad (21)$$

$$N_{s\Lambda} = \sum_{\Lambda \in REC_i} \sum_{S \in \Lambda} \delta_{REC_i}(S, \Lambda), \quad (22)$$

$$\delta_{REC_i}(S, \Lambda) = \begin{cases} 1 & (\text{If } c^{REC_i}(\Lambda, S) \neq 0) \\ 0 & (\text{else}) \end{cases}. \quad (23)$$

The double summations $\sum_{\Lambda \in REC_i} \sum_{S \in \Lambda}$ and $\sum_{\Lambda \in GA_j} \sum_{S \in \Lambda}$ are taken respectively over all subzones in REC_i and GA_j . In Eq. 21, the summation $\sum_{\Lambda \in REC_i}$ operates not only to $c^{REC_i}(\Lambda, S)$ but also to $c^{GA_j}(\Lambda, S)$: The same relative positions of Λ in GA_j and REC_i are summed in a coordinated manner.

We convert $c_{cor}(REC_i, GA_j)$ to a score function $E_{score}(REC_i, GA_j)$ as

$$E_{simi}(REC_i, GA_j) = \begin{cases} \frac{1}{c_{cor}(REC_i, GA_j)} - 1 & (\text{if } c_{cor}(REC_i, GA_j) > c_{sh}) \\ \frac{1}{c_{sh}} - 1 & (\text{if } c_{cor}(REC_i, GA_j) \leq c_{sh}) \end{cases}. \quad (24)$$

The delta function $\delta_{REC_i}(\mathbf{S}, \mathbf{\Lambda})$ is used in $c_{cor}(REC_i, GA_j)$. Therefore, a non-zero c^{GA_j}

$(\mathbf{\Lambda}, \mathbf{S})$ in GA_j is eliminated from the computation of E_{simi} if $c^{REC_i}(\mathbf{S}, \mathbf{\Lambda}) = 0$.

Furthermore, the REC zone of REC_i is always eliminated from E_{simi} because δ_{REC_i}

$(\mathbf{S}, \mathbf{\Lambda}) = 0$ for $\mathbf{\Lambda} = [0, 0, 0]$. Parameter c_{sh} is introduced to avoid a negative value of

E_{simi} by a negative correlation $c_{cor}(REC_i, GA_j)$. Consequently, c_{sh} can be set to a

small positive value ($c_{sh} = 0.01$ for instance).

One may replace the SS counts of the REC zone of REC_i by those of the central GA zone of GA_j if E_{simi} is smaller than a certain value. However, we assess the quality of GA_j in the next section to judge if the replacement is appropriate.

Overlap of SS-count patterns among zones in a GA block

This subsection explains a method to assess the quality of a GA block GA_j by introducing E_{phys} , which quantifies the spatial overlap of SS counts between

overlapping subzones within the block. In fact, E_{phys} is a quantity expressed only by the SS counts in GA_j independent of REC_i .

One can consider two subzones $[\mathbf{S}, \mathbf{\Lambda}]$ and $[\mathbf{S}', \mathbf{\Lambda}']$ in GA_j . First, we introduce a δ function $\delta_{ov}(\mathbf{L}, \mathbf{S}; \mathbf{L}', \mathbf{S}')$, which is used as a flag to detect overlapping subzones in GA_j :

$$\delta_{ov}(\mathbf{S}, \mathbf{\Lambda}; \mathbf{S}', \mathbf{\Lambda}') = \begin{cases} 1 & ([\mathbf{S}, \mathbf{\Lambda}] \text{ and } [\mathbf{S}', \mathbf{\Lambda}'] \text{ are overlapping}) \\ 0 & (\text{else}) \end{cases}. \quad (25)$$

For instance, $\delta_{ov}(\mathbf{L}, \mathbf{S}; \mathbf{L}', \mathbf{S}') = 1$ for any subzone pair taken from the shaded ones in Fig. 1B.

Similarly to Eq. 9, the SS counts $c^{GA_j}(\mathbf{S}, \mathbf{\Lambda})$ can be scaled by a factor $\sigma_{\mathbf{\Lambda}}$, and generate modulated SS counts as $\rho^{GA_j}(\mathbf{S}, \mathbf{\Lambda}; \sigma_{\mathbf{L}}) = \sigma_{\mathbf{\Lambda}} c^{GA_j}(\mathbf{S}, \mathbf{\Lambda})$. Then, we define a function E_{phys} as

$$E_{phys}(GA_j) = \sum_{\mathbf{\Lambda} \in GA_j}^{\mathbf{\Lambda} \neq \mathbf{0}} \sum_{\mathbf{\Lambda}' \in GA_j}^{\mathbf{\Lambda}' \neq \mathbf{0}, \mathbf{\Lambda}' \neq \mathbf{\Lambda}} \delta_{ov}(\mathbf{S}, \mathbf{\Lambda}; \mathbf{S}', \mathbf{\Lambda}') P(\sigma_{\mathbf{L}}, \sigma_{\mathbf{L}'}), \quad (26)$$

where

$$P(\sigma_L, \sigma_{L'}) = \frac{\max[\rho^{GA_j}(\mathbf{s}, \mathbf{\Lambda}; \sigma_L), \rho^{GA_j}(\mathbf{s}, \mathbf{\Lambda}; \sigma_{L'})]}{\min[\rho^{GA_j}(\mathbf{s}, \mathbf{\Lambda}; \sigma_L), \rho^{GA_j}(\mathbf{s}, \mathbf{\Lambda}; \sigma_{L'})]} - 1. \quad (27)$$

The summation $\sum_{\mathbf{\Lambda}' \in GA_j, \mathbf{\Lambda}' \neq \mathbf{\Lambda}}$ in Eq. 26 is taking pairs of different zones ($\mathbf{\Lambda}' \neq \mathbf{\Lambda}$). The summations are taken with eliminating zone pairs related to the central GA zone ($\mathbf{\Lambda} \neq \mathbf{0}$ and $\mathbf{\Lambda}' \neq \mathbf{0}$). The number of zones in a GA block depends on the place where the block is taken from the RC space (Fig. 5). Therefore, the summation of Eq. 26 is taken over existing zones in the GA block. For instance, a surface GA block involves 18 zones in Fig. 5B.

Although $F(\sigma)$ (Eq. 10) was defined for the whole RC space, E_{phys} is defined for a single GA block. Finally, we minimize E_{phys} with modulating $\{\sigma_L\}$ using a Monte Carlo scheme. The smaller the minimized $E_{phys}(GA_j)$ becomes, the better the quality of the block GA_j is. The E_{phys} used in the next section is this minimized one.

First generation of GA

To start the GA procedure, the first generation, denoted as Gen_1 , is set by taking GA blocks from the GA database. Now, we specifically examine block REC_i , which means that the SS counts in REC_i are determined by the GA operation. Then, we calculate $E_{simi}(REC_i, GA_j)$ and $E_{phys}(GA_j)$ for all the GA blocks in the GA database, and define the score function as

$$E_{score}(REC_i, GA_j) = \omega_s E_{simi}(REC_i, GA_j) + \omega_p E_{phys}(GA_j), \quad (28)$$

where ω_s and ω_p are weights set by the user, such as 1.0 and 2.0, respectively, as described in our earlier report [9]. The smaller the E_{score} is, the more similar the SS-count patterns of GA_j to REC_i and the better the overlap of SS counts among zones in GA_j .

Next, we sort the resultant scores $\{E_{score}(REC_i, GA_j), \forall j\}$ in ascending order and collect N_{GA}^{all} GA blocks from the first score. Those N_{GA}^{all} GA blocks construct the first generation, denoted as Gen_1 .

The first generation is prepared for each of the REC blocks. The size of a REC block depends on its place in the RC space, as described previously. We express the size of REC_i and GA_j as $n_{\alpha}^{REC_i} \times n_{\beta}^{REC_i} \times n_{\gamma}^{REC_i} \times \dots$ and $n_{\alpha}^{GA_j} \times n_{\beta}^{GA_j} \times n_{\gamma}^{GA_j} \times \dots$, respectively. Then, the GA blocks used in E_{score} can be expected to satisfy the following relations: $n_h^{REC_i} \leq n_h^{GA_j}$ ($h = \alpha, \beta, \gamma, \dots$).

GA generation update

Presuming that we have the k -th GA generation, denoted as Gen_k , with respect to a recovery block REC_i , then the generation update from Gen_k to Gen_{k+1} is presented in Fig. 7. Remember that the first generation of GA consists of N_{GA}^{all} blocks. This number N_{GA}^{all} is kept in all generations of GA. Next, one can select the N_{GA}^{top} members from the first to N_{GA}^{top} -th scored members from Gen_k ($N_{GA}^{all} > N_{GA}^{top}$) and pass them to the next generation Gen_{k+1} , as shown in the red-framed rectangle in Fig. 7. Consequently, N_{GA}^{top} members in Gen_{k+1} are taken from Gen_k without modulation.

Next, we randomly select N_{GA}^{mut} members from Gen_k , which are denoted as $MEM_k^{(1)}, MEM_k^{(2)}, \dots, MEM_k^{(N_{GA}^{mut})}$, and the same number of GA blocks from the GA database, denoted as $GA^{(1)}, GA^{(2)}, \dots, GA^{(N_{GA}^{mut})}$. Then, we replace a part of $MEM_k^{(l)}$

by a part of $GA^{(l)}$ ($l = 1, \dots, N_{GA}^{mut}$). This procedure corresponds to *mutation* in GA. In Fig. 7, the cyan-colored zones of a member in Gen_k are replaced by the cyan-colored zones in a GA block. Similarly, the yellow ones of a block in Gen_k are replaced by the yellow ones in a GA block. The mutated members are put in Gen_{k+1} .

Up to this point, $N_{GA}^{top} + N_{GA}^{mut}$ generation members are set for Gen_{k+1} . Here, we randomly select $N_{GA}^{exc} (= N_{GA}^{all} - N_{GA}^{top} - N_{GA}^{mut})$ from Gen_k and generate $N_{GA}^{exc}/2$ pairs of the selected members: N_{GA}^{top} and N_{GA}^{mut} should be set so that N_{GA}^{exc} is an even number. We denote the m -th pair by $MEM_k^{(m)(1)}-MEM_k^{(m)(2)}$ ($m = 1, \dots, N_{GA}^{exc}/2$), where superscripts (1) and (2) show that the two members are pairing. Then, a part of $MEM_k^{(m)(1)}$ and a part of $MEM_k^{(m)(2)}$ are exchanged. In Fig. 7, two magenta parts are mutually exchanged. Two green parts are done too. The members generated by the exchange are put in Gen_{k+1} . This procedure corresponds to *crossover* or *exchange* in GA. Up to this point, N_{GA}^{all} members are prepared for Gen_{k+1} . Details for mutation and exchange are presented later.

To generate Gen_{k+2} , we repeat similar procedures to Gen_{k+1} as done for Gen_k : (1) Calculate the score function between each REC block (say REC_i) and all N_{GA}^{all} members in Gen_{k+1} (not all blocks in the GA database):

$$E_{score}(REC_i, MEM_{j,k+1}) = w_s E_{simi}(REC_i, MEM_{j,k+1}) + w_p E_{phys}(MEM_{j,k+1}), \quad (29)$$

where $MEM_{j,k+1}$ stands for the j -th member ($j = 1, \dots, N_{GA}^{all}$) in Gen_{k+1} . (2) The first

to N_{GA}^{top} scored members of Gen_{k+1} are passed to the next generation Gen_{k+2} . (3)

Randomly selecting N_{GA}^{mut} members from Gen_{k+1} and N_{GA}^{mut} GA blocks from the GA

database, mutations are executed. The generated N_{GA}^{mut} members are passed to Gen_{k+2} .

(4) Selecting N_{GA}^{exc} members randomly from Gen_{k+1} , generate $N_{GA}^{exc}/2$ pairs of

members. Then exchange is done in each pair. The resultant N_{GA}^{exc} members are passed

to Gen_{k+2} . Up to this point, N_{GA}^{all} members are prepared for Gen_{k+2} .

We repeat this cycle N_{cycle} times, where N_{cycle} is set by the user. Finally, the SS counts of the REC zone in REC_i (i.e., the central zone of REC_i) are replaced by those of the central zone of the best-scored member of $Gen_{N_{cycle}}$. The user might exit

the cycle before reaching the N_{cycle} cycle if E_{simi} and E_{phys} of a member become smaller than the respective thresholds set by the user.

Multiple GA procedure

Figure 8A is a part of 2D RC space, where the magenta-colored circles are REC zones. The SS counts for these REC zones are estimated by the GA (Fig. 8B). Then, one might calculate $Q_{cano}^{[M]}(\mathbf{L})$ and $J_{L(i)}^{[M]}$ from these SS counts and then proceed to the $(M + 1)$ -th iteration. In Fig. 8B, however, two zones indicated by cyan-colored circles emerge as REC zones anew. Therefore, we can advance the second GA procedure to estimate the cyan-colored zones. Figure 8C shows that the cyan-colored zones are determined. However, another REC zone (light-yellow zone) appears in Fig. 8C. Therefore, we can multiply the GA procedure. When a new REC zone does not appear anymore, we quit the procedure. Then we proceed to the $(M + 1)$ -th iteration. Supplementary Figure S3 depicts the procedures used for the GA-guided mD-VcMD method as a flow chart.

Mutation and exchange

We explain details for mutation and exchange. In mutation, a pair of $MEM_k^{(i)}$ ($i = 1, \dots, N_{GA}^{mut}$) and a GA block $GA^{(i)}$ is treated. In exchange, a pair of $MEM_k^{(m)(1)}$ ($m = 1, \dots, N_{GA}^{exc}/2$) and $MEM_k^{(m)(2)}$ is treated. In either operation, we select a slice from $MEM_k^{(i)}$, $GA^{(i)}$, $MEM_k^{(m)(1)}$, and $MEM_k^{(m)(2)}$. Imagine slices in a block as

$$\Lambda^{(h)} = \begin{cases} -1 \\ +1 \end{cases} \quad (h = \alpha, \beta, \gamma, \dots). \quad (30)$$

Figure 9 presents an image of slices in the 3D RC space to help understanding of the slices. Each slice consists of nine zones.

In mutation, a slice, say $\Lambda^{(x)}$, is selected randomly from slices of $MEM_k^{(i)}$. Then, a slice parallel to $\Lambda^{(x)}$ is selected randomly from slice $\Lambda^{(x)} = -1$ or $\Lambda^{(x)} = +1$ of $GA^{(i)}$. Then, the selected slice of $MEM_k^{(i)}$ is replaced by that of $GA^{(i)}$, which generates a new member for Gen_{k+1} .

In exchange, after selecting a slice randomly from slices of $MEM_k^{(m)(1)}$, a slice is selected randomly from slices of $MEM_k^{(m)(2)}$ in the same manner as that used

for mutation. Then, the selected slices are mutually exchanged. This procedure generates two members for Gen_{k+1} .

No middle slice ($\Lambda^{(x)} = 0$) is used for the mutation and exchange. The GA procedure is used to estimate the SS counts of a REC zone. The REC zone is involved in the middle slices of a REC block. To maintain logical consistency between the REC zone and the member in Gen_k , we eliminated the middle slice.

A technical point must be noted: In mutation and exchange, the size of the operated slice of a REC block should be included in or should be the same as that of a block in Gen_k . The size of the operated slice of the block in Gen_k should be included in or should be the same as that of a GA block.

Initial Conformations of Simulation

Evenly distributed initial conformations (EDIC)

The initial conformation of the first iteration can be any conformation for all the original, subzone-based and GA-guided mD-vcMD methods. However, to discover pathways from the unstable state to the most stable conformation (i.e., the native tertiary

structure), the initial conformation should be far from the most stable conformation. Then, it is readily apparent that such sampling is free from knowledge of the native tertiary structure.

We have performed multiple runs for the mD-VcMD method to obtain the thermodynamically plausible distribution (the canonical distribution) of a biological system [10,11]. One can assume that iteration M , which consists of N_{run} runs, is finished. The initial conformation of the i -th run of iteration $M + 1$ can be the last conformation of the i -th run of iteration M . The first iteration is an exception, where the initial conformations are a single conformation as described above.

A more appropriate method to prepare the initial conformations is the following: Imagine an ensemble of the last conformations of the first to M -th iterations. The ensemble comprises of $N_{run} \times M$ conformations. Supplementary Figure S4A schematically presents those conformations in the RC space, which is presented two-dimensionally for convenience. Then, we divide the RC space into small patches of equal size and count patches that involve one or more conformations. The patches are called “occupied patches.” A patch is not the virtual zone (ζ_L). Eighteen occupied

patches are presented by shaded rectangles in Supplementary Fig. S4. Some occupied patches may involve a single conformation. The others involve more conformations. Next, we select N_{run} patches randomly from the occupied ones, and select one conformation randomly from each of the selected patches. Red-filled circles are shown in the figure assuming that $N_{run} = 10$. Those conformations are used for the initial conformations for iteration $M + 1$. We divide the RC space into smaller patches if the number of occupied patches is smaller than N_{run} . However, if the number of occupied patches is considerably larger than N_{run} , then we increase the patch size. We designate this method as the “evenly distributed-initial-conformation (EDIC) method.”

Selectively distributed initial conformations (SDIC)

Another method can be used to set the initial conformations. It is simple and useful when a poorly sampled region emerged in the RC space. Assuming again that the M -th iteration has finished and assuming that an RC region was poorly sampled in all iterations, then the accuracy of $Q_{cano}^{[M]}(\mathbf{L})$ for the poorly sampled RC region can be increased using a simple procedure: We then select conformations which are in or near

the poorly sampled RC region, from trajectories done previously (iterations 1 to M).

Then, the $(M + 1)$ -th iteration is done starting from those conformations.

This selective sampling procedure does not work if $Q_{cano}^{[M]}(\mathbf{L})$ is calculated using the original method (Eq. S8 in SI) because this procedure increases $Q_{entire}^{[M]}(\mathbf{L})$ locally. This local inflation of $Q_{entire}^{[M]}(\mathbf{L})$ breaks the balance of $Q_{entire}^{[M]}$, resulting in errors of $Q_{cano}^{[M]}$ over a wide RC region.

By contrast, in the subzone-based and GA-guided methods, the artificial increase of $Q_{entire}^{[M]}(\mathbf{L})$ vanishes by fitting $\rho^{[M]}(\mathbf{S}, \mathbf{L}; \sigma_L^{min})$ among overlapping zones. This lack of an artificial increase is an important advantage of the subzone-based and GA methods against the original method. We designate this method as “selectively distributed-initial-conformation (SDIC).”

We introduced a selective sampling procedure in an earlier report [8]. However, the selective sampling introduced herein differs from that described in that earlier report.

Thermodynamic Weight Assigned to Sampled Conformations

It is fundamentally important to assign a thermodynamic weight to each snapshot for analyzing the resultant conformational ensemble. Given a zone ζ_L , the thermodynamic weight assigned to ζ_L is $Q_{cano}(\mathbf{L})$. If the simulation is at the M -th iteration, then $Q_{cano}(\mathbf{L}) = Q_{cano}^{[M]}(\mathbf{L})$. In the original mD-VcMD method, the weight assigned to the snapshot detected in ζ_L is proportional to $Q_{cano}(\mathbf{L})$ if $Q_{cano}(\mathbf{L}) \approx \text{const}$ is satisfied in the simulation because the number of snapshots detected in zones is approximately equivalent.

In the subzone-based and GA methods, however, sampling is flexible. For instance, according to the selective sampling proposed above, the number of snapshots in zones might be considerably uneven, which means that the thermodynamic weight assigned to a snapshot is not equivalent to $Q_{cano}(\mathbf{L})$. We present a simple method for assigning weight: Consider a snapshot, as detected in ζ_L , and presume that this snapshot is the i -th one in the conformational ensemble. Then, the thermodynamic weight, w_i , assigned to the i -th snapshot is

$$w_i = Q_{cano}(\mathbf{L}) / n_{snap}^{[1;M]}(\mathbf{L}), \quad (31)$$

where $n_{snap}^{[1;M]}(\mathbf{L})$ signifies the number of snapshots detected in $\zeta_{\mathbf{L}}$ in all iterations.

Although w_i is dependent on $\mathbf{L}$ in Eq. 31, once the weight is assigned, we can forget

$\mathbf{L}$. If a relation $n_{snap}^{[1;M]}(\mathbf{L}) \approx \text{const}$ is satisfied, then $w_i \approx Q_{cano}(\mathbf{L})$ substantially.

Application to a Simple System

We earlier applied the original mD-VcMD method and a conventional Monte Carlo method to a model system [12]. Although the analytical conformational distribution of the system was not obtained because of the complexity of its potential energy function, both methods mutually converged. Importantly, the convergence from the original mD-VcMD was 200 times quicker than that from the conventional Monte Carlo method.

As described below, we check the efficiency of the GA-guided and original mD-VcMD methods using another model system for which the conformational distribution is obtainable analytically.

Model system

In the model, a single particle with coordinates expressed by $[x, y, z]$, moves in a 3D space at a temperature on a potential-energy surface, E_{model} , defined as

$$E_{model}(r) = D r^2(r^2 - a)(r^2 - b), \quad (32)$$

where r is the distance of the particle from the origin $[0,0,0]$:

$$r^2 = x^2 + y^2 + z^2. \quad (33)$$

Actually, E_{model} has a spherical symmetry around the origin in the 3D space.

Parameters D , a , and b in Eq. 32 are set respectively to 0.045, 10, and 20. We set the temperature T to 1.5 or 3.0 for the simulations. Parameters D , a , and b and temperature T are introduced as non-dimensional quantities in this application. Then, the Boltzmann factor is defined simply as $\exp [-E_{model}/T]$.

Originally, the RCs $\lambda^{(\alpha)}$, $\lambda^{(\beta)}$, and $\lambda^{(\gamma)}$ were defined by the inter-atom-group distances in a molecular system (Section 1 of SI), and the motions of

RCs were calculated from the motions of the molecular conformation. By contrast, in this application, $\lambda^{(\alpha)}$, $\lambda^{(\beta)}$, and $\lambda^{(\gamma)}$ respectively correspond directly to x , y , and z without assuming the molecular system. The function $E_{model}(r)$ is simple as shown in Supplementary Fig. S5A of SI. This figure shows two energy basins at $r = 0.0$ and $r \approx 4.0$, which are mutually separated by an energy barrier at $r \approx 2.0$. Supplementary Figure S6 demonstrates the potential energy in the 2D space ($r^2 = x^2 + y^2$; $z = 0$). We designate the energy basins at $r = 0.0$ and $r \approx 4.0$ respectively as the “central basin” and “periphery basin.” The periphery basin is deeper than the central basin. The energy barrier at $r \approx 2.0$ is called “major barrier.”

We restrict the sampling region in a cubic box by setting the lower and upper limits to the coordinates as $-3.0 \leq x \leq 3.0$, $-3.0 \leq y \leq 3.0$, and $-3.0 \leq z \leq 3.0$. In the outside of the cubic box, E_{model} increases rapidly. By this restriction, this model system has nine energy basins in a strict sense. One is the central basin at the origin ($x = y = z = 0$). The periphery basin at $r \approx 2.0$ in Supplementary Fig. S5A is split into eight sub-basins near the edges of the cubic box. The energy barriers among the eight sub-basins are considerably lower than the major barrier. They are overcome

easily during simulation at the examined temperatures. Therefore, we do not specifically examine differences among the eight sub-basins but specifically examine traversal between the central and periphery basins in this study.

The virtual zones are set as listed in Supplementary Table S2. We divided each of x -, y -, and z -axes equally into 11 virtual zones ($\zeta_1^{(h)}, \dots, \zeta_{11}^{(h)}$; $h = x, y, z$). The width of each zone is 1.0, and the lower and upper limits for the axes are, respectively, -3.0 and 3.0 .

The difficulty of sampling can be modulated by varying the system temperature. Supplementary Figure S5B shows the Boltzmann factor, $\exp[-E_{model}(r)/T]$, assigned to a microscopic state located at r . We note that the factor is not $4\pi r^2 \exp[-E_{model}(r)/T]$. Measuring from the most probable position ($r \approx 2.0$), the probability at the major barrier is about 10^{-5} at $T = 3.0$ and 10^{-10} at $T = 1.5$. This difference of the probability strongly influences the sampling efficiency of the original mD-VcMD method, as demonstrated later. We used a computer program myPresto/omegagene for the simulations [13].

Simulations

We moved the particle by a Monte Carlo (MC) simulation for both the original and GA-guided mD-VcMD simulations. The current simulation should be called “VcMC” because MC is done instead of molecular dynamics (MD). However, “VcMD” is used for convenience in this study. Step size $\Delta r = [\Delta x^2 + \Delta y^2 + \Delta z^2]^{1/2}$ of the MC simulation was 0.02: Δr distributed evenly from 0.0 to 0.02 isotropically in the 3D space. The initial position of the particle for the first iteration was set at the origin $[0, 0, 0]$. We performed 30 runs in each iteration ($N_{run} = 30$). The number of simulation steps for a run was 1×10^7 . The total simulation length for an iteration was 3×10^8 steps. We used the EDIC method to prepare the initial conformations for iteration M ($M \neq 1$).

In the GA-guided method, when an iteration finishes, CS, IS, and E zones are detected. Then, REC zones are set and the SS counts in the REC zones are calculated to proceed into the next iteration. Generally, with the progress of the iterative simulation, the CS zones increases. Therefore, the IS or E zones decreases, which means that the REC zones decrease. Finally, when the IS or E zones vanished (i.e., the REC zones vanished), we switched the GA-guided method to the subzone-based method.

For the GA-guided VmMD method at $T = 1.5$ and 3.0 , respectively, 10 and 5 iterations were performed, whereas 13 and 5 iterations were done, respectively, for the original mD-VcMD at $T = 1.5$ and 3.0 . The interval for attempting inter-virtual state transition ($\Delta\tau$; see section 4 of SI) is set to 10 steps of MC simulation for both the GA-guided and original VmMD methods.

To assess the sampling, we calculated correlation coefficients between the resultant $Q_{cano}^{[M]}(\mathbf{L})$ and the analytical $Q_{cano}(\mathbf{L})$ denoted as $Q_{cano}^{anal}(\mathbf{L})$:

$$C_{cor}^{normal}(M) = \frac{\langle f^{[M]} \times f^{anal} \rangle - \langle f^{[M]} \rangle \langle f^{anal} \rangle}{SD[f^{[M]}] \times SD[f^{anal}]}, \quad (34)$$

where

$$f^{[M]}(\mathbf{L}) = Q_{cano}^{[M]}(\mathbf{L}), \quad (35)$$

and

$$f^{anal}(\mathbf{L}) = Q_{cano}^{anal}(\mathbf{L}). \quad (36)$$

The terms in Eq. 34 are given as

$$\langle f^{[M]} \rangle = \frac{1}{N_{site}} \sum_{\mathbf{L}} f^{[M]}(\mathbf{L}), \quad (37)$$

$$\langle f^{anal} \rangle = \frac{1}{N_{site}} \sum_{\mathbf{L}} f^{anal}(\mathbf{L}), \quad (38)$$

$$\langle f^{[M]} \times f^{anal} \rangle = \frac{1}{N_{site}} \sum_{\mathbf{L}} \{f^{[M]}(\mathbf{L}) \times f^{anal}(\mathbf{L})\}, \quad (39)$$

$$SD[f^{[M]}] = \left(\frac{1}{N_{site}} \sum_{\mathbf{L}} [f^{[M]}(\mathbf{L})]^2 - \langle f^{[M]}(\mathbf{L}) \rangle^2 \right)^{1/2}, \quad (40)$$

and

$$SD[f^{anal}] = \left(\frac{1}{N_{site}} \sum_{\mathbf{L}} [f^{anal}(\mathbf{L})]^2 - \langle f^{anal} \rangle^2 \right)^{1/2}. \quad (41)$$

Therein, N_{site} represents the number of virtual zones in the system ($N_{site} = 11 \times 11 \times 11$). Supplementary Figure S5B shows that the Boltzmann factor in the periphery basin ($r \approx 4$) is considerably larger than that of the other regions at both temperatures. Therefore, C_{cor}^{normal} is determined mostly by $Q_{cano}^{[M]}$ in the periphery basin, ignoring the other regions.

To assess the overall shape of $Q_{cano}(\mathbf{L})$ in the correlation calculation, we defined another coefficient with setting $g^{[M]}$ and g^{anal} as

$$g^{[M]}(\mathbf{L}) = \log_{10}[Q_{cano}^{[M]}(\mathbf{L})], \quad (42)$$

and

$$g^{anal}(\mathbf{L}) = \log_{10}[Q_{cano}^{anal}(\mathbf{L})]. \quad (43)$$

Then, the new correlation coefficient is defined as

$$C_{cor}^{log10}(M) = \frac{\langle g^{[M]} \times g^{anal} \rangle - \langle g^{[M]} \rangle \langle g^{anal} \rangle}{SD[g^{[M]}] \times SD[g^{anal}]}. \quad (44)$$

The terms involved in Eq. 44 are defined by the same equations as Eqs. 37–41 with replacement of $f^{[M] \rightarrow g^{[M]}}$ and $f^{anal \rightarrow g^{anal}}$.

Figures 10A and 10B demonstrate the correlation functions as a function of iteration No. M calculated respectively from the GA-guided mD-VcMD at $T = 1.5$ and 3.0. Figures 10C and 10D respectively show those from the original mD-VcMD at $T = 1.5$ and 3.0. Figures 10B and 10D indicate that both the GA-guided and original methods provided a quick convergence of $Q_{cano}^{[M]}$ to Q_{cano}^{anal} at $T = 3.0$, which means that both methods are useful when the free-energy barrier is low.

At the lower temperature ($T = 1.5$), the free-energy barrier became considerably higher than that at $T = 3.0$. For the GA-guided method, $Q_{cano}^{[M]}$ converged again quickly to Q_{cano}^{anal} : Even in iteration 3, $C_{cor}^{normal}(3) = 0.94$ and $C_{cor}^{log10}(3) = 0.94$. For the original method, it is interesting that $Q_{cano}^{[M]}$ converged to Q_{cano}^{anal} once: $C_{cor}^{normal}(3) = 0.99$ and $C_{cor}^{log10}(3) = 0.99$. However, C_{cor}^{normal} and C_{cor}^{log10} started decreasing for $M \geq 4$. Consequently, if we can quit the sampling at iteration 3, then the original

method is also a good method, even at $T = 1.5$. However, determination of the optimal number of iterations is not straightforward in general because the analytical $Q_{cano}^{anal}(\mathbf{L})$ is unknown a priori for a molecular system.

Figure 11 displays the spatial patterns of $Q_{cano}^{[M]}$ calculated from the simulations as well as the analytical Q_{cano}^{anal} at $T = 1.5$. The analytical one has spherical symmetry around the origin (Fig. 11A). Figure 11B shows that $Q_{cano}^{[10]}$ from iteration 10 of the GA-guided method is similar to the analytical one: $C_{cor}^{normal}(10) = 0.96$ and $C_{cor}^{log10}(10) = 1.00$. Figure 11C is $Q_{cano}^{[3]}$ from iteration 3 of the original method, which is also similar to the analytical one, whereas the distribution shape around the central basin is somewhat deviated from the spherical shape. By contrast, Fig. 11D shows $Q_{cano}^{[30]}$ from iteration 30 of the original method, which is largely deviated from the analytical one. At $T = 3.0$, the special patterns of $Q_{cano}^{[M]}$ were similar to Q_{cano}^{anal} at iteration 3. The patterns were not broken in further iterative simulations (data not shown).

SDIC method to assess the GA-guided mD-VcMD method

Lastly in this application section, we assessed the SDIC method by assuming a situation that the central region (i.e., the central basin) was insufficiently sampled during preceding iterations. Note that the SDIC method is not applicable to the original mD-VcMD method, in theory.

In the subsection presented above, we obtained $Q_{cano}^{[10]}$ from the 10-th iteration of the GA-guided method. To test the SDIC method efficiency, we generated $Q_{cano}^{modi}(\mathbf{L})$ by modifying $Q_{cano}^{[10]}(\mathbf{L})$ as follows: $Q_{cano}^{[10]}(\mathbf{L})$ for zones in the central region ($L^{(h)} = 4, \dots, 8; h = x, y, z$) was replaced by a small positive value ε . This procedure assumes that the central region was not sampled or poorly sampled in iterations $1, \dots, M$. The white region in Supplementary Fig. S7 of SI is the central region. Actually, ε should be neither zero nor negative because such a value of ε results in an infinite or negative IVT probability $J_{L(i)}^{[M]}$ (Eq. 13). Here, we set ε by the smallest value of $Q_{cano}^{[10]}(\mathbf{L})$ at the top of the major barrier. The $Q_{cano}^{[10]}(\mathbf{L})$ for the shaded region of Supplementary Fig. S7 was taken from the original $Q_{cano}^{[10]}(\mathbf{L})$. Then $Q_{cano}^{modi}(\mathbf{L})$ is defined as

$$Q_{cano}^{modi}(\mathbf{L}) = \begin{cases} \varepsilon & \text{(for zones in the central region)} \\ Q_{cano}^{[10]}(\mathbf{L}) & \text{(for outside of the central region)} \end{cases} \quad (45)$$

Subsequently, we performed the 11-th iteration of the GA-guided method at $T = 1.5$ using Q_{cano}^{modi} . We denote Q_{cano} resultant from the 11-th iteration as $Q_{cano}^{[11]}$. The number of runs and the simulation length of a run are the same as those for the iterations done earlier ($N_{run} = 30$ and 1×10^7 , respectively). All of the 30 runs started from a single conformation at the origin (red filled circle in Supplementary Fig. S7), which is involved in the central region. In other words, the initial conformations were selected from the poorly sampled region. This is a simple example of SDIC.

Results show high correlation coefficients between $Q_{cano}^{[11]}(\mathbf{L})$ and $Q_{cano}^{anal}(\mathbf{L})$:

$C_{cor}^{normal} = 0.97$ and $C_{cor}^{log10} = 1.0$. However, it is noteworthy that the coefficients

between $Q_{cano}^{modi}(\mathbf{L})$ and $Q_{cano}^{anal}(\mathbf{L})$ were also high: $C_{cor}^{normal} = 0.96$ and $C_{cor}^{log10} = 0.97$

because the central region is small compared with the whole region: $\frac{5^3}{11^3} = 0.094$. In

other words, the correlation coefficients obtained as presented above originated from

outside of the central region (shaded region in Supplementary Fig. S7).

Therefore, we re-calculated the correlation coefficients locally taking zones with $L^{(h)} = 3, \dots, 9$ ($\forall h$), as shown two-dimensionally by the region surrounded by the cyan-colored line in Supplementary Fig. S7. The volume ratio is: $\frac{5^3}{7^3} = 0.36$. The local correlation coefficients are denoted as B_{cor}^{normal} and $B_{cor}^{log^{10}}$. By this calculation, we can check not only the accuracy of $Q_{cano}^{[11]}(\mathbf{L})$ in the central region but also the connection of $Q_{cano}^{[11]}(\mathbf{L})$ between the central and outside regions. The resultant local correlations between $Q_{cano}^{modi}(\mathbf{L})$ and $Q_{cano}^{anal}(\mathbf{L})$ were $B_{cor}^{normal} = 0.74$ and $B_{cor}^{log^{10}} = 0.11$. In contrast, those between $Q_{cano}^{[11]}(\mathbf{L})$ and $Q_{cano}^{anal}(\mathbf{L})$ were $B_{cor}^{normal} = 0.98$ and $B_{cor}^{log^{10}} = 0.99$. This increment of B_{cor}^{normal} and $B_{cor}^{log^{10}}$ manifests that the central region (the poorly sampled region) was sampled appropriately using the SDIC method without breaking the entire shape of $Q_{cano}(\mathbf{L})$. This means that $Q_{cano}(\mathbf{L})$ was connected correctly between the central and outside regions.

Closing Remarks

Benefits of the subzone-based and GA-guided methods are the following: (1)

All snapshots from all iterations are useful for analysis: no production run is necessary,

or all simulations are production run. By contrast, in the original mD-VcMD method or other enhanced sampling methods, the production run is mandatory usually. Only the production run is used for analysis. (2) To proceed with sampling, no fine tuning of a weight function is used, which is also a benefit in comparison to multicanonical or other adaptive umbrella sampling methods. (3) The statistical weight w_i (Eq. 31) is found after sampling. The method used to compute w_i is simple. (4) Statistical significance of a poorly sampled region can be enhanced by selective sampling. In the original mD-VcMD method, the selective sampling accumulates local errors in $Q_{cano}(\mathbf{L})$, which results in slow convergence.

We applied the GA-guided and original mD-VcMD methods to the model system. The model system simplicity enables control of the sampling problem difficulty by tuning the height of the free-energy barrier (i.e., by varying the simulation temperature). At the higher temperature, both the original and GA-guided methods worked well, showing quick convergence of $Q_{cano}(\mathbf{L})$ to the analytical one. However, at the lower temperature, only the GA-guided method showed quick convergence. Furthermore, we demonstrated that the SDIC method worked well to provide $Q_{cano}(\mathbf{L})$

in the poorly sampled region without breaking the global shape of $Q_{cano}(\mathbf{L})$ in the RC space.

GA-guided mD-VcMD is applicable to any complicated biomolecular system for which the potential energy is defined for an MD simulation. We generated computer programs for the GA-guided mD-VcMD method, and applied them to molecular binding of a flexible medium-sized ligand (a peptide of about ten amino-acid residues long) to a receptor protein in an explicit solvent [9]. The resultant conformational ensemble produced a binding free-energy landscape in which various structural clusters (the unbound state, fragile encounter complexes, and the most stable native-like complex) are distributed. This result demonstrated that the GA-guided mD-VcMD simulation sampled the wide conformational space.

Conflict of Interest

All authors declare that they have no conflict of interest.

Author Contributions

All authors contributed equally to this study.

A preliminary version of this work was deposited in the arXiv (2006.06950v2) on August 13, 2020.

Acknowledgments

We deeply appreciate Prof. Haruki Nakamura from Osaka Univ. for insightful comments. J. H. was supported by JSPS KAKENHI Grant No. 16K05517 and by the Development of Core Technologies for Innovative Drug Development Based Upon IT from Japan Agency for Medical Research and Development (AMED). N. K. was supported by a JSPS KAKENHI Grant (No. JP20H03229). K. K. was supported by JSPS KAKENHI Grant No. 16K18526. This work was supported as an HPCI System Research Project (Project IDs: hp190017, hp190018, hp190027, hp200063, hp200090, and hp200025). Simulations were performed on the TSUBAME3.0 supercomputers at the Tokyo Institute of Technology, in part under the Cooperative Research Program of the Institute for Protein Research, Osaka University, CR-19-05 and CR-20-05.

References

- [1] Mitsutake, A., Sugita, Y. & Okamoto, Y. Generalized-ensemble algorithms for molecular simulations of biopolymers. *Biopolymers* **60**, 96–123 (2001). DOI: 10.1002/1097-0282(2001)60:2<96::AID-BIP1007>3.0.CO;2-F
- [2] Higo, J., Ikebe, J., Kamiya, N. & Nakamura, H. Enhanced and effective conformational sampling of protein molecular systems for their free energy landscapes. *Biophysical Rev.* **4**, 27–44 (2012). DOI: 10.1007/s12551-011-0063-6
- [3] Higo, J., Dasgupta, B., Mashimo, T., Kasahara, K., Fukunishi, Y. & Nakamura, H. Virtual-system-coupled adaptive umbrella sampling to compute free-energy landscape for flexible molecular docking. *J. Comput. Chem.* **36**, 1489–1501 (2015). DOI: 10.1002/jcc.23948
- [4] Higo, J., Kasahara, K., Dasgupta, B. & Nakamura, H. Enhancement of canonical sampling by virtual-state transitions. *J. Chem. Phys.* **146**, 044104 (2017). DOI: 10.1063/1.4974087
- [5] Paine, G. H. & Scheraga, H. A. Prediction of the native conformation of a polypeptide by a statistical-mechanical procedure. I. Backbone structure of enkephalin. *Biopolymers* **24**, 1391–1436 (1985). DOI: 10.1002/bip.360240802

- [6] Mezei, M. Adaptive umbrella sampling: Self-consistent determination of the non-Boltzmann bias. *J. Comput. Phys.* **68**, 237–248 (1987). DOI: 10.1016/0021-9991(87)90054-4
- [7] Hayami, T., Higo, J., Nakamura, H. & Kasahara, K. Multidimensional virtual-system coupled canonical molecular dynamics to compute free-energy landscapes of peptide multimer assembly. *J. Comput. Chem.* **40**, 2453–2463 (2019). DOI: 10.1002/jcc.26020
- [8] Higo, J., Kasahara, K., Wada, M., Dasgupta, B., Kamiya, N., Hayami, T., et al. Free-energy landscape of molecular interactions between endothelin 1 and human endothelin type B receptor: Fly casting mechanism. *Protein Engineering, Design & Selection (PESD)* **32**, 297–308 (2019). DOI: 10.1093/protein/gzz029
- [9] Higo, J., Kawabata, T., Kusaka, A., Kasahara, K., Kamiya, N., Fukuda, I., Mori, K., Hata, Y., Fukunishi, Y. & Nakamura, H. Molecular interaction mechanism of 14-3-3 ϵ protein with phosphorylated Myeloid leukemia factor 1 revealed by an enhanced conformational sampling. *J. Chem. Inf. Model.* **60**, 4867–4880 (2020). <https://doi.org/10.1021/acs.jcim.0c00551>

- [10] Higo, J., Kamiya, N., Sugihara, T., Yonezawa, Y. & Nakamura, H. Verifying trivial parallelization of multicanonical molecular dynamics for conformational sampling of a polypeptide in explicit water. *Chem. Phys. Lett.*, **473**, 326–329 (2009). DOI: 10.1016/j.cplett.2009.03.077
- [11] Ikebe, J., Umezawa, K., Kamiya, N., Sugihara, T., Yonezawa, Y., Takano, Y., Nakamura, H. & Higo, J. Theory for trivial trajectory parallelization of multicanonical molecular dynamics and application to a polypeptide in water. *J. Comput. Chem.*, **32**, 1286–1297 (2011). DOI: 10.1002/jcc.21710
- [12] Higo, J., Kasahara, K. & Nakamura, H. Multi-dimensional virtual system introduced to enhance canonical sampling. *J. Chem. Phys.*, **147**, 134102 (2017). DOI: 10.1063/1.4986129
- [13] Kasahara, K., Ma, B., Goto, K., Dasgupta, B., Higo, J., Fukuda, I., Mashimo, T., Akiyama, Y. & Nakamura, H. myPresto/omegagene: a GPU-accelerated Molecular Dynamics Simulator Tailored for Enhanced Conformational Sampling Methods with a Non-Ewald Electrostatic Scheme. *Biophysics and Physicobiology.*, **13**, 209–216 (2016). DOI: 10.2142/biophysico.13.0_209

Figure legends

Figure 1. 2D RC space is divided into zones. Panel (A) is equivalent to Supplementary Fig. S2C. Black arrows indicate directions to shift zones to remove inter-zone overlap (shaded region). (B) Extended RC space resulted from the zone shifts. L_i ($i = 1, \dots, 9$) and S_j ($j = 1, \dots, 4$) are the respective indices for zones and subzones. Shaded and checked subzones are described in the text.

Figure 2. Time development of sampled and non-sampled subzones in a zone along three (first, second and third) iterations. Four types of time development are presented as panels (A)–(D). “Average [1;3]” is $c^{[1;M]}(\mathbf{S}, \mathbf{L})$ (Eq. 5) over the three iterations ($M = 3$). Shaded and white subzones respectively represent sampled and non-sampled subzones. Checked subzones for “Zone correction” are corrected ones by Eq. 7 (procedure 2) for panel (C) and by Eq. 8 (procedure 3) for panel (D).

Figure 3. 2D RC space presented in extended form as Fig. 1B. Four subzones in red circles overlap to a single intersection. Six red lines represent six subzone pairs. Similarly, four subzones in blue circles overlap to another intersection with six subzone pairs indicated by blue lines. Green lines represent subzone pairs that are related to a zone ζ_{L_7} , as described in the text.

Figure 4. In both panels, the x-axis is presented by three measures: λ , L , and S . They are scalar because the RC space is 1D in this figure. (A) Open circles represent SS counts $c^{[1;M]}(L,S)$. Pairing subzones are connected by broken-line rectangles. Broken line represents the level of $c^{[1;M]}(L,S) = 1$. (B) Green-colored filled circles represent modulated SS counts $\rho^{[M]}(S_k, L_j; \sigma_L^{min})$. Red filled circles are $Q_{cano}^{[M]}(L)/2 = \{\rho^{[M]}(S_1, L_k; \sigma_{L_k}^{min}) + \rho^{[M]}(S_2, L_k; \sigma_{L_k}^{min})\}/2$. Pairing subzones, which are unified in broken-line rectangles, have the same modulated SS count.

Figure 5. (A) GA blocks in 2D RC space. A GA block, which consists of CS zones, is classified into three depending on the position of the central GA zone (marked with a red circle) in the RC space: blocks for which the central GA zones are at the side and corner of the RC space are called side and corner GA blocks, respectively. A block is called an inside GA block if the central GA zone is inside the RC space. (B) In 3D RC space, blocks of four types are possible: inside, face, side, and corner. The central GA zone for the inside GA block is hidden. Although this figure is presented for blocks in GA database, the same block classification is possible for recovery blocks and blocks in GA generation Gen_k .

Figure 6. (A) GA blocks and recovery blocks in 2D RC space. Six GA blocks $GA_1, \dots, GA_6$ are shown by frames of different colors. Its central GA zone is marked by a circle with the same color. CS, IS, and E zones in GA blocks are shown by rectangles of different tones. Broken-line frames represent recovery blocks. (B) Four recovery blocks ($REC_1, \dots, REC_4$) and their recovery zones (REC zone 1, ..., 4) are selected from panel (A). Although more recovery zones are possible in panel (A), they are not discussed herein. (C) Reset recovery blocks $REC_1, \dots$, and REC_4 .

Figure 7. GA process to update the k -th GA generation Gen_k to the $(k + 1)$ -th one Gen_{k+1} . Details for the process are explained in the text.

Figure 8. Part of 2D RC space for which the axes are $L^{(i)}$ (or $\lambda^{(i)}$) and $L^{(j)}$ (or $\lambda^{(j)}$). Three zone types are represented by different tones. Colored circles are REC zones described in text.

Figure 9. Image of slices in the 3D RC space.

Figure 10. Correlation coefficients, $C_{cor}^{normal}[M]$ and $C_{cor}^{log10}[M]$ as a function of iteration ordinal number M between analytical Q_{cano}^{anal} and the simulated $Q_{cano}^{[M]}$. Red and blue lines respectively represent C_{cor}^{log10} and C_{cor}^{normal} for all panels. (A) and (B)

Correlation coefficients between $Q_{cano}^{[M]}$ from the GA-guided method and Q_{cano}^{anal} , respectively, at $T = 1.5$ and 3.0 . (C) and (D) Those between $Q_{cano}^{[M]}$ from the original method and Q_{cano}^{anal} , respectively, at $T = 1.5$ and 3.0 .

Figure 11. Spatial patterns of $Q_{cano}(\mathbf{L})$ (i.e., dependence of Q_{cano} on $\mathbf{L}$) at $T = 1.5$.

Although three sides of the box are parallel to $L^{(x)}$ -, $L^{(y)}$ -, and $L^{(z)}$ -axes, because potential energy E_{model} has spherical symmetry, we do not specify the axes. (A) Patterns for analytical Q_{cano}^{anal} , (B) those from the 10-th iteration of the GA-guided method, (C) those from the third iteration of the original method, and (D) those from the 30-th iteration of the original method. Red and blue contours respectively correspond to $\log_{10}(Q_{cano}(\mathbf{L})) = -3$ and -9 , where Q_{cano} is normalized as $\sum_{\mathbf{L}} Q_{cano}(\mathbf{L}) = 1$.

Figure 1

296x209mm (300 x 300 DPI)

Figure 2

296x209mm (300 x 300 DPI)

Figure 3

296x209mm (300 x 300 DPI)

Figure 4

593x419mm (72 x 72 DPI)

Figure 5

296x209mm (300 x 300 DPI)

Figure 6
593x419mm (72 x 72 DPI)

Figure 7

296x209mm (300 x 300 DPI)

Figure 8

296x209mm (300 x 300 DPI)

Figure 9

296x209mm (300 x 300 DPI)

Figure 10
593x419mm (72 x 72 DPI)

Figure 11

593x419mm (72 x 72 DPI)

Supplementary text S1

Supplementary Table S1. Terminology

This table indicates where technical terms, parameters, and functions are defined in the main text or SI.

Term/quantity	Section
Reaction coordinate (RC)	Introduction of main text or section 1 of SI
Multi-dimensional (mD) RCs	Introduction of main text or section 2 of SI
Atom groups G_h^A and G_h^B ($h = \alpha, \beta, \gamma, \dots$)	Section 1 of SI (see Fig. S1)
$\lambda^{(h)}$ ($h = \alpha, \beta, \gamma, \dots$)	Section 1 of SI (see Fig. S1)
$\mathbf{R}$: system's conformation	Section 2 of SI
$\boldsymbol{\lambda} = [\lambda^{(\alpha)}(\mathbf{R}), \lambda^{(\beta)}(\mathbf{R}), \lambda^{(\gamma)}(\mathbf{R}), \dots]$	Section 2 of SI (see Fig. S2C)
$\zeta_i^{(h)}$ ($h = \alpha, \beta, \gamma, \dots; i = 1, \dots, n_{vs}(h)$)	Section 2 of SI (see Fig. S2A and S2B)
$n_{vs}(h)$	Section 2 of SI (see Fig. S2A)
$\Delta\lambda_i^{(h)}$ and $\Delta\lambda^{(h)}$	Section 2 of SI (see Fig. S2A)
$[\zeta_i^{(h)}]_{min}$ and $[\zeta_i^{(h)}]_{max}$	Section 2 of SI (see Fig. S2A and S2B)
$L^{(h)}$ and $\mathbf{L} = [L^{(\alpha)}, L^{(\beta)}, L^{(\gamma)}, \dots]$: zone index	Section 2 of SI (see Fig. S2C)
$\boldsymbol{\zeta}_L = \boldsymbol{\zeta}_{[L^{(\alpha)}, L^{(\beta)}, L^{(\gamma)}, \dots]} = [\zeta_{L^{(\alpha)}}^{(\alpha)}, \zeta_{L^{(\beta)}}^{(\beta)}, \zeta_{L^{(\gamma)}}^{(\gamma)}, \dots]$	Section 2 of SI (see Fig. S2C, and Eq. S1)
$\mathbf{L}$: virtual-state variable (or virtual-state)	Section 3 of SI (equivalent to zone index)
intersection	Section 3 of SI (see Fig. S2C)
$\boldsymbol{\zeta}_{L(1)}, \boldsymbol{\zeta}_{L(2)}, \dots$, and $\boldsymbol{\zeta}_{L(n_{link})}$: linked RC zones	Section 3 of SI (see Fig. S2C)
$\mathbf{L}_{(1)}, \dots, \mathbf{L}_{(n_{link})}$: linked virtual states	Sections 3 and 4 of SI
n_{link} : number of linked virtual states (zones)	Section 3 of SI (see Fig. S2C)
$E_{entire}(\mathbf{R}, \mathbf{L})$: potential energy	Section 3 of SI (see Eq. S2)
$E_R(\mathbf{R})$: original potential energy	Section 3 of SI (see Eq. S2)
$E_V(\mathbf{L})$	Section 3 of SI (see Eq. S5)
$E_{RV}(\boldsymbol{\lambda}(\mathbf{R}), \mathbf{L})$	Section 3 of SI (see also Eq. S3)
inter-virtual state transition (IVT) = inter-zone transition	Sections 3 and 4 of SI
$P_t(\mathbf{L} \rightarrow \mathbf{L}')$: formal IVT probability	Section 4 of SI (see Eq. S6)
$J_{L(i)}$: optimal IVT probability	Section 4 of SI (see Eq. 7)
configurational motion (CFM)	Section 4 of SI

$Q_{entire}(\mathbf{L})$: virtual state-partitioned probability	Section 4 of SI
$Q_{cano}(\mathbf{L})$: virtual state-partitioned canonical probability	Section 4 of SI
$\mathbf{S}_j$: subzone index	Section 1.1 of main text (see Fig. 1B)
$c^{[M]}(\mathbf{S}, \mathbf{L})$: subzone's snapshot count (SS count)	Section 1.1 of main text
empty zone (E zone)	Section 1.1 of main text
completely-sampled zone (CS zone)	Section 1.1 of main text
incompletely sampled zone (IS zone)	Section 1.1 of main text
n_{is}	Section 1.2 of main text
subzone pair	Section 1.2 of main text
$n_{pair}(j)$	Section 1.2 of main text
modulated SS count	Section 1.2 of main text
$p_{k,j}^{(1)}$ and $p_{k,j}^{(2)}$: pairing subzone indices	Section 1.2 of main text
$q_{k,j}^{(1)}$ and $q_{k,j}^{(2)}$: pairing zone indices	Section 1.2 of main text
$[\mathbf{S}_{p_{k,j}^{(1)}}, \mathbf{L}_{q_{k,j}^{(1)}}]$ and $[\mathbf{S}_{p_{k,j}^{(2)}}, \mathbf{L}_{q_{k,j}^{(2)}}]$:	Section 1.2 of main text
σ_L and σ : multiplication factor to SS count	Section 1.2 of main text
F : objective function to be minimized	Section 1.2 of main text
σ_L^{min} and σ^{min} : multiplication factor of minimize F	Section 1.2 of main text
GA database	Section 2.1 of main text
GA block	Section 2.1 of main text
central GA zone	Section 2.1 of main text
inside, face, side, or corner GA block	Section 2.1 of main text (see Fig. 5)
GA_j : the j -th GA block	Section 2.1 of main text (see Fig. 6)
Λ : relative position of a zone in a block	Section 2.1 of main text
$[\mathbf{S}, \Lambda]$	Section 2.1 and section 2.2 of main text
$c^{GA_i}(\mathbf{S}, \Lambda)$	Section 2.1 of main text
N_{surr}	Section 2.2 of main text
recovery zone (REC zone)	Section 2.2 of main text
recovery block (REC block)	Section 2.2 of main text
REC_i : the i -th REC block	Section 2.2 of main text
inside, face, side, and corner REC blocks	Section 2.2 of main text (see Fig. 5)
$c^{REC_i}(\mathbf{S}, \Lambda)$	Section 2.3 of main text
$c_{cor}(REC_i, GA_j)$: correlation coefficient between REC_i and GA_j	Section 2.4 of main text

$E_{simi}(REC_i, GA_j)$: score function	Section 2.4 of main text
$P(\sigma_L, \sigma_{L'})$	Section 2.5 of main text
E_{phys}	Section 2.5 of main text
Gen_k : the k -th generation	Sections 2.6 and 2.7 of main text (see Fig. 7)
$E_{score}(REC_i, GA_j)$	Section 2.6 of main text
N_{GA}^{all} : number of members in a generation	Section 2.6 of main text
N_{GA}^{top}	Section 2.7 of main text (see Fig. 7)
N_{GA}^{mut}	Section 2.7 of main text
N_{GA}^{exc}	Section 2.7 of main text
$MEM_k^{(1)}, MEM_k^{(2)}, \dots, MEM_k^{(N_{GA}^{mut})}$	Section 2.7 of main text
$GA^{(1)}, GA^{(2)}, \dots, GA^{(N_{GA}^{mut})}$	Section 2.7 of main text
mutation	Section 2.7 of main text
crossover/exchange	Section 2.7 of main text
EDIC (evenly-distributed-initial-conformations)	Section 3.1 of main text
SDIC (Selectively distributed initial conformations)	Section 3.2 of main text
$n_{snap}^{[1:M]}(\mathbf{L})$	Section 4 of main text

1. Definition of reaction coordinates

Consider two atom groups G_h^A and G_h^B ($h = \alpha, \beta, \gamma, \dots$) in a molecular system. A reaction coordinate (RC), $\lambda^{(h)}$, is defined by the distance between centers of mass of G_h^A and G_h^B (Supplementary Fig. S1). Superscripts A and B indicate simply that the two atom groups are pairing to define $\lambda^{(h)}$, and then, one can exchange the superscripts as: $G_h^A \rightarrow G_h^B$ and $G_h^B \rightarrow G_h^A$ without changing the value of $\lambda^{(h)}$. The RC can be defined between two different molecules as in Supplementary Fig. S1A or between two portions in a molecule (Supplementary Fig. S1B). In Supplementary Fig. S1C, the atom group G_h^A is composed of two different portions in a molecule. As shown in this figure, there are various ways to define atom groups. This means that the current method is applicable to various problems (folding and binding).

Supplementary Figure S1. Two atom groups, G_h^A and G_h^B , are indicated by red-colored and blue-colored rectangles, respectively. Atoms in G_h^A and G_h^B are presented by small black filled circles. Center of mass of each atom group is presented by a cross. The distance between the two centers of mass is $\lambda^{(h)}$ (broken-line with arrows).

2. Division of RC space and definition of virtual zones

In this section, we use a one-dimensional (1D) or two-dimensional (2D) RC space for explanation. Outcomes from the simplified RC space are extendable readily to those for a multi-dimensional (mD) RC space.

The RCs are arbitrary but should be computable uniquely from the system's conformation $\mathbf{R}$ as $\lambda^{(h)} = \lambda^{(h)}(\mathbf{R})$ ($h = \alpha, \beta, \gamma, \dots$), where $\mathbf{R} = [x_1, y_1, z_1, x_2, y_2, z_2, \dots]$ and x_i , y_i , and z_i are the x-, y-, and z-coordinates of atom i , respectively. Then, we express the RCs in a vector as $\boldsymbol{\lambda} = [\lambda^{(\alpha)}, \lambda^{(\beta)}, \lambda^{(\gamma)}, \dots] = [\lambda^{(\alpha)}(\mathbf{R}), \lambda^{(\beta)}(\mathbf{R}), \lambda^{(\gamma)}(\mathbf{R}), \dots]$. According to the motion of $\mathbf{R}$, $\boldsymbol{\lambda}$ moves in the mD RC space.

We divide each 1D RC axis into zones (called *RC zones* or simply *zones*) $\{\zeta_1^{(h)}, \dots, \zeta_{n_{vs}(h)}^{(h)}\}$ as shown in Supplementary Fig. S2A, where $\zeta_i^{(h)}$ is the i -th zone along the $\lambda^{(h)}$ -axis ($i = 1, \dots, n_{vs}(h)$), $n_{vs}(h)$ is the number of zones set to $\lambda^{(h)}$, and $\Delta\lambda_i^{(h)}$ is the width of the i -th zone. Usually we set $\Delta\lambda_i^{(h)} = \text{const}$ ($\forall i$), although $\Delta\lambda_i^{(h)}$ and $\Delta\lambda_j^{(h)}$ can be different for $i \neq j$ in general. Thus, we denote simply the width as $\Delta\lambda^{(h)}$ ($= \Delta\lambda_1^{(h)} = \dots = \Delta\lambda_{n_{vs}(h)}^{(h)}$) in this paper. The case of $\Delta\lambda_i^{(h)} \neq \Delta\lambda_j^{(h)}$ is discussed in ref. 1. The lower and upper bounds of $\zeta_i^{(h)}$ are denoted as $[\zeta_i^{(h)}]_{\min}$ and $[\zeta_i^{(h)}]_{\max}$, respectively: $[\zeta_i^{(h)}]_{\min} \leq \lambda^{(h)} \leq [\zeta_i^{(h)}]_{\max}$ if $\lambda^{(h)} \in \zeta_i^{(h)}$. The entire sampling range for $\lambda^{(h)}$ is $[\zeta_1^{(h)}]_{\min}$ to $[\zeta_{n_{vs}(h)}^{(h)}]_{\max}$. Importantly, the adjacent zones

$\zeta_i^{(h)}$ and $\zeta_{i\pm 1}^{(h)}$ overlap to each other, whereas $\zeta_{i-1}^{(h)}$ and $\zeta_{i+1}^{(h)}$ do not. This overlap is essentially important for inter-zone transitions as explained later.

Supplementary Figure S2B illustrates division of a 2D RC space constructed by $\lambda^{(\alpha)}$ and $\lambda^{(\beta)}$ into 2D zones. A 2D zone generated by the i -th and j -th 1D zones along $\lambda^{(\alpha)}$ and $\lambda^{(\beta)}$, respectively, is expressed as $[\zeta_i^{(\alpha)}, \zeta_j^{(\beta)}]$. For instance, the 2D zone enclosed by the broken line in Supplementary Fig. S2B is expressed as $[\zeta_2^{(\alpha)}, \zeta_3^{(\beta)}]$. A higher dimensional RC space can be divided similarly: In the m D RC space constructed by $\lambda^{(\alpha)}$, $\lambda^{(\beta)}$, and $\lambda^{(\gamma)}$, a m D zone is expressed as $[\zeta_i^{(\alpha)}, \zeta_j^{(\beta)}, \zeta_k^{(\gamma)}, \dots]$, where $\zeta_k^{(\gamma)}$ is the k -th 1D zone along $\lambda^{(\gamma)}$. Now, we introduce an index parameter $L^{(h)}$ ($h = \alpha, \beta, \gamma, \dots$) to specify an RC zone along $\lambda^{(h)}$. A 2D index, $\mathbf{L} = [L^{(\alpha)}, L^{(\beta)}]$, is used to specify a 2D zone. In Supplementary Fig. S2B, for instance, the index for the 2D zone enclosed by the broken line is presented as $\mathbf{L} = [2, 3]$. Similarly, a m D index parameter is given as $\mathbf{L} = [L^{(\alpha)}, L^{(\beta)}, L^{(\gamma)}, \dots]$.

Using the index parameter, a 2D zone is expressed as a vector: $\boldsymbol{\zeta}_{\mathbf{L}} = \boldsymbol{\zeta}_{[L^{(\alpha)}, L^{(\beta)}]} = [\zeta_{L^{(\alpha)}}^{(\alpha)}, \zeta_{L^{(\beta)}}^{(\beta)}]$. Similarly, a m D zone is expressed as:

$$\boldsymbol{\zeta}_{\mathbf{L}} = \boldsymbol{\zeta}_{[L^{(\alpha)}, L^{(\beta)}, L^{(\gamma)}, \dots]} = [\zeta_{L^{(\alpha)}}^{(\alpha)}, \zeta_{L^{(\beta)}}^{(\beta)}, \zeta_{L^{(\gamma)}}^{(\gamma)}, \dots], \quad (\text{S1})$$

where $\zeta_{L^{(h)}}^{(h)}$ represents the $L^{(h)}$ -th RC zone along $\lambda^{(h)}$. For instance, an index with $L^{(\alpha)} = i$, and $L^{(\beta)} = j$ in the 2D RC space is presented as $\mathbf{L} = [i, j]$, and the corresponding 2D zone is $\boldsymbol{\zeta}_{\mathbf{L}} = \boldsymbol{\zeta}_{[i, j]} = [\zeta_i^{(\alpha)}, \zeta_j^{(\beta)}]$.

Supplementary Figure S2. Schematic illustration of RC space. (A) Division of a 1D RC axis $\lambda^{(h)}$, where $h = \alpha, \beta$, or γ , into RC zones $\{\zeta_i^{(h)}\}$ ($i = 1, \dots, n_{vs}(h)$), where $\Delta\lambda_i^{(h)}$ is the width of the i -th zone, and $n_{vs}(h)$ is the number of zones along $\lambda^{(h)}$. The lower and upper bounds of the i -th zone are expressed respectively as $[\zeta_i^{(h)}]_{\min}$ and $[\zeta_i^{(h)}]_{\max}$. (B) Division of 2D RC space constructed by $\lambda^{(\alpha)}$ and $\lambda^{(\beta)}$ into 2D zones. Periphery and corner regions are designated respectively as “phr” and “cnr”. Broken-line frame is mentioned in text. (C) Close-up of 2D RC space, where four zones of differently colored frames overlap in the shaded region, which is called an intersection. System’s conformation is moving from λ to λ' . Colored indices $L_{(1)}$, $L_{(2)}$, $L_{(3)}$, and $L_{(4)}$ are mentioned in text. A color used for a character and frame correspond to each other. For instance, $L_{(4)}$ specifies the red-colored zone. Shaded region is RC-zone intersection.

3. Potential energy and mD-VcMD

We set walls at the zone boundaries to confine λ in a zone and allow inter-zone transitions occasionally during a simulation. Figure 2C is a portion of the 2D RC space, where the current zone is surrounded by red-colored frame. The shaded region (called *intersection*) in Fig. 2C is shared by four zones, which are referred to as *linked RC zones*, and expressed as $\zeta_{L_{(1)}}$ (cyan-colored frame), $\zeta_{L_{(2)}}$ (bright yellow), $\zeta_{L_{(3)}}$ (brown), and $\zeta_{L_{(4)}}$ (red). The indices $L_{(1)}$, $L_{(2)}$, $L_{(3)}$, and $L_{(4)}$ are those for the

linked zones. Later, we renamed $\{\mathbf{L}_{(i)}\}$ ($i = 1, \dots$) as linked virtual states. In general, we express the linked zones sharing an intersection as: $\mathbf{L}_{(1)}, \dots, \mathbf{L}_{(n_{link})}$, where n_{link} is the number of the linked RC zones for the intersection. The value of n_{link} depends on the RC-space dimensionality and the position of the intersection in the RC space. For instance, in Fig. 2C, $n_{link} = 4$ if the intersection is inside of the 2D RC space, although $n_{link} = 2$ for intersections at periphery (“phr” in Fig. 2B), and $n_{link} = 1$ for intersections at corner (“cnr”). For a higher-dimensional RC space, n_{link} increases. If the current RC zone is $\boldsymbol{\zeta}_{\mathbf{L}_{(4)}}$ in Fig. 2C, then $\boldsymbol{\lambda}$ moves within the red-colored frame. Members of the linked RC zones change according to the motion: When $\boldsymbol{\lambda}$ moves to $\boldsymbol{\lambda}'$ in $\boldsymbol{\zeta}_{\mathbf{L}_{(4)}}$, then $\boldsymbol{\zeta}_{\mathbf{L}_{(1)}}$ and $\boldsymbol{\zeta}_{\mathbf{L}_{(2)}}$ are eliminated from the linked RC zones, and other two zones (blue- and green-colored framed zones) join in the linked RC zones.

We defined the potential energy E_{entire} of the system as

$$E_{entire}(\mathbf{R}, \mathbf{L}) = E_R(\mathbf{R}) + E_V(\mathbf{L}) + E_{RV}(\boldsymbol{\lambda}(\mathbf{R}), \mathbf{L}), \quad (\text{S2})$$

where $\mathbf{R}$ and $\mathbf{L}$ are dynamic variables to specify E_{entire} . Therefore, we rename the current-zone index $\mathbf{L}$ a *virtual-state variable* (or simply, *virtual state*) to express that the system’s motion is controlled by varying $\mathbf{L}$. The term $E_R(\mathbf{R})$ is the original potential energy described by the system’s conformation $\mathbf{R}$, $E_{RV}(\boldsymbol{\lambda}, \mathbf{L})$ is introduced to confine $\boldsymbol{\lambda}$ in the current RC zone $\boldsymbol{\zeta}_{\mathbf{L}}$, and $E_V(\mathbf{L})$ controls a transition of the current zone from $\boldsymbol{\zeta}_{\mathbf{L}}$ to $\boldsymbol{\zeta}_{\mathbf{L}'}$.

The form of E_{RV} is a well-type potential expressed as:

$$E_{RV}(\boldsymbol{\lambda}; \mathbf{L}) = \begin{cases} 0 & (\text{for } \boldsymbol{\lambda} \in \boldsymbol{\zeta}_{\mathbf{L}}) \\ c_{RV} \sum_h \left(\delta_{\mathbf{L}^{(h)}}^{(h)} \right)^2 & (\text{for } \boldsymbol{\lambda} \notin \boldsymbol{\zeta}_{\mathbf{L}}) \end{cases} \quad (\text{S3})$$

where

$$\delta_{\mathbf{L}^{(h)}}^{(h)} = \begin{cases} \lambda^{(h)} - [\zeta_{\mathbf{L}^{(h)}}^{(h)}]_{min} & (\text{for } \lambda^{(h)} < [\zeta_{\mathbf{L}^{(h)}}^{(h)}]_{min}) \\ \lambda^{(h)} - [\zeta_{\mathbf{L}^{(h)}}^{(h)}]_{max} & (\text{for } \lambda^{(h)} > [\zeta_{\mathbf{L}^{(h)}}^{(h)}]_{max}) \end{cases}. \quad (\text{S4})$$

The summation for h in Eq. S3 is taken over $\alpha, \beta, \gamma, \dots$. Remember that $[\zeta_{L^{(h)}}^{(h)}]_{min}$ and $[\zeta_{L^{(h)}}^{(h)}]_{max}$ are the lower and upper boundaries of the $L^{(h)}$ -th RC zone along the $\lambda^{(h)}$ axis, respectively. The restoring force, $-\nabla E_{RV}$, acts on λ only when λ is outside the zone ζ_L , and the coefficient c_{RV} is set to a large value so that the restoring force increases rapidly outside ζ_L . Usually we set c_{RV} to 100 kcal/mol. We count the number of snapshots detected in each RC zone. However, we eliminate the outside-the-zone conformations from the count. Therefore, the effect of the artificial force (i.e., restoring force) is removed completely in the statistical analyses.

We define E_V as follows:

$$E_V(L) = g_L, \quad (S5)$$

where g_L is a constant in ζ_L independent of λ and R , but varies when L moves to another zone L' : $g_L \neq g_{L'}$ if $L \neq L'$. Because L varies discretely, we use a Monte-Carlo method to move L , which means that E_V is related to transition probabilities among virtual states.

4. Inter-virtual state transition probability

When an inter-zone transition is applied to the system at a time t ($= t_0 + n \Delta\tau$), where t_0 is the initial time of simulation and n is an integer. In other words, R is confined in a current zone ζ_L for a time interval of $[t_0 + (n-1)\Delta\tau; t_0 + n\Delta\tau]$. Now, we explain the simulation procedure concretely: In a time interval $[t_0, t_0 + \Delta\tau]$, the *configurational motion* (CFM); the motion of R is performed with an MD protocol where forces acting on atoms are derived by $-\nabla E_{entire}$ with fixing the current virtual state at L (i.e., the current RC zone is ζ_L). Suppose that λ is in an intersection shared by n_{link} linked RC zones $\zeta_{L(1)}, \zeta_{L(2)}, \dots$, and $\zeta_{L(n_{link})}$ at time $t_0 + \Delta\tau$, and note that the current RC zone ζ_L is one of the n_{link} linked RC zones. The virtual states whose zones are linked are called *linked virtual states*. Then, *inter-virtual state transition* (IVT) from L to L' (i.e., L' is also one of $L(1), \dots, L(n_{link})$) is executed at time $t_0 + \Delta\tau$ by selecting one of the n_{link} linked RC zones with using a transition probability between L and L' . If $\zeta_{L'} = \zeta_L$, this transition is a self-transition. Note that R and λ are fixed when executing IVT. In the next interval $[t_0 + \Delta\tau, t_0 + 2\Delta\tau]$, CFM is performed using E_{entire} with fixing L' (i.e., λ moves in $\zeta_{L'}$). Then, at time $t_0 +$

$2\Delta\tau$, IVT is achieved from $\mathbf{L}'$ to $\mathbf{L}''$ with fixing $\mathbf{R}$ and $\boldsymbol{\lambda}$, and so on. Although the interval $\Delta\tau$ can be set arbitrary in theory, we usually set $\Delta\tau$ to a constant value (say 10 ps).

Needless to say, the transition probability controls the sampling efficiency. Here, we present an optimal set of the transition probability. First, the formal expression for the transition probability is given as:

$$\begin{aligned} P_t(\mathbf{L} \rightarrow \mathbf{L}') &= \exp[-\beta\Delta E_{entire}] \\ &= \exp[-\beta(E_{entire}(\mathbf{R}, \mathbf{L}') - E_{entire}(\mathbf{R}, \mathbf{L}))] \\ &= \exp[-\beta\Delta E_V] \end{aligned} \quad (\text{S6})$$

where $P_t(\mathbf{L} \rightarrow \mathbf{L}')$ is the *inter-virtual state transition probability (IVT probability)* from $\mathbf{L}$ to $\mathbf{L}'$ with fixing $\mathbf{R}$ and $\boldsymbol{\lambda}$ at t , and $\Delta E_V = E_V(\mathbf{L}') - E_V(\mathbf{L})$. Note that $\boldsymbol{\zeta}_{\mathbf{L}}$ and $\boldsymbol{\zeta}_{\mathbf{L}'}$ are zones sharing an intersection, which involves $\boldsymbol{\lambda}$. The terms $E_R(\mathbf{R})$ and $E_{RV}(\boldsymbol{\lambda}, \mathbf{L})$ are canceled out in Eq. S6: $\Delta E_R = \Delta E_{RV} = 0$. Therefore, $P_t(\mathbf{L} \rightarrow \mathbf{L}')$ is controlled only by E_V . The term “inter-zone transition” used in the above sections is replaced by “inter-virtual state transition” here. Equation 6 is a formal expression for IVT transitions, where E_V (or $g_{\mathbf{L}}$) is not derived from a physicochemical law but can be set according to our purpose. This is why we can modulate $P_t(\mathbf{L} \rightarrow \mathbf{L}')$ to increase sampling efficiency.

We introduce a quantity named *virtual state-partitioned probability* $Q_{entire}(\mathbf{L})$, which is the number of snapshots detected in the zone $\boldsymbol{\zeta}_{\mathbf{L}}$ in the simulation. Suppose that the system is at $\boldsymbol{\lambda}$ in an intersection at a time and that its virtual state is $\mathbf{L}_{(i)}$, which is one of linked virtual zones $\{\mathbf{L}_{(1)}, \dots, \mathbf{L}_{(n_{link})}\}$. Then, one count of detection is added to $Q_{entire}(\mathbf{L}_{(i)})$ not to those of the other linked virtual states. Furthermore, we introduce another quantity $Q_{cano}(\mathbf{L})$, called a *virtual state-partitioned canonical probability* [1,2]. If we perform a long mD-VcMD simulation with setting all IVT probabilities to a constant (i.e., $P_t(\mathbf{L} \rightarrow \mathbf{L}') = \text{const}$ for $\forall \mathbf{L}$ and $\forall \mathbf{L}'$), then the resultant $Q_{entire}(\mathbf{L})$ is $Q_{cano}(\mathbf{L})$. Thus, $Q_{cano}(\mathbf{L})$ is a special case of $Q_{entire}(\mathbf{L})$. However, such a simulation takes considerably long computation time for convergence of $Q_{cano}(\mathbf{L})$. Last in this paragraph, we note that $Q_{entire}(\mathbf{L})$ and $Q_{cano}(\mathbf{L})$ are not continuous but discrete function of $\mathbf{L}$ because the zone index is a discrete quantity.

Determination of $Q_{cano}(\mathbf{L})$ is essentially important because we can define the IVT probability using $Q_{cano}(\mathbf{L})$. Ref. [3] showed that sampling enhancement is realized when $Q_{entire}(\mathbf{L}) \approx \text{const}$ is obtained during sampling, and Supporting Information of Ref. 1 presents that the IVT probability is given as:

$$J_{L(i)} = \left[Q_{cano}(\mathbf{L}_{(i)}) \sum_{m=1}^{n_{link}} \frac{1}{Q_{cano}(\mathbf{L}_{(m)})} \right]^{-1}. \quad (\text{S7})$$

In general, the current virtual zone is one of linked zones $\zeta_{L(1)}, \zeta_{L(2)}, \dots, \zeta_{L(n_{link})}$. Then, $J_{L(i)}$ provides an IVT probability from any of the linked virtual states to $\mathbf{L}_{(i)}$. If λ is in the corner zone (see Fig. 2B), n_{link} may be 1, and then, the only self-transition is allowed. We note that Eq. S6 is a formal expression for the IVT probability and Eq. S7 is the actually used one to optimize the sampling.

One may consider that another expression, $E_V(\mathbf{L}) = -RT \ln[Q_{cano}(\mathbf{L})]$, can result in $Q_{entire}(\mathbf{L}) \approx \text{const}$. Equation S7 is another expression to obtain $Q_{entire}(\mathbf{L}) \approx \text{const}$ with a small rejection rate in IVT [3].

Whereas Eq. 7 requires a set of $Q_{cano}(\mathbf{L})$, it is difficult to obtain accurate $Q_{cano}(\mathbf{L})$ directly from a simulation with setting $P_t(\mathbf{L} \rightarrow \mathbf{L}') = 1$ in a realistic time. Instead, we use a method to convert $Q_{entire}(\mathbf{L})$ to $Q_{cano}(\mathbf{L})$ via iterative mD-VcMD simulations. Section 1 of SI explains briefly the “original mD-VcMD method”[1,2] for the conversion. Then, below we present two methods: “a subzone-guided mD-VcMD method”, which is an extension of the original method, and a “GA-guided mD-VcMD method”, which is introduced to expand sampling to un-sample regions through iterative simulation.

5. Original method to compute $Q_{cano}(\mathbf{L})$

Assume that we have performed the M -th iteration of mD-VcMD. Then, $Q_{cano}(\mathbf{L})$ is set as [1,2]:

$$Q_{cano}^{[M]}(\mathbf{L}) = Q_{cano}^{[M-1]}(\mathbf{L}) Q_{entire}^{[M]}(\mathbf{L}), \quad (\text{S8})$$

where $Q_{cano}^{[M-1]}(\mathbf{L})$ is $Q_{cano}(\mathbf{L})$ obtained from the $(M-1)$ -th iteration, and $Q_{entire}^{[M]}(\mathbf{L})$ is the resultant $Q_{entire}(\mathbf{L})$ calculated numerically from the M -th iteration (i.e., the currently done simulation). From $Q_{cano}^{[M]}(\mathbf{L})$ computed by Eq. S8, $J_{L(i)}$ is calculated (Eq. S7), and the $(M+1)$ -th iteration is performed. This iteration is continued till a relation $Q_{entire}^{[M]}(\mathbf{L}) \approx \text{const} (\forall \mathbf{L})$ is satisfied approximately [1,2]. Because $Q_{cano}^{[0]}(\mathbf{L})$ is not prepared in advance for the first iteration, we assume $Q_{cano}^{[0]}(\mathbf{L}) = \text{const} (\forall \mathbf{L})$ and $Q_{cano}^{[1]}(\mathbf{L}) = Q_{entire}^{[1]}(\mathbf{L})$ as a guess.

Supplementary Figure S3. Flow chart of the GA-guided mD-VcMD method

Supplementary Figure S3. Schematic illustration of GA-guided mD-VcMD. Given a system, RCs are set, and the multi-dimensional (mD) RC space is divided into zones. Then, the zones are divided into subzones. Next, iteration M (current iteration) is performed, and $Q_{entire}^{[M]}(\mathbf{L})$ is obtained. After that, GA blocks are generated using data from iterations 1 to M . The ensemble of GA blocks is GA database. Sampled zones from iteration M are classified into three: CS, IS, and E zones. Then, REC zones and REC blocks are detected from the patterns of spatial patterns of CA, IC, and E zones. Next, the first GA procedure is done with comparing the spatial patterns of the SS counts between REC blocks and GA blocks, from which SS counts of the REC zones are determined. Then, new REC zones are detected, and the second GA procedure is done to determine the new REC zones, and so on. When the SS counts of all REC zones are determined, the multiple GA procedure is finished. Resultant SS counts are used to

calculate $Q_{entire}^{[M]}(\mathbf{L})$, from which IVT probability $J_{L(i)}^{[M]}$ is calculated. Iteration $M + 1$ is performed using $J_{L(i)}^{[M]}$, and so on.

Supplementary Figure S4. EDIC Method

Supplementary Figure S4. EDIC method is explained using a 2D RC space (the $\lambda^{(\alpha)}$ - $\lambda^{(\beta)}$ plane) for convenience. (A) Distribution of the last conformations (black filled circles) of runs from the first to M -th iterations, which consists of $N_{run} \times M$ conformations. The space is divided into small patches of an equal size. Shaded patches, called occupied patches, are those involving one or more conformations. (B) After randomly picking up N_{run} patches from the occupied ones, one conformation is selected from each of the picked patch, which are shown by red-colored filled circles.

Supplementary Figure S5. Potential energy for the simple model

Supplementary Figure S5. (A) Potential energy function $E_{\text{model}}(r)$ (Eq. 32 of main text) with respect to r for the simple model. The r is the distance from the origin of the space: $r = [x^2 + y^2 + z^2]^{1/2}$. Shown upper range, 5.2, for r is the limit for r , because the ranges for x , y , and z are: $-3.0 \leq x \leq 3.0$, $-3.0 \leq y \leq 3.0$, and $-3.0 \leq z \leq 3.0$. (B) $P_{\text{model}}(r)$ is the Boltzmann factor of a microscopic state at r : $P_{\text{model}}(r) = \exp [-E_{\text{model}}(r)/T]$ at two temperatures $T = 1.5$ (red-colored line) and $T = 3.0$ (blue-colored line). $P_{\text{model}}(r)$ is normalized so that the largest probability is set to 1.0.

Supplementary Figure S6. Potential surface in 2D space

Supplementary Figure S6. Shown is a cross section of $E_{\text{model}}(r)$ on the x - y plane ($z = 0$). The x - and y axes correspond to $\lambda^{(\alpha)}$ and $\lambda^{(\beta)}$, respectively, and the z -axis does to the energy axis to represent the value of E_{model} .

Supplementary Figure S7. Scheme for the SDIC method

Supplementary Figure S7. RC space presented two-dimensionally. The coordinate axes $L^{(x)}$ and $L^{(y)}$ are divided into 11 virtual zones (see Table S2 of SI) in the extended form. The white region is the central region,

and the shaded region is the outside of the central region. Red-colored filled circle is the position of the origin. Region surrounded by cyan-colored line is that used for the local correlation coefficients B_{cor}^{normal} and B_{cor}^{log10} .

Supplementary Table S2. Virtual zones

Table 1. Virtual zones $\zeta_i^{(h)}$ ($i = 1, \dots, 11$; $h = x, y, z$)

i	range for $\zeta_i^{(h)}$
1	[−3.0, −2.0]
2	[−2.5, −1.5]
3	[−2.0, −1.0]
4	[−1.5, −0.5]
5	[−1.0, 0.0]
6	[−0.5, 0.5]
7	[0.0, 1.0]
8	[0.5, 1.5]
9	[1.0, 2.0]
10	[1.5, 2.5]
11	[2.0, 3.0]

References

- [1] Higo, J., Kasahara, K., Wada, M., Dasgupta, B., Kamiya, N., Hayami, T., *et al.* Free-energy landscape of molecular interactions between endothelin 1 and human endothelin type B receptor: Fly-casting mechanism. *Protein Engineering, Design & Selection (PESD)* **32**, 297–308 (2019). DOI: 10.1093/protein/gzz029
- [2] Hayami, T., Higo, J., Nakamura, H. & Kasahara, K. Multidimensional virtual-system coupled canonical molecular dynamics to compute free-energy landscapes of peptide multimer assembly. *J Comput. Chem.* **40**, 2453–2463 (2019). DOI: 10.1002/jcc.26020

[3] Higo, J., Kasahara, K., Dasgupta, B. & Nakamura, H. Enhancement of canonical sampling by virtual-state transitions. *J. Chem. Phys.* **146**, 044104 (2017). DOI: 10.1063/1.4974087

Graphical Abstract

296x209mm (300 x 300 DPI)