Lacunary-Substituted Undecatungstosilicates Sensitize Methicillin-Resistant *Staphylococcus aureus* to β -Lactams

Yutaka TAJIMA*

Clinical Laboratory, Saga Medical School Hospital, Nabeshima, Saga 849–8501, Japan.

Received February 21, 2001; accepted June 18, 2001

Some Keggin-structural (hetero)polyoxotungstates with a lacunary hole, such as undecatungstosilicate ($[\text{SiW}_{11}\text{O}_{39}]^{8-}$, SiW_{11}), greatly sensitize methicillin-resistant *Staphylococcus aureus* (MRSA) to β -lactams. In this study, the effects of lacunary-substituted derivatives of SiW_{11} were tested to determine if the lacunary hole is required for the sensitizing effect. Initially, it was supposed that the hole functioned as a binding site that interacts with some bacterial components. However, most of the lacunary-substituted species had stronger effects than SiW_{11} , indicating that the hole is not essential for the sensitizing effect. Undecatungsto(ferri)ferrosilicate ($[\text{SiW}_{11}\text{O}_{39}\text{Fe}]^{6(5-)}$) was the most potent compound tested, but undecatungstocobaltosilicate ($[\text{SiW}_{11}\text{O}_{39}\text{Co}]^{6-}$) gave better results according to the fractional inhibitory concentration index. Small lipophilic cations enhanced the sensitizing effect, but bulky cations inhibited it. This suggests that a charge interaction between the tungsten compounds and some unknown materials in the culture medium or bacterial cells plays an important role in the sensitizing effect.

Key words *Staphylococcus aureus*; methicillin-resistant; β -lactam; tungstosilicate; lacunary hole

In a previous study, the author found a factor (called “Factor T”) in aged mixtures of tungstate and phosphate, and found that this factor greatly sensitizes methicillin-resistant *Staphylococcus aureus* (MRSA) to β -lactams.^{1–3} Although Factor T alone does not show significant antibacterial activity, minimum inhibitory concentrations (MICs) of various β -lactams to MRSA strains are reduced to be 1/10–1/1000 in the presence of the Factor T.^{1,2}

Subsequently, Factor T was purified and identified as undecatungstophosphate ($[\text{PW}_{11}\text{O}_{39}]^{7-}$, PW_{11}).⁴ This is one of the compounds called “Keggin”-structural (hetero)polyoxotungstates,^{5–7} which can be easily prepared by partial hydrolysis of dodecatungstophosphate ($[\text{PW}_{12}\text{O}_{40}]^{3-}$, PW_{12}).⁴ Undecatungstosilicate ($[\text{SiW}_{11}\text{O}_{39}]^{8-}$, SiW_{11}), a compound closely related to PW_{11} , has a similar sensitizing effect on MRSA strains.⁴

Initially, the author used PW_{11} for experiments, but recently, SiW_{11} has been used, for several reasons:

(1) The molecular structure and preparation procedure of SiW_{11} are almost the same as those of PW_{11} .^{5,6} The differences are that (a) the core part of the SiW_{11} molecule is silicate instead of phosphate (Fig. 1), (b) the anionic valency is increased from -7 to -8 , and (c) the starting material is dodecatungstosilicate ($[\text{SiW}_{12}\text{O}_{40}]^{4-}$, SiW_{12}) instead of PW_{12} .

(2) The sensitizing effect of SiW_{11} is almost equivalent to that of PW_{11} , but SiW_{11} has better pH-stability than PW_{11} : Around 10% of SiW_{11} is decomposed with every increase of 0.1 pH unit from pH 8.0 (completely hydrolyzed over pH 9), whereas PW_{11} starts to decompose from pH 7.0.⁷ This better stability is useful for experiments.

(3) Structurally, PW_{11} (or SiW_{11}) has a “pit” called “lacunary hole” in its molecule (Fig. 1), which is made by releasing one tungstate unit from the symmetrical $\text{PW}_{12}/\text{SiW}_{12}$ molecule upon partial hydrolysis. PW_{11} or SiW_{11} can hold another molecule (such as a cobalt ion (Co^{2+})) in the lacunary hole, like porphyrins can, forming a new compound called a “lacunary-substituted” undecatungstophosphate ($[\text{PW}_{11}\text{O}_{39}\text{X}]^{n-7}$) or undecatungstosilicate ($[\text{SiW}_{11}\text{O}_{39}\text{X}]^{n-8}$,

SiW_{11}X).^{5,6} Because the stability constant ($K=[\text{SiW}_{11}\text{X}]/([\text{SiW}_{11}]\cdot[\text{X}^{n+}])$) between SiW_{11} and metal cation X^{n+} is generally higher (*ca.* 10^7) than that of PW_{11} and X^{n+} (*ca.* 10^4),⁸ SiW_{11} will give more stable complexes than PW_{11} (*i.e.*, the effect of free metal cations “ X^{n+} ” released from SiW_{11}X can be ignored).

Of the related tungsten compounds tested, Keggin-structural polyoxotungstates and their mono- or tri-vacant lacunary species have the strongest sensitizing effects.^{9,10} Interestingly, these effective compounds are highly accumulated in the membrane fraction, and only a small portion is incorporated inside the bacterial cells.¹⁰ However, ionic substances are generally insoluble in such a hydrophobic region. Therefore, it was supposed that the lacunary hole might function as a binding site that interacts with some components of the bacterial membrane.

If the lacunary hole is essential to produce the sensitizing effect, the effect of lacunary-substituted species would be reduced, since the binding site is already occupied. In this report, the effects of several lacunary-substituted compounds on MRSA strains were studied to determine if the lacunary hole is required for the sensitizing effect.

MATERIALS AND METHODS

Chloride salts of tetraphenylphosphonium (TPP) and bis[2-(5-chloro-2-pyridylazo)-5-diethylaminophenolato]cobalt(III) ((CPADAP)₂Co) are products of Dojin Chemicals (Kumamoto, Japan). The other chemicals, culture media, established bacterial strains used in this study, and the method of preparing PW_{11} are described in our previous reports.^{1,2,11} Medium pH was kept at $\text{pH } 7.0 \pm 0.1$ to avoid decomposition of the polyoxotungstates. Although all the tungsten compounds used in the experiments are heat-stable, WO_4^{2-} and/or SiO_3^{2-} were mixed with the agar medium below about 45 °C. Otherwise, they can polymerize and form a certain effective compound.

Antimicrobial susceptibility was measured by the microdi-

* To whom correspondence should be addressed. e-mail: tajima@post.saga-med.ac.jp

lution broth method or the paper disk method approved by the National Committee for Clinical Laboratory Standards. The effect of various compounds was expressed as an 'Enhancement Index' = $\log_2(\text{MIC without agent}/(\text{MIC with agent}))$.¹⁾ To evaluate the synergism, the fractional inhibitory concentration (FIC) index was measured by a checkerboard study based on the microdilution broth method.

To prepare SiW_{11} , a modification of Komura's method was used according to the following equation: $[\text{SiW}_{12}\text{O}_{40}]^{4-} + 6\text{OH}^- \rightarrow [\text{SiW}_{11}\text{O}_{39}]^{8-} + \text{WO}_4^{2-} + 3\text{H}_2\text{O}$.¹²⁾ Briefly describing, SiW_{12} (33 g as $\text{H}_4[\text{SiW}_{12}\text{O}_{40}] \cdot 24\text{H}_2\text{O}$) was dissolved in 1 M CH_3COOH (200 ml), and this mixture (heated to 95 °C with vigorous stirring) was neutralized to pH 6.0 by gradual addition of solid KHCO_3 (ca. 25 g).¹³⁾ This neutralized solution was filtered once, and the filtrate was kept at 4 °C for a few days to allow a white powder to sediment. The precipitate was recovered by filtering, and then dissolved in hot water and recrystallized twice. Finally, the SiW_{11} thus obtained was washed with cold water (4 °C) and air-dried at room temperature. Purified SiW_{11} was dissolved in water to 20 mM (pH became ca. 6) and kept at -20 °C until use. If precipitation occurred during storage, the solution was warmed to 60 °C.

To prepare SiW_{11}X , SiW_{12} solution was neutralized to make SiW_{11} in a similar way to that above. Then, this acetate-buffered solution (pH 6.0) was filtered once, and the filtrate was kept at 95 °C. Next, 0.01 mol of a suitable compound of desired element " X^{n+} " (e.g., $\text{Co}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$) was dissolved in water (10 ml), and this was gradually added with vigorous stirring to the solution of SiW_{11} . Because lanthanide elements give a "2:1" complex of the form $\{[\text{SiW}_{11}\text{O}_{39}]\text{-X-}[\text{SiW}_{11}\text{O}_{39}]\}^{n-16}$ (Fig. 1),¹⁴⁾ the amount of the lanthanide solution mixed was 5 ml.

If the pH of the solution was significantly altered from 6.0, 2 M CH_3COOK solution was added dropwise to adjust the pH back to 6.0. After this mixture had been filtered and cooled, the crystal/powder was further purified by recrystallization as described above. An additional amount of the compound precipitated on adding an equal volume of methanol to the solution, and this increased the yield.

Because the water of crystallization content may vary depending on the drying procedure, elemental analysis was required for every preparation. To check the purity and concentration of $\text{SiW}_{11}/\text{SiW}_{11}\text{X}$, an inductively coupled plasma spectrometer and infrared spectrophotometer were used. Calculated/found weight contents (%) for SiW_{11} ($\text{K}_8[\text{SiW}_{11}\text{O}_{39}] \cdot 13\text{H}_2\text{O}$) were: K, 9.71/8.7; Si, 0.872/0.9; W, 62.78/63.5. Those for SiW_{11}Co ($\text{K}_6[\text{SiW}_{11}\text{O}_{39}\text{Co}(\text{H}_2\text{O})] \cdot 14\text{H}_2\text{O}$) were: K, 7.25/6.6; Si, 0.867/1.0; Co, 1.82/1.7; W, 62.45/63.6.

To prepare a solution of SiW_{11}Mg , 10 ml of 20 mM SiW_{11} solution was mixed with 200 μl of 1 M MgCl_2 immediately before use, and this mixture was used as a solution of 20 mM SiW_{11}Mg without further purification. This is because the stability constant ($K = [\text{SiW}_{11}\text{Mg}]/([\text{SiW}_{11}] \cdot [\text{Mg}^{2+}])$) between SiW_{11} and Mg^{2+} is so low (ca. 10^3) that purification by recrystallization will make no sense in this case.⁸⁾

RESULTS AND DISCUSSION

Effects of the Lacunary-Substituted SiW_{11} on MRSA Strains Table 1 shows the effects of the lacunary-substituted derivatives of SiW_{11} (SiW_{11}X) on several typical

MRSA strains. Most of the lacunary-substituted species had a (generally 2—4 times) stronger sensitizing effect than SiW_{11} . SiW_{11}Mg , SiW_{11}Al , and SiW_{11}Ca were almost equivalent to SiW_{11} , probably because of their low stability constant ($K = 10^2\text{—}10^3$): i.e., they easily dissociate to SiW_{11} and X^{n+} at a low concentration.

As shown in Table 2, all of the lacunary-substituted species had very strong synergism with oxacillin (MIPIC) on an MRSA strain. However, there was no relationship between the sensitizing potency of SiW_{11}X and the location of metal "X" in the periodic table of the elements. In terms of dose dependency, undecatungsto(ferri)ferrosilicate ($[\text{SiW}_{11}\text{O}_{39}\text{Fe}]^{6.5-}$, $\text{SiW}_{11}\text{Fe}(\text{II/III})$) was the most potent compound tested, followed by SiW_{11}Bi , SiW_{11}VO , and undecatungstocobaltosilicate ($[\text{SiW}_{11}\text{O}_{39}\text{Co}]^{6-}$, SiW_{11}Co).

The sensitizing effect of $\text{SiW}_{11}/\text{SiW}_{11}\text{X}$ was not due to the effects of their constituents. For example, almost no effect occurred when WO_4^{2-} , SiO_3^{2-} and metal cations such as Co^{2+} were mixed with the medium alone or in any combination (Fig. 2a). This means that the steric structure as a polyoxotungstate (Fig. 1) is essential to produce the sensitizing effect to MRSA strains, and the further enhancement of the sensitizing effect produced by lacunary-substitution can not be explained by simple toxicity of the substituents as a heavy metal.

Although its sensitizing effect was less than that of SiW_{11}Fe , the author considers that SiW_{11}Co is better for experimental use for the following reasons:

(1) When compared by selective toxicity (=MIC/effective concentration minimizing FIC index), SiW_{11}Co (2000 $\mu\text{M}/50 \mu\text{M}=40$) was superior to SiW_{11}Fe (200 $\mu\text{M}/10 \mu\text{M}=20$).

(2) In terms of FIC index, SiW_{11}Co gave the best result (Table 2). This was mainly due to milder toxicity of SiW_{11}Co , as described above, and MICs of MIPIC were minimized by any lacunary-substituted species; i.e., the higher the concentration of SiW_{11}X was, the smaller the MICs of MIPIC became. The compound with the lowest direct toxicity will be more advantageous than the others.

(3) In aqueous solution at around neutral pH, SiW_{11}Fe will partially form an oxo-bridged dimer as follows: $2[\text{SiW}_{11}\text{O}_{39}\text{Fe}]^{n-} + \text{O}^{2-} \leftrightarrow \{[\text{SiW}_{11}\text{O}_{39}\text{Fe}]\text{-O-}[\text{SiW}_{11}\text{O}_{39}\text{Fe}]\}^{2n-2}$.¹⁵⁾ Thus, it is unclear which molecular species is more effective. SiW_{11}Co is a single compound and can be easily purified, which is also an advantage.

The Role of the Lacunary Hole in the Sensitizing Effect

The sensitizing effects of the lacunary-substituted species of SiW_{11} (SiW_{11}X) were generally stronger than that of the original compound (SiW_{11}), indicating that occupation of the hole enhanced the sensitizing effect. To confirm this, the lacunary hole was blocked with more bulky molecules. Some ligands (e.g., CN^-) can coordinate with the metal ions in the lacunary hole of SiW_{11} , and one water molecule (H_2O) is usually bound there loosely (e.g., $[\text{SiW}_{11}\text{O}_{39}\text{Co}(\text{H}_2\text{O})]^{6-}$, see Fig. 1).^{5,6,12)} However, almost no change occurred in the advanced enhancing effect of SiW_{11}Co from SiW_{11} when CN^- , SCN^- , NO_2^- or pyridine was mixed with SiW_{11}Co to make a further coordinated complex, such as $[\text{SiW}_{11}\text{O}_{39}\text{Co}(\text{CN})]^{7-}$ (data not shown).

Initially, the author had supposed that $\text{PW}_{11}/\text{SiW}_{11}$ bound to some bacterial components with the lacunary hole, and the

Table 1. The Sensitizing Effect of Several Polyoxotungstates on MRSA Strains

Compounds ^{a)}	Color	log ₁₀ K ^{b)}	Concentration (μM)			
			10	25	50	100
			(MRSA Strains: ATCC43300/MRS394-1/SR3605) ^{c)} Enhancement Index of MPIPC ^{d)}			
PW ₁₁	—	—	+2/+2/+2	+4/+4/+4	+6/ +6/ +6	+7/ +8/ +7
SiW ₁₁	—	—	+3/+3/+2	+5/+4/+4	+7/ +6/ +6	+8/ +8/ +8
SiW ₁₁ Mg(II) ^{e)}	—	3.2	+3/+3/+2	+5/+4/+4	+7/ +6/ +6	+8/ +8/ +8
SiW ₁₁ Al(III)	—	—	+3/+3/+2	+5/+4/+4	+7/ +6/ +6	+8/ +8/ +8
SiW ₁₁ Ca(II)	—	4.0	+3/+3/+2	+5/+4/+4	+7/ +6/ +6	+8/ +8/ +8
SiW ₁₁ V(IV)O	Dark Violet	—	+4/+4/+4	+6/+6/+6	+8/ +8/ +8	+9/ +9/ +9
SiW ₁₁ Cr(III)	Green-Yellow	—	+3/+3/+2	+5/+5/+6	+8/ +8/ +8	+9/ +9/ +9
SiW ₁₁ Mn(II)	Yellow	6.2	+3/+3/+2	+6/+6/+6	+8/ +8/ +8	+9/ +9/ +9
SiW ₁₁ Fe(II) ^{f)}	Dark Brown	7.2	+6/+6/+6	+8/+8/+7	+10/+11/+9	>+10/>+11/+11 ^{g)}
SiW ₁₁ Fe(III) ^{f)}	Light Yellow	7.2	+6/+6/+6	+8/+8/+7	+10/+11/+9	>+10/>+11/+11 ^{g)}
SiW ₁₁ Co(II)	Wine Red	7.2	+4/+3/+3	+5/+5/+5	+8/ +7/+7	+9/ +9/ +8
SiW ₁₁ Ni(II)	Yellow	7.2	+3/+3/+3	+5/+5/+5	+7/ +7/+7	+8/ +9/ +8
SiW ₁₁ Cu(II)	Light Blue	8.0	+3/+3/+2	+5/+5/+4	+7/ +6/+6	+8/ +8/ +8
SiW ₁₁ Zn(II)	—	7.2	+3/+3/+2	+5/+5/+4	+7/ +6/+6	+8/ +8/ +8
SiW ₁₁ La(III) ^{h)}	—	—	+3/+4/+3	+5/+5/+5	+7/ +8/+7	+9/ +9/ +8
SiW ₁₁ Tm(III) ^{h)}	—	—	+3/+3/+3	+5/+4/+4	+6/ +7/+6	+8/ +9/ +8
SiW ₁₁ Yb(III) ^{h)}	—	—	+3/+3/+3	+5/+4/+5	+7/ +7/+7	+8/ +8/ +8
SiW ₁₁ Bi(III) ^{h)}	—	—	+5/+5/+4	+6/+7/+5	+8/ +8/+7	+10/ +9/ +9

a) The atomic valency of each substituents is shown as a roman number in parentheses after the element. b) Estimated from the graph shown in Ref. 8. This seems to be a function of atomic radius of the substituents. The stability constants (K) of elements not listed here were assumed to be 10⁶–10⁷. c) Penicillinase activity of ATCC43300/MRS394-1/SR3605 was +/–/–, and their resistance to MPIPC was (hetero and inducible)/(homo and constitutive)/(homo and constitutive), respectively. d) For example, ‘+7’ means that a compound made this MRSA strain (2⁷)128-fold susceptible to the drug. e) This is an unpurified mixture of SiW₁₁ and MgCl₂ (see text). Other compounds were purified by recrystallization. f) An oxo-bridged dimer of the form {[SiW₁₁O₃₉Fe]-O-[SiW₁₁O₃₉Fe]}¹²⁻ was partially formed. g) ‘>’ means that MICs in the presence of SiW₁₁Fe were below the lower measurement range (≅2⁻⁵ mg/l). h) A ‘2:1’ complex such as {[SiW₁₁O₃₉]-X-[SiW₁₁O₃₉]}¹²⁻ is formed.

Table 2. FIC Index of Several Polyoxotungstates and MPIPC on an MRSA Strain

Compounds ^{a)}	MICs of MPIPC (mg/l)	MICs of Polyoxotungstates (μM)	FIC Index ^{b)}
	(Alone/Combined)	(Alone/Combined)	
PW ₁₁	64/1	2000/50	0.041
SiW ₁₁	64/1	2000/50	0.041
SiW ₁₁ V(IV)O	64/1	500/25	0.066
SiW ₁₁ Cr(III)	64/0.25	1000/50	0.054
SiW ₁₁ Mn(II)	64/1	1000/25	0.041
SiW ₁₁ Fe(II/III)	64/1	200/10	0.066
SiW ₁₁ Co(II)	64/0.5	2000/50	0.033
SiW ₁₁ Ni(II)	64/0.5	1500/50	0.041
SiW ₁₁ Cu(II)	64/2	1500/25	0.048
SiW ₁₁ La(III)	64/2	500/25	0.081
SiW ₁₁ Yb(III)	64/0.5	1000/50	0.058
SiW ₁₁ Bi(III)	64/2	500/10	0.051

a) Details are shown in the legend of Table 1. b) Results shown here are the data which minimized the FIC index. This is a result obtained from a typical MRSA strain (MRS394-1), and SiW₁₁Co also generally gave the minimum FIC index in other MRSA strains (data not shown).

effect of lacunary-substituted species (SiW₁₁X) would be reduced because the binding site was already occupied. However, the result was opposite, which suggests that the hole is not essential to produce the sensitizing effect on MRSA strains, although polyoxotungstates with a lacunary hole are generally more effective than other molecular species.^{9,10)}

When small and lipophilic cations, such as TPP and (CPADAP)₂Co, were mixed with the culture medium containing SiW₁₁, the sensitizing effect of SiW₁₁ was further enhanced, whereas a bulky polycation (such as protamine) inhibited it (Fig. 2b). When SiW₁₁Co was used instead of

Fig. 1. Structure of “Keggin”-Structural (Hetero)polyoxotungstates

The left panel shows the structure of the SiW₁₂ molecule ([SiW₁₂O₄₀]⁴⁻). One silicate (SiO₄²⁻) unit, which is located in the center of the molecule (shaded), is surrounded by twelve tungstate (WO₄²⁻) units. This is a typical example of a “Keggin”-structural (hetero)polyoxotungstate. The right panel shows the structure of the SiW₁₁ molecule ([SiW₁₁O₃₉]⁸⁻). One tungstate unit is released from the SiW₁₂ molecule by partial hydrolysis, and this site is called a “lacunary hole”. This type of compound is sometimes known as a “mono-vacant” lacunary-Keggin. Some metal cations (such as Co²⁺) can be trapped by the lacunary hole and to form a new compound called a “lacunary-substituted” (hetero)polyoxotungstate. Lanthanide elements give a “2:1” complex with a sandwich-like structure.

SiW₁₁, TPP/(CPADAP)₂Co still enhanced the sensitizing effect, but hydrophilic cations such as ethanolamine showed no enhancement effect (data not shown).

It has been reported that such a lipophilic cation functions as a phase-transfer agent to many polyanions^{16,17)}; i.e., SiW₁₁ makes an ion-pair with TPP/(CPADAP)₂Co and the complex becomes extractable to organic solvents. Some investigators

Fig. 2. The Effects of Several Compounds on MRSA Strains

General outline of the effects of several compounds on MRSA strains (MRS394-1/ATCC43300) is shown. Various reagents such as SiW₁₁ were mixed directly with the agar medium at the final concentrations shown here, and photographs were taken at 24 h after the inoculation. Similar patterns were obtained from all the MRSA strains, and typical examples are shown here.

have applied this feature to potentiometric measurement.^{18,19} Therefore, the author considers that a charge interaction between these polyoxotungstates and some unknown (perhaps positively charged) materials in the culture medium or bacterial cells plays an important role on the sensitizing effect (e.g., amino acids including oligopeptides and aliphatic amines such as phosphatidylethanolamine are candidates for making ion-pairs with SiW₁₁). It is possible that metal cations in the lacunary hole may only neutralize the high negative charge of SiW₁₁, like TPP/(CPADAP)₂Co did, thereby further enhancing the sensitizing effect.

Suggested Mechanism of Action of Polyoxotungstates

As shown in Table 3 and our previous reports,^{1,2} the sensitizing effect of polyoxotungstates on MRSA strains occurred in all the β -lactam antibiotics (including cepheims and carbapenems but not monobactams). Furthermore, there was no synergism of the polyoxotungstates with other classes of antibiotics including vancomycin and teicoplanin (both inhibitors of cell wall synthesis, like β -lactams²⁰), and there was no effect against bacterial strains other than staphylococcal species.^{1,2}

Although the mechanism of action of polyoxotungstates has not been completely elucidated, they seem to reduce the expression of penicillin-binding proteins (PBPs), thereby making MRSA strains greatly susceptible to β -lactams. These compounds neither directly inactivated PBPs nor increased in their affinities for β -lactams.^{2,3,9,10}

The resistance of MRSA strains to β -lactams is principally

attributed to acquisition of an altered PBP (PBP2') with lowered affinity to β -lactams.^{20,21} Although the suppressive effect was stronger against PBP2', polyoxotungstates also reduced the expression of ordinary PBPs (PBP1—4).^{2,10} Therefore, polyoxotungstates had a weak sensitizing effect on methicillin-susceptible *S. aureus* (MSSA) strain (Table 3), but this might not be obvious in the "Enhancement Index" because β -lactams acted strongly on such strains even before sensitization by the polyoxotungstates.^{1,2} Probably for the same reason, MICs of β -lactams were minimized even in MRSA strains and sometimes reduced below the lower measurement limits (Tables 1 and 3).

It seems likely that polyoxotungstates inhibited the activity of enzymes with nucleic acids as their substrates. Nucleic acids can be regarded as a type of polyanion, and the polyoxotungstates probably inhibited the enzymes by acting as a nucleic acid-analogue in a charge interaction.^{3,22–24} Other enzymes (such as transaminases and proteases) were generally insensitive to these tungsten compounds.³ The DNA-binding activity of anti-DNA antibody was recently found to be inhibited by low concentration of PW₁₁ (ca. 1 mM). This also suggests that polyoxotungstates act as nucleic acid-analogues.²⁵

Because PBP2' is a very fragile protein and probably has rapid turnover,²⁶ cells of MRSA strains seem to be required to constantly produce enough PBP2' for full expression of resistance to β -lactams. Thus, the author now considers that this resistance may be more easily affected than other pheno-

Table 3. Effect of $\text{PW}_{11}/\text{SiW}_{11}/\text{SiW}_{11}\text{Co}$ on Various Types of Antibiotics

Antimicrobial Agents	Strains								
	MSSA NCTC 8325			MRSA (inducible) ATCC 43300			MRSA (constitutive) MRS 394-1		
	Enhancement Index ^{a)} (100 μM PW ₁₁ /SiW ₁₁ /SiW ₁₁ Co)								
Oxacillin	+1/	+1/	+2	+7/	+8/	+9	+8/	+8/	+9
Cefazolin	+1/	+1/	+1	+7/	+7/	+8	+5/	+6/	+7
Cefmetazole	+1/	+1/	+1	+6/	+6/	+7	+5/	+6/	+7
Flomoxef	+1/	+1/	+1	+5/	+5/	+6	+3/	+4/	+5
Cefozopran	0/	0/	0	+4/	+4/	+6	+4/	+5/	+6
Imipenem	0 ^{b)/}	0 ^{b)/}	0 ^{b)}	>+6 ^{c)/} >+6 ^{c)/} >+6 ^{c)}			+5/	+5/	+6
Fosfomycin	0/	0/	0	0/	0/	0	0/	0/	0
Erythromycin	0/	0/	0	0/	0/	0	0/	0/	0
Clindamycin	0/	0/	0	0/	0/	0	0/	0/	0
Gentamicin ^{d)}	0/	0/	0	-1/	-1/	-1/	-1/	-1/	-1
Minocycline ^{d)}	-2/	-2/	-2	-2/	-2/	-2/	-2/	-2/	-2
Tosufloxacin	0 ^{b)/}	0 ^{b)/}	0 ^{b)}	0 ^{b)/}	0 ^{b)/}	0 ^{b)}	0 ^{b)/}	0 ^{b)/}	0 ^{b)}
Vancomycin	0/	0/	0	0/	0/	0	0/	0/	0
Teicoplanin	0/	0/	0	0/	0/	0	0/	0/	0

a) Bold lettering (such as '+3') indicates resistance; i.e., MIC without tungsten compounds was $\geq 2^4$ mg/l. The MICs were measured up to 2^{10} mg/l. b) Both MICs with and without tungsten compounds are $\leq 2^{-5}$ mg/l (below the lower measurement range). c) MIC with tungsten compounds is $\leq 2^{-5}$ mg/l. d) These tungsten compounds made insoluble complexes with these antibiotics, thereby reducing their antimicrobial activities (see text).

typic characteristics, even though several (anion-sensitive) ordinary cell functions such as transcription and translation activity are only weakly inhibited by polyoxotungstates.

The Effects of Polyoxotungstates on Aminoglycoside and Tetracycline Polyoxotungstates rather reduced the efficacy of aminoglycoside and tetracycline (Table 3). This is probably because these tungsten compounds are a type of polyanion, and they may neutralize such a cationic substance. For instance, when 100 μM SiW_{11} solution was mixed with 20–50 mg/l gentamicin or 100–200 mg/l minocycline, the mixture became turbid. Furthermore, metachromatic reaction of toluidine blue induced by SiW_{11} was also inhibited by these antibiotics.

However, these polyoxotungstates did not precipitate other classes of antibiotics at a working concentration: For instance, >5 mM SiW_{11} caused almost no precipitation with >5000 mg/l vancomycin (or teicoplanin), and β -lactams did not make insoluble complex with these tungsten compounds at any concentration.

Suggested Use of the Polyoxotungstates Unfortunately, polyoxotungstates such as SiW_{11} can not be used clinically, because they are toxic to animals. When directly injected to the vein of an animal, pulmonary embolism occurred, probably due to platelet aggregation induced by the compounds (details to be published elsewhere).

The author now intends to use these tungsten compounds as a "tool" in laboratory tests: e.g., screening media for resistant bacterial strains. Interestingly, MRSA strains with a high resistance to β -lactams were easily detected by adding these compounds to the medium. When they were mixed with the culture medium, the growth inhibitory zone formed around a paper disk containing a β -lactam antibiotic such as MIPIC was significantly enlarged. However, colonies appeared inside the circle if the inoculated MRSA strain was hetero-resistant to the agent and contained subclones with a higher resistance.

For example, the MIC of MIPIC in a typical MRSA strain

ATCC43300 was around 32 mg/l, and this was reduced to 0.25–0.5 mg/l in the presence of 50 μM SiW_{11} . However, we can just see some colonies inside the growth inhibitory zone (indicated by an arrow in Fig. 2b), and their MICs of MIPIC alone were increased to 64–128 mg/l. When such selection was repeated, the MIC of MIPIC became ca. 256 mg/l. It is noteworthy that the geometric pattern of the colonies was roughly related to the result of the population analysis (details to be published elsewhere).

CONCLUSION

Since the lacunary-substituted species of SiW_{11} generally had a stronger effect than SiW_{11} , the lacunary hole does not seem to be required for the sensitizing effect on MRSA strains. Of the related compounds tested, SiW_{11}Co was found to give the best FIC index. As a small and lipophilic cations further enhanced the sensitizing effect of SiW_{11} , it was suggested that SiW_{11} made an ion-pair to interact with the bacterial cells.

REFERENCES

- 1) Tajima Y., Nagasawa Z., Tadano J., *Microbiol. Immunol.*, **37**, 695–703 (1993).
- 2) Tajima Y., Nagasawa Z., Tanabe I., Kusaba K., Tadano J., *Microbiol. Immunol.*, **38**, 639–648 (1994).
- 3) Tajima Y., Nagasawa Z., Tanabe I., Kusaba K., Tadano J., *Res. Microbiol.*, **147**, 279–286 (1996).
- 4) Tajima Y., *J. Inorg. Biochem.*, **68**, 93–99 (1997).
- 5) Pope M. T., "Comprehensive Coordination Chemistry," Vol. 3, ed. by Wilkinson G., Gillard R. D., McCleverty J. A., Pergamon Press, Oxford, 1987, pp. 1023–1058.
- 6) Pope M. T., Müller A., *Angew. Chem. Int. Ed. Engl.*, **30**, 34–48 (1991).
- 7) Kepert D. L., "Comprehensive Inorganic Chemistry," Vol. 4, ed. by Bailar J. C., Jr., Emeléus H. J., Nyholm R., Trotman-Dickenson A. F., Pergamon Press, Oxford, 1973, pp. 607–672.
- 8) Constant R., Ciabrini J. P., *J. Chem. Res. (S)*, 50–51 (1982).
- 9) Yamase T., Fukuda N., Tajima Y., *Biol. Pharm. Bull.*, **19**, 459–465

- (1996).
- 10) Fukuda N., Yamase T., Tajima Y., *Biol. Pharm. Bull.*, **22**, 463—470 (1999).
- 11) Tajima Y., *J. Biochem. Biophys. Methods*, **38**, 217—223 (1999).
- 12) Komura A., Hayashi M., Imanaga H., *Bull. Chem. Soc. Jpn.*, **49**, 87—91 (1976).
- 13) Weakley T. J. R., Malik S. A., *J. Inorg. Nucl. Chem.*, **29**, 2935—2944 (1967).
- 14) Peacock R. D., Weakley T. J. R., *J. Chem. Soc. (A)*, 1836—1839 (1971).
- 15) Zonnevijlle F., Tourné C. M., Tourné G. F., *Inorg. Chem.*, **21**, 2751—2757 (1982).
- 16) Ripan R., Calu C., *Talanta*, **14**, 887—891 (1967).
- 17) Taga M., Kan M., *Bull. Chem. Soc. Jpn.*, **62**, 1482—1485 (1989).
- 18) Bouklouze A. A., El-Jammal A., Patriarche G. J., Christian G. D., *J. Pharm. Biomed. Anal.*, **9**, 393—399 (1991).
- 19) Wen M. L., Zhao Y. B., Chen X., Wang C. Y., *J. Pharm. Biomed. Anal.*, **18**, 957—961 (1999).
- 20) Hiramatsu K., Ito T., Hanaki H., “Baillière’s Clinical Infectious Disease,” Vol. 5, ed. by Finch R. G., Williams R. J., Baillière Tindall, London, 1999, pp. 221—242.
- 21) Hiramatsu K., *Microbiol. Immunol.*, **39**, 531—543 (1995).
- 22) Jasmin C., Raybaud N., Chermann J. C., Haapala D., Sinoussi F., Loustau C. B., Bonissol C., Kona P., Raynaud M., *Biomedicine*, **18**, 319—327 (1973).
- 23) Haapala D. K., Jasmin C., Sinoussi F., Chermann J. C., Raynaud M., *Biomedicine*, **19**, 7—11 (1973).
- 24) Inouye Y., Tokutake Y., Yoshida T., Yamamoto A., Yamase T., Nakamura S., *Chem. Pharm. Bull.*, **39**, 1638—1640 (1991).
- 25) Tajima Y., *Clin. Lab.*, **47**, 289—292 (2001).
- 26) Utsui Y., Yokota T., *Antimicrob. Agents Chemother.*, **28**, 397—403 (1985).