Biol. Pharm. Bull.

Regular Article

**Indomethacin, a non-steroidal anti-inflammatory drug, induces skin
dryness via PPAR γ in mice**

Kiyoko Maruyama,^a Kenji Goto,^b Keiichi Hiramoto,^a Shota Tanaka^a and Kazuya Ooi^{a,*}

^a*Department of Pharmaceutical Sciences, Suzuka University of Medical Science, 3500-3*

Minamitamagaki, Suzuka, Mie 513-8607, Japan

^b*Research Laboratories, Nichinichi Pharmaceutical Co., Ltd., 239-1 Tominaga, Iga,*

Mie 518-1417, Japan.

*To whom correspondence should be addressed. e-mail: zooi@suzuka-u.ac.jp

Department of Pharmaceutical Sciences, Suzuka University of Medical Science,

3500-3 Minamitamagaki, Suzuka, Mie 513-8607, Japan

SUMMARY

Cyclooxygenase (COX)-1-selective inhibitors have side effects such as itching and dryness of the skin. In this study, the degree of skin dryness and the onset mechanism of this condition were investigated by comparing the effects of three non-steroidal anti-inflammatory drugs (NSAIDs) in mice. Mice were orally administered either indomethacin, loxoprofen sodium, or celecoxib (n = 5 per group) once daily for four consecutive days, and blood samples as well as skin and jejunal tissues were isolated on day 5. In the mice treated with indomethacin, transepidermal water loss was significantly increased, and dry skin was observed. In addition, the expression of matrix metalloproteinase (MMP)-I, mast cells, CD163, CD23, CD21, histamine, and peroxisome proliferation-activated receptor (PPAR) γ in the skin and jejunum was increased, and the blood levels of interleukin-10 and immunoglobulin E were also increased. In contrast, the expression of collagen type I in the skin was decreased. These results show that indomethacin activates PPAR γ in the skin and jejunum, changes the polarity of macrophages, increases the secretion of MMP-1 from mast cells, and decomposes collagen type I, leading to dry skin.

Key words: indomethacin; dry skin; macrophages; mast cell; peroxisome proliferation-activated receptor γ

INTRODUCTION

The number of elderly people in Japan is increasing, and the prevalence of chronic pain is also increasing. Non-steroidal anti-inflammatory drugs (NSAIDs) are widely used to treat pain and many inflammatory diseases.¹⁾ NSAIDs decrease the production of prostaglandins by inhibiting cyclooxygenase (COX). COX has two main isozymes, COX-1 and COX-2. COX-1 is constitutively expressed in most tissues. Specifically, it has a protective effect on the mucosa of the stomach and intestine.²⁾ COX-2 is primarily localized to inflammatory cells and tissues. It induces inflammation, fever, and pain.³⁾ Many NSAIDs inhibit COX-2 to suppress inflammation and the progression of inflammatory reactions. However, it has been reported that some NSAIDs, which are not selective toward COX, cause adverse effects in digestive organs upon long-term use.⁴⁾

Gut, including small intestine, influences skin conditions through various mechanisms. The intestine often causes systemic inflammation by release of inflammatory materials. This ability has important implications in skin disease. In addition, this ability is affected by intestinal conditions, such as increase of intestinal permeability.

We previously reported that dry skin may be associated with indomethacin-induced small intestine injury in mice.⁵⁾ In that study, we found increased expression of mast

cells in the small intestine. Mast cells are known to release tryptase and histamine.⁶⁾ Therefore, it has been suggested that indomethacin-induced small intestinal mast cells may be involved in skin dryness. However, the detailed induction mechanism of skin dryness by indomethacin is unknown.

Furthermore, indomethacin acts as an agonist of peroxisome proliferation-activated receptor (PPAR) γ , a subtype of PPAR.⁷⁾ PPAR γ is a nuclear receptor and a nuclear transposable element belonging to the steroid hormone receptor superfamily. It is also involved in the differentiation and proliferation of adipose tissue cells, inflammation, and immune response.^{8,9)} However, the effect of PPAR γ on mast cell degranulation, which plays an important role in inflammation and dry skin, is unknown.

Therefore, in this study, we administered indomethacin, loxoprofen sodium, and celecoxib to mice to examine whether the degree of intestinal damage is correlated with the extent of skin disruption. In addition, the effects of indomethacin and mast cells on skin dryness were investigated.

MATERIALS AND METHODS

Chemicals. Indomethacin was purchased from Sigma-Aldrich (St. Louis, MO, USA). Loxoprofen sodium was purchased from Wako Pure Chemical Industries, Ltd. (Osaka, Japan). Celecoxib was purchased from TOCRIS Bioscience (Avonmouth, Bristol, UK). All other reagents were of analytical grade. Indomethacin selectively inhibits COX-1, whereas loxoprofen sodium inhibits both COX-1 and COX-2 to the same degree. Celecoxib selectively inhibits COX-2 expression.

Animals. Hairless mice (HOS: HR-1) were purchased from SLC (Hamamatsu,

Shizuoka, Japan). The mice were housed under a 12-h light cycle at a constant temperature of 23 ± 2 °C and relative humidity of $55 \pm 10\%$. The mice were fed laboratory chow (CE-2; Oriental Yeast Co., Tokyo, Japan) and water *ad libitum*.

Experimental design. At eight weeks of age, hairless mice were randomly assigned (n = 5 per group) to oral administration of indomethacin (40 mg/kg, suspended in saline), loxoprofen sodium (30 mg/kg, suspended in distilled water), or celecoxib (10 mg/kg, dissolved in 0.25% dimethyl sulfoxide) once a day for four days. The doses of indomethacin, loxoprofen and celecoxib (40, 30, and 10 mg/kg) used in the preliminary examination were based on the method of Kuzumoto et al. (2021),¹⁰⁾ Shimada et al. (2020)¹¹⁾, and Liang et al. (2021).¹²⁾ Each value is the maximum concentration that does not affect the health changes (weight loss, behavioral abnormalities) of mice after long-term administration. Control mice were administered the corresponding delivery vehicles according to the same schedule. This study strictly followed the recommendations and guidelines for the care and use of laboratory animals at the Suzuka University of Medical Science (approval number: 34). All surgical procedures were performed under anesthesia induced by pentobarbital, and every effort was made to minimize animal suffering.

PPAR γ antagonist treatment. In the PPAR γ antagonist-treated group, mice were

intraperitoneally injected with a PPAR γ antagonist (GW9662: 1 mg/mL; Wako Pure Chemical Industries) in saline (0.1 mL) daily for four days. Animals in the control group were administered saline only.¹³⁾

Measurement of transepidermal water loss (TEWL) and capacitance. On day 5, the TEWL and capacitance in the dorsal skin of each mouse were measured. TEWL and capacitance were used as a marker of skin permeability, reflecting the barrier function of the skin; decreased TEWL and increased capacitance indicates dry skin¹⁴⁻¹⁶⁾. TEWL measurements were carried out using a Tewameter TM300 (Courage + Khazaka Electronic GmbH, Cologne, Germany).¹⁷⁾ Values were recorded once the reading had stabilized, typically 10 s after the probe was placed on the skin. The data are presented as the average of three independent measurements. The level of hydration in the outermost layer of the skin, the stratum corneum, was measured on day 5 using a Corneometer CM825 (Courage + Khazaka Electronic GmbH).¹⁸⁾ The Corneometer probe was applied to the dorsal skin surface of each mouse, and the degree of skin hydration was determined by the electrical capacitance of the surface of the skin, expressed in arbitrary units. The reported data are the average of three independent measurements per test area.

Measurement of histamine, interleukin (IL)-10, and immunoglobulin (Ig)E levels.

Skin, jejunum, and plasma samples were collected on the last experimental day. We rinsed 100 mg of skin and jejunum tissues in phosphate-buffered saline (PBS) to remove excess blood before homogenization. The tissues were finely minced and homogenized in 1 mL of PBS with a glass homogenizer on ice. The cells were lysed by ultrasonication at room temperature two to three times. The samples were centrifuged at $1,500 \times g$ for 15 min. The supernatant was collected for assay. The levels of histamine in the skin and jejunum, and the plasma levels of IL-10 and IgE were determined using commercial ELISA kits (histamine: Bertin Pharmacol, Montigny-le-Bretonnux, France; IL-10: Proteintech, Rosemont, IL, USA; IgE, Yamasa Shoyu Co., Ltd., Chiba, Japan) according to the manufacturer's instructions. Optical density was measured using a microplate reader (Molecular Devices, Sunnyvale, CA, USA).

Tissue section staining. Jejunal and dorsal skin tissue samples were isolated and fixed in PBS containing 4% paraformaldehyde (Wako Pure Chemical Industries). Fixed tissue specimens were embedded in a frozen Tissue-Tek OCT Compound (Sakura Finetek, Tokyo, Japan) and sliced into 5- μ m-thick sections. The skin and jejunal specimens were stained with hematoxylin-eosin (HE) in accordance with established procedures to enable histological analysis. We microscopically evaluated HE stained skin tissues. To determine overall skin thickness, 10 regions in which the skin appeared

flat in acquired images were randomly selected. The length from the outer layer of the epidermis to the border of the subcutis was measured, and the average value was calculated. The cells were stained with toluidine blue to evaluate the number of mast cells via fluorescence microscopy using ImageJ software. Mast cells were quantified by counting cell numbers per mm² field in 10 randomly selected regions. In addition, the skin specimens were stained using antibodies for immunological analysis according to a previously published method.⁵⁾ Briefly, the specimens were incubated with either mouse monoclonal anti-CD23 (B cell marker, 1:100; Proteintech) or rabbit polyclonal anti-CD21 (1:100; Abcam, Cambridge, UK) primary antibodies. The samples were then incubated with fluorescein isothiocyanate-conjugated anti-mouse and tetramethylrhodamine isothiocyanate-conjugated anti-rabbit (1:30; Dako Cytomation, Glostrup, Denmark) secondary antibodies. The expression of CD23 and CD21 was evaluated immunohistochemically using fluorescence microscopy.

Western blot analysis. The dorsal skin samples were homogenized in a lysis buffer (Kurabo, Osaka, Japan) and then centrifuged at 8,000 ×g for 10 min. The supernatant from each sample was collected and stored at -80 °C until analysis. After thawing, equal amounts of protein (5 µg/lane) were loaded onto a 4%–12% BIS-TRIS Bolt gel (Life Technologies, Carlsbad, CA, USA) and the gel was subjected to electrophoresis at

200 V for 30 min. Following separation, proteins were transferred onto a nitrocellulose membrane using an iBlot western blotting system (Life Technologies). The membranes were blocked with 5% skim milk at 4 °C overnight. After blocking, the membranes were incubated at 25 °C for 1 h with primary antibodies against type I collagen (1:1,000; EMD Chemicals Inc., Gibbstown, NJ, USA), matrix metalloproteinase (MMP)-1 (1:1,000; Santa Cruz Biotechnology Inc., Santa Cruz, CA, USA), mast cell tryptase (mast cell marker, 1:500; Santa Cruz Biotechnology Inc.), chemokine receptor 7 (M1 macrophage marker, 1:1,000; Santa Cruz Biotechnology Inc.), CD163 (M2 macrophage marker, 1:1000; Abcam), PPAR γ (1:1,000; Santa Cruz Biotechnology Inc.), p-PPAR γ (1:1,000; Bioss Inc, Woburn, MA, USA), CD23 (1:1000; Proteintech), and β -actin (1:5,000; Sigma-Aldrich). The membranes were incubated with a horseradish peroxidase-labeled secondary antibody (Life Technologies), and the signal was detected with ImmunoStar Zeta (Wako Pure Chemical Industries) using a lumino-image analyzer (LAS-4000; Fuji Film, Tokyo, Japan). Protein bands were subjected to densitometric analysis, and the values were normalized to those of β -actin.

Statistical analysis. All data are presented as the mean $\pm$ standard deviation. Data were analyzed using Microsoft Excel 2010 and one-way analysis of variance followed by Tukey's *post-hoc* test with SPSS software version 20 (IBM, Armonk, NY, USA).

Heatmap analysis and hierarchical clustering were performed using R software.

Differences were considered significant at $P < 0.05$.

RESULTS

The delivery vehicle of each NSAID was also administered to the control mice.

However, no differences were observed following treatment (data not shown).

Effect of NSAIDs on skin dryness in mice. First, we examined various indicators of dry skin. The TEWL and capacitance in the dorsal skin showed that the mice in the indomethacin and loxoprofen sodium treatment groups had a higher TEWL than that in the control mice. In particular, the TEWL increased significantly in the indomethacin-treated mice compared to that in the control mice (Fig. 1A). The capacitance was only reduced in the indomethacin-treated mice (Fig. 1B). The skin of indomethacin-treated mice was thicker than that of control mice; however, the skin of loxoprofen or celecoxib treated mice were not different from control mice (Fig. 1C). In addition, indomethacin treatment markedly reduced the expression of collagen type 1 in the skin (Fig. 1D) and markedly increased the expression of MMP-1 (Fig. 1E).

Effect of NSAIDs on the expression of mast cells in the skin and the histamine levels in the skin and jejunum. Toluidine blue staining of the skin and jejunum

sections indicated that the number of mast cells increased in each treatment group compared with that in the control group. In particular, the increase was the greatest in the indomethacin-treated group, followed by the loxoprofen sodium and celecoxib treatment groups (Figs. 2A, B). The results of measuring mast cells using Western blot showed the same tendency as the results of toluidine blue staining (Figs. 2C, D). In contrast, histamine levels were greatly increased by indomethacin treatment. The histamine levels with the other NSAID treatments were unchanged from those in the control group (Figs. 2C, D).

Effect of NSAIDs on the expression of PPAR γ and p-PPAR γ in the skin and jejunum. Next, we examined the expression of PPAR γ , for which indomethacin acts as an agonist. The PPAR γ expression in the skin was reduced by indomethacin treatment, and PPAR γ expression in the jejunum was the same as that in the control group after all NSAID treatments (Figs. 3C, D). Conversely, activated PPAR γ and p-PPAR γ showed high values in both the skin and jejunum (Figs. 3A, B).

Effect of NSAIDs on the expression of CCR7 and CD163 in the skin and jejunum. We investigated the localization of two types of macrophages. The expression of CCR7, an indicator of M1 macrophages, was unchanged in all groups (Figs. 4A, B). However, the expression of CD163, an indicator of M2 macrophages, was very high in

the skin and jejunum of the indomethacin-treated group (Figs. 4C, D).

Effect of NSAIDs on the plasma levels of IL-10 and IgE and the expression of CD23 in the skin and jejunum. We measured the plasma levels of IL-10 and IgE, as well as the expression of CD23 (mature B cells), in the skin and jejunum, which are closely associated with mast cells. The plasma levels of IL-10 and IgE were greatly increased by indomethacin treatment (Figs. 5A, B). However, in the other treatment groups, there was no difference from the levels in the control group. Similarly, the expression of CD23 in the skin and jejunum was also greatly increased by indomethacin treatment (Figs. 5C, D). In addition, we observed an increase in CD23 expression as well as co-localization of CD23 with CD21 in the mice treated with indomethacin (Fig. 5E).

Effect of PPAR γ antagonist treatment on the TEWL and the expression of PPAR γ and p-PPAR γ in the skin. We administered an antagonist of PPAR γ to confirm the role of indomethacin in PPAR γ . The TEWL, an indicator of skin dryness, increased due to the administration of indomethacin but was improved by administration of a PPAR γ antagonist (Fig. 6A). In contrast, expression of p-PPAR γ , which is the active form of PPAR γ , was greatly increased by treatment with indomethacin, but decreased by administration of an antagonist, and was the same as that of the control group (Figs. 6B,

C).

DISCUSSION

In the present study, we examined the interaction between dry skin and small intestinal injury induced by NSAIDs. We observed that treatment with NSAIDs (indomethacin, loxoprofen sodium, or celecoxib) increased the TEWL and decreased the capacitance in the skin of mice. However, these effects were most prominent in the indomethacin-treated group.

Generally, TEWL increases and capacitance decreases in dry skin.¹⁹⁾ Collagen, along with other proteins in the skin, act as natural moisturizers, and mast cell-induced inflammation decreases their levels to dry the skin.^{20, 21)} Mast cells cause skin dryness induced by lesions in the digestive tract.^{5, 22)} Histamine is a well-known bioactive amine that is released from mast cells when IgE binds to the Fcε receptor on mast cells.^{23, 24)} Additionally, tryptase released from mast cells activates MMPs. MMP-1 decomposes type I collagen, and MMP-9 decomposes type IV collagen, which then causes skin disruption.^{20, 21)} Therefore, we confirmed the number of mast cells in the skin of mice treated with NSAIDs (Fig. 2). The number of mast cells in the skin increased in each NSAID treatment group, and this increase was the greatest in the

239 indomethacin-treated mice.

240 The extent of skin dryness caused by NSAIDs depends on the COX selectivity of
241 each NSAID. COX-1-selective inhibitors have a greater potential to cause
242 gastrointestinal tract disturbances.²⁵⁾ Indomethacin selectively inhibits COX-1.
243 Accordingly, in our study, indomethacin induced gastrointestinal tract disturbance by
244 increasing the number of neutrophils to the greatest extent among all the tested NSAIDs.
245 As celecoxib is a COX-2-selective inhibitor, it is not expected to induce gastrointestinal
246 tract disturbance.²⁶⁾ Consistently, small intestinal injury was observed in the
247 celecoxib-treated mice. Moreover, skin dryness in the celecoxib-treated mice was mild
248 compared to that in the mice of the other groups.

249 The second reason why the indomethacin-treated mice experienced the driest skin
250 in this study is considered to be the relationship between indomethacin and PPAR γ . It is
251 generally known that indomethacin activates PPAR γ without suppressing the expression
252 of the transcription factors AP-1 and NF κ B.²⁷⁾ The role of PPAR γ in inflammation has
253 been advocated. An increase in bone marrow mononuclear cells has been observed
254 when examining cell proliferation of mouse bone marrow cells with the PPAR γ agonist
255 troglitazone, indicating that PPAR γ may be involved in an increase in mast cells.²⁸⁾
256 Indomethacin may have caused an increase in mast cells in the skin and jejunum in this

study via PPAR γ . Furthermore, activation of PPAR γ changes the polarity of M1/M2 macrophages and is important for inducing differentiation into M2 macrophages.^{29, 30)} M2 macrophages activate the secretion of IL-10, an anti-inflammatory cytokine that promotes B-cell maturation. CD23 was examined as a B-cell marker in this study. It is known that CD23 is mainly expressed on B cells, binds to an antigen-IgE immune complex, and is involved in the regulation of IgE production.³¹⁾ In particular, CD23 interacts with CD21 in B cells and preferentially promotes IgE production.³²⁾ Mast cells release various mediators (histamine, tumor necrosis factor- α , and matrix metalloproteinase) through IgE-mediated stimulation. These mediators have been suggested to cause dry skin. To prove this data, a PPAR γ antagonist was administered instead of indomethacin and drying of the skin was induced, suggesting that this pathway may be involved in skin dryness (Fig. 6).

In this study, administration of indomethacin increased histamine. Histamine H1 and H2 receptors expressed on keratinocytes are activated to delay the recovery of skin barrier function.³³ It has also been reported that TNF- α has many effects on the skin barrier function.^{34, 35)} However, this study does not examine histamine receptors or TNF- α . Therefore, further studies are needed on histamine receptors and TNF- α . In addition, topical application of PPAR γ normalized skin barrier homeostasis.³⁶⁾ Therefore,

275 it is necessary to apply indomethacin locally and study it.

276

277 CONCLUSION

278 From the above results, it is evident that small intestinal injury caused by
279 indomethacin induces skin dryness. In addition, it is suggested that indomethacin
280 changes the M2 type of macrophages by activating PPAR γ in the skin and jejunum, and
281 moves the downstream signal for IL-10/CD23 (B cell)/IgE/mast cells, leading to dry
282 skin. Additionally, we confirmed that COX-2-selective NSAIDs may be preferred over
283 other NSAIDs for their anti-inflammatory and analgesic effects, as they do not cause
284 significant skin dryness. This information may be critical when selecting NSAIDs for
285 patients predisposed to skin disorders.

Acknowledgments. This study was supported by JSPS KAKENHI (Grant No.

18K06082).

Conflict of interest The authors declare no conflict of interest.

REFERENCES

- 1) Brooks P, Emery P, Evans JF, Fenner H, Hawkey CJ, Patrono C, Smolen J, Breedveld F, Day R, Dougados M, Ehrich EW, Gijon-Baños J, Kvien TK, Van Rijswijk MH, Warner T, Zeidler H. Interpreting the clinical significance of the differential inhibition of cyclooxygenase-1 and cyclooxygenase-2. *Rheumatology (Oxford)*, **38**, 779–788 (1999).
- 2) Vane J. Towards a better aspirin. *Nature*, **367**, 215–216 (1994).
- 3) Sekiguchi H, Inoue G, Nakazawa T, Imura T, Saito W, Uchida K, Miyagi M, Takahira N, Takaso M. Loxoprofen sodium and celecoxib for postoperative pain in patients after spinal surgery: a randomized comparative study. *J. Orthop. Sci.*, **20**, 617–623 (2015).
- 4) Graham DY, Opekun AR, Willingham FF, Qureshi WA. Visible small-intestinal mucosal injury in chronic NSAID users. *Clin. Gastroenterol. Hepatol.*, **3**, 55–59 (2005).
- 5) Yokoyama S, Hiramoto K, Koyama M, Ooi K. Skin disruption is associated with indomethacin-induced small intestinal injury in mice. *Exp. Dermatol.*, **23**, 659–663 (2014).

- 322 6) Gruber BL, Schwartz LB, Ramamurthy NS, Irani AM, Marchese MJ. Activation of
323 latent rheumatoid synovial collagenase by human mast cell tryptase. *J. Immunol.*,
324 **140**, 3936–3942 (1988).
- 325 7) Issemann I, Green S. Activation of a member of the steroid hormone receptor
326 superfamily by peroxisome proliferators. *Nature*, **347**, 645–650 (1990).
- 327 8) Ricote M, Li AC, Willson TM, Kelly CJ, Glass CK. The peroxisome
328 proliferator-activated receptor- γ is a negative regulator of macrophage activation.
329 *Nature*, **391**, 79–82 (1998).
- 330 9) Jiang C, Ting AT, Seed B. PPAR- γ agonists inhibit production of monocyte
331 inflammatory cytokines. *Nature*, **391**, 82–86 (1998).
- 332 10) Kuzumoto T, Tanigawa T, Higashimori A, Kitamura H, Nadatani Y, Otani K,
333 Fukunaga S, Hosomi S, Tanaka F, Kamata N, Nagami Y, Taira K, Watanabe T,
334 Fujiwara Y. Protective role of resolvin D1, a pro-resolving lipid mediator, in
335 nonsteroidal anti-inflammatory drug-induced small intestinal damage. *PLoS One*, **16**,
336 e0250862 (2021).
- 337 11) Shimada N, Sakata A, Igarashi T, Takeuchi M, Nishimura S. M1 macrophage
338 infiltration exacerbate muscle/bone atrophy after peripheral nerve injury. *BMC*
339 *Musculoskelet. Disord.*, **21**, 44 (2020).

- 340 12) Liang S, Shi L-Y, Duan J-Y, Liu H-H, Wang T-T, Li C-Y. Celecoxib reduces
341 inflammation and angiogenesis in mice with adenomayosis. *Am. J. Transl. Res.*, **13**,
342 2858-1866 (2021).
- 343 13) Chen J, Yang H, Sheng Z. Ellagic acid activated PPAR signaling pathway to
344 protect ileums against castor oil-induced diarrhea in mice: application of
345 transcriptome analysis in drug screening. *Front. Pharmacol.*, **10**, 1681 (2020).
- 346 14) Washizaki K, Kanto H, Yoshida K, Chung L, Takada Y, Motomura A, Nomura S,
347 Akatsuka M, Ishikawa A. The objective technique for evaluating of dry skin
348 symptom in the skin disease. *J. Med. Soc. Toho Univ.*, **63**, 39-42 (2016).
- 349 15) Ohno H. Effects of ionized water nanodroplets (mist) released from an
350 air-conditioner on the biophysical and biomechanical propertyes of skin in low
351 humidity rooms. *Jpn. J. Biometeor.*, **53**, 113-121 (2016).
- 352 16) Piao X, Miura R, Miyake S, Komazawa-Sakon S, Koike M, Shindo R, Takeda J,
353 Hasegawa A, Abe R, Nishimura C, Mikami T, Yagita H, Uchiyama Y, Nakao H.
354 Blockade of TNF receptor superfamily 1 (TNFR1)-dependent and
355 TNFR1-independent cell death is crucial for normal epidermal differentiation. *J.*
356 *Allergy Clin. Immunol.*, **143**, 213-228 (2019).
- 357 17) Barel AO, Clarys P. Study of the stratum corneum barrier function by

358 transepidermal water loss measurements: comparison between two commercial
359 instruments: Evaporimeter and Tewameter. *Skin Pharmacol.*, **8**, 186–195 (1995).

360 18) Berardesca E, European Group for Efficacy Measurements on Cosmetics and
361 Other Topical Products (EEMCO). EEMCO guidance for the assessment of stratum
362 corneum hydration: electrical methods. *Skin Res. Technol.*, **3**, 126–132 (1997).

363 19) Lodén M, Olsson H, Axéll T, Linde YW. Friction, capacitance and transepidermal
364 water loss (TEWL) in dry atopic and normal skin. *Br. J. Dermatol.*, **126**, 137–141
365 (1992).

366 20) Varani J, Warner RL, Gharaee-Kermani M, Phan SH, Kang S, Chung JH, Wang ZQ,
367 Datta SC, Fisher GJ, Voorhees JJ. Vitamin A antagonizes decreased cell growth and
368 elevated collagen-degrading matrix metalloproteinases and stimulates collagen
369 accumulation in naturally aged human skin. *J. Invest. Dermatol.*, **114**, 480–486
370 (2000).

371 21) Iddamalgoda A, Le QT, Ito K, Tanaka K, Kojima H, Kido H. Mast cell tryptase and
372 photoaging: possible involvement in the degradation of extra cellular matrix and
373 basement membrane proteins. *Arch. Dermatol. Res.*, **300**, S69–S76 (2008).

374 22) Yokoyama S, Hiramoto K, Koyama M, Ooi K. Impaired skin barrier function in
375 mice with colon carcinoma induced by azoxymethane and dextran sodium sulfate.

- 376 *Biol. Pharm. Bull.*, **38**, 947–950 (2015).
- 377 23) Bax HJ, Keeble AH, Gould HJ. Cytokinergic IgE Action in Mast Cell Activation.
378 *Front Immunol.*, **6**, 229 (2012).
- 379 24) Metcalfe DD, Baram D, Mekori YA. Mast cells. *Physiol. Rev.*, **77**, 1033–1079
380 (1997).
- 381 25) Higuchi K, Yoda Y, Amagase K, Kato S, Tokioka S, Murano M, Takeuchi K,
382 Umegaki E. Prevention of NSAID-Induced Small Intestinal Mucosal Injury:
383 Prophylactic Potential of Lansoprazole. *J. Clin. Biochem. Nutr.*, **45**, 125–130 (2009).
- 384 26) Coppelli G, Guaita E, Spaggiari S, Coruzzi G. Gastric effects of the selective
385 cyclooxygenase-2 inhibitor, celecoxib, in the rat. *Dig. Liver Dis.*, **36**, 265–270
386 (2004).
- 387 27) Tegeder I, Pfeilschifter J, Geisslinger G. Cyclooxygenase-independent actions of
388 cyclooxygenase inhibitors. *FASEB J.*, **15**, 2057–2072 (2001).
- 389 28) Maeyama K, Emi M, Chihara K. Role of PPAR γ in inflammatory response related
390 to mast cells. *Nihon Yakurigaku Zasshi*, **122**, 325–330 (2003).
- 391 29) Bouhlef MA, Derudas B, Rigamonti E, Dièvert R, Brozek J, Haulon S, Zawadzki C,
392 Jude B, Torpier G, Marx N, Staels B, Chinetti-Gbaguidi G. PPAR γ activation primes
393 human monocytes into alternative M2 macrophages with anti-inflammatory

- 394 properties. *Cell Metab.*, **6**, 137–143. (2007).
- 395 30) Odegaard JI, Ricardo-Gonzalez RR, Goforth MH, Morel CR, Subramanian V,
396 Mukundan L, Eagle AR, Vats D, Brombacher F, Ferrante AW, Chawla A.
397 Macrophage-specific PPAR γ controls alternative activation and improves insulin
398 resistance. *Nature*, **447**, 1116–1120 (2007).
- 399 31) Engeroff P, Fellmann M, Yerly D, Bachmann MF, Vogel M. A novel recycling
400 mechanism of native IgE-antigen complexes in human B cells facilitates transfer of
401 antigen to dendritic cells for antigen presentation. *J. Allergy Clin. Immunol.*, **142**,
402 557–568.e6 (2018).
- 403 32) Bonnefoy JY, Gauchat JF, Life P, Graber P, Aubry JP, Lecoanet-Henchoz S.
404 Regulation of IgE synthesis by CD23/CD21 interaction. *Int. Arch. Allergy Immunol.*,
405 **107**, 40–42 (1995).
- 406 33) Ashida Y, Denda M, Hirao T. Histamine H1 and H2receptor antagonists accelerate
407 skin barrier repair and prevent epidermal hyperplasia induced by barrier disruption
408 in a dry environment. *J. Invest. Dermatol.*, **116**, 261-265 (2001).
- 409 34) Wood LC, Elias PM, Sequeira-Martin SM, Grunfeld C, Feingold KR. Occlusion
410 lowers cytokine mRNA levels in essential fatty acid-deficient and normal mouse
411 epidermis, but not after acute barrier disruption. *J. Invest. Dermatol.*, **103**, 834-838

(1994).

35) Kim BE, Howell MD, Guttman-Yassky E, Gilleaudeau PM, Cardinale IR, Boguniewicz M, Krueger JG, Leung DY. TNF- α downregulates filaggrin and loricrin through c-Jun N-terminal kinase: role for TNF- α antagonists to improve skin barrier. *J. Invest. Dermatol.*, **131**, 1272-1279 (2011).

36) Demerjian M, Man M-Q, Choi E-H, Brown BE, Crumrine D, Cang S, Mauro T, Elias PM, Feingold KR. Topical treatment with thiazolidinediones, activators of peroxisome proliferator-activated receptor-gamma, normalizes epidermal homeostasis in a murine hyperproliferative disease model. *Exp. Dermatol.*, **15**, 154-160 (2006).

Fig. 1. Changes in the skin condition of mice treated with non-steroidal anti-inflammatory drugs (NSAIDs). Measurement of transepidermal water loss (TEWL) (A), capacitance (B), histological damage (C), and the expression of Collagen type I (D) and MMP-1 (E) in the dorsal skin of mice treated with indomethacin, loxoprofen sodium, or celecoxib. Values represent the mean $\pm$ standard deviation. * $p < 0.05$. Capacitance is represented in arbitrary units (a.u.). Scale bar = 100 μ m.

a

442

443

b

446

447

Fig. 2. Effect of non-steroidal anti-inflammatory drug (NSAID) treatment on the levels of mast cells (A, B, C, D) and histamine (E, F) in the skin and jejunum. The number of mast cells in the control and NSAID-treated mice was counted following toluidine blue staining of the skin specimens. Values represent the mean $\pm$ standard deviation. * $p < 0.05$. Scale bar = 100 μm .

Fig. 3. Effect of non-steroidal anti-inflammatory drug treatment on the expression of peroxisome proliferation-activated receptor (PPAR) γ and p-PPAR γ in the skin and jejunum. Values represent the mean $\pm$ standard deviation. * p < 0.05.

Fig. 4. Effect of non-steroidal anti-inflammatory drug treatment on the expression of CCR7 (A, B) and CD163 (C, D) in the skin and jejunum. CCR7 is a marker of M1 macrophages and CD163 is a marker of M2 macrophages. Values represent the mean $\pm$ standard deviation. * $p < 0.05$.

Fig. 5. Effect of non-steroidal anti-inflammatory drug treatment on the plasma levels of interleukin (IL)-10 (A) and immunoglobulin (Ig)E (B), the expression of CD23 in the skin (C, D), and the expression of CD23 and CD21 in the skin and jejunum (E). Values represent the mean $\pm$ standard deviation. * $p < 0.05$. Scale bar = 100 μm .

Fig. 6. Effect of GW9662 (peroxisome proliferation-activated receptor (PPAR) γ antagonist) treatment on the transepidermal water loss (TEWL) (A) and expression of PPAR γ and p-PPAR γ in the skin (B, C). Values represent the mean $\pm$ standard deviation.

* $p < 0.05$.

508

509

510 Figure 7. Mechanism underlying indomethacin treatment and skin dryness in mice.