Letter to the Editor

Japanese Criteria of Metabolic Syndrome

To the Editor:

We appreciate the labor-some quadratic regression analyses by Dr Matsuzawa.¹ However, we are afraid that our intention was not properly conveyed to the author. He stated in Author's Reply to our letter¹ that there were minor differences between quadratic and linear regression analyses in order to determine the optimal cut-points of waist circumference (WC) to discriminate persons with visceral fat area (VFA) $\geq 100\text{cm}^2$ from those with VFA $< 100\text{cm}^2$ and that the Japanese cut-points of WC² are the only ones supported by evidence.¹ However, he mentioned neither receiver operating characteristic (ROC) curves, nor areas under ROC curves (AUC). If he did ROC and AUC analyses, WC might be revealed as a poor discriminator of VFA especially in women, even if these 2 factors are well correlated. After all, WC is a marker of abdominal (central) obesity, not of visceral obesity (which is assessed by VFA using CT scanning, exposing subjects to irradiation). At least he should reveal the sensitivities and specificities of the cut-points of WC as a discriminator of VFA. Without his revelation, one can easily see in his VFA-WC distribution graphs² that the sensitivity of the female cut-point of 90 cm is less than 50%. Furthermore, he did not separately analyze the optimal cut-points of VFA by gender. If he did so, the optimal cut-point of VFA for women might be lower than that for men, because more abundant subcutaneous fat in women than in men may also contribute to multiple risk factors, even if the degree of the contribution is less than that of visceral fat. Therefore, this female cut-point of WC (90 cm) is not acceptable.

Regarding criteria of metabolic syndrome, the American Heart Association/National Heart, Lung, and Blood Institute (AHA/NHLBI) criteria³ stand on the concept of a syndrome resulting from multiple causes, the International Diabetes Federation (IDF) criteria⁴ stand on the concept of an entity due to central obesity, and the Japanese criteria⁵ intended to, but do not really, stand on the concept of an entity due to visceral obesity, and excluded women with HDL cholesterol levels of 40–49 mg/dl, which constitutes a significant risk category for both coronary events and total mortality in Japan.⁶ The American Diabetes Association/European Association for the Study of Diabetes jointly stated that doctors should not label patients with metabolic syndrome because the concept is premature as a syndrome. And Reaven called off debating the pros and cons of various definitions of metabolic syndrome.⁸ Nevertheless, no one can stop endless debates upon new definitions of it^{9–12} except in Japan where the Japanese criteria⁵ was forced upon people ex cathedra. Therefore, we wish the Examination Committee of Criteria for Metabolic Syndrome would

withdraw their unique Japanese criteria of metabolic syndrome⁵ based on no prospective prognostic evidence and really cooperate internationally. At least one should understand that this Japanese definition is not the only one based on prospective prognostic evidence, but one of many arbitrary premature definitions⁷ and the concept of which differs from those of 2 recently published worldwide AHA/NHLBI³ and IDF⁴ definitions, thus hindering global cooperation, contrary to the intention of the Committee.⁵

References

1. Matsuzawa Y. Author's reply; Oda E. Letter. Cutoff point of waist circumference in Japan. *Circ J* 2006; **70**: 150.
2. The Examination Committee of Criteria for "Obesity Disease" in Japan: New Criteria for "Obesity Disease" in Japan. *Circ J* 2002; **66**: 987–992.
3. Grundy SM, Cleeman JJ, Daniels SR, Donato KA, Eckel RH, Franklin BA, et al. Diagnosis and Management of the Metabolic Syndrome: An American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation* 2005; **112**: 2735–2752.
4. The IDF consensus worldwide definition of the metabolic syndrome. www.idf.org/home/index.cfm?unode=1120071E-AACE-41D2-9FA0-BAB6E25BA072
5. The Examination Committee of Criteria for Metabolic Syndrome. Definition and Criteria of Metabolic Syndrome. *J Jpn Soc Int Med* 2005; **94**: 794–809 (in Japanese).
6. Matsuzaki M, Kita T, Mabuchi H, Matsuzawa Y, Nakaya N, Oikawa S, et al. Large scale cohort study of the relationship between serum cholesterol concentration and coronary events with low-dose simvastatin therapy in Japanese patients with hypercholesterolemia: Primary prevention cohort study of the Japan Lipid Intervention Trial (J-LIT). *Circ J* 2002; **66**: 1087–1095.
7. Kahn R, Ferrannini E, Buse J, Stern M. The metabolic syndrome: Time for a critical appraisal: Joint statement from the American Diabetes Association and the European Association for the Study of Diabetes. *Diabetes Care* 2005; **28**: 2289–2304.
8. Reaven G. Insulin resistance, type 2 diabetes mellitus, and cardiovascular disease: The end of the beginning. *Circulation* 2005; **112**: 3030–3032.
9. Balkau B, Eschwege E. The metabolic syndrome: Correspondence. *Lancet* 2005; **366**: 1921.
10. Kahn R, Buse J, Ferrannini E, Stern M. The metabolic syndrome: Correspondence. *Lancet* 2005; **366**: 1921–1922.
11. Jarret RJ. The metabolic syndrome: Correspondence. *Lancet* 2005; **366**: 1922.
12. Neuhauser HK. The metabolic syndrome: Correspondence. *Lancet* 2005; **366**: 1922–1923.

Eiji Oda, MD

Department of Internal Medicine,
Niigata Prefectural Yoshida Hospital,
Niigata, Japan

Kenichi Watanabe, MD

Department of Clinical Pharmacology,
Niigata University of Pharmacy
and Applied Life Sciences,
Niigata, Japan