CORRIGENDUM

Clinical Significance of High-Molecular Weight Form of Adiponectin in Male Patients With Coronary Artery Disease

Noriyuki Komura, MD; Shinji Kihara, MD, PhD; Mina Sonoda, BS; Masahiro Kumada, MD, PhD; Koichi Fujita, MD; Aki Hiuge, MD; Takuya Okada, MD; Yasuhiko Nakagawa, MD; Sachiko Tamba, MD; Yohei Kuroda, MD; Noriyuki Hayashi, MD; Satoru Sumitsuji, MD; Toshiharu Kawamoto, MD, PhD; Satoru Matsumoto, MD; Noriyuki Ouchi, MD, PhD; Yukio Arita, MD, PhD; Yoshihisa Okamoto, MD, PhD; Ichiro Shimomura, MD, PhD; Tohru Funahashi, MD, PhD; Yuji Matsuzawa, MD, PhD for the Osaka CAD Group

(*Circ J* 2008; **72**: 23–28)

The author (N. Komura) apologizes for errors which appeared in the Table 1, page 25, Vol. 72. The numbers and proportion of hypertension or current smokers were 72 (55%), 61 (47%) in non-diabetic patients, and 39 (70%), 26 (46%) in diabetic patients, respectively. The corrected Table 1 is given below.

Table 1 Clinical and Biochemical Characteristics

	Non-diabetic (n=130)	Diabetic (n=56)	p value
Age (years)	64.2±0.9	63.2±1.1	0.532
BMI (kg/m ²)	24.3±0.3	24.4±0.4	0.920
Total adiponectin (μg/ml)	6.3±0.3	5.5±0.4	0.094
HMW adiponectin (μg/ml)	5.3±0.3	4.6±0.4	0.161
Fasting glucose (mg/dl)	98.9±1.0	144.2±4.5	<0.0001
Hemoglobin A _{1c} (%)	5.3±0.1	6.9±0.2	<0.0001
Dyslipidemia, n (%)	76 (58)	33 (59)	0.953
TC (mg/dl)	201.3±3.4	195.7±4.7	0.346
TG (mg/dl)	147.0±7.5	146.4±10.5	0.961
HDL-C (mg/dl)	45.3±1.2	45.6±2.0	0.909
LDL-C (mg/dl)	126.6±2.8	120.8±4.1	0.246
Hypertension, n (%)	72 (55)	39 (70)	0.069
SBP (mmHg)	131.7±1.7	135.1±2.8	0.282
DBP (mmHg)	75.0±1.1	74.5±1.5	0.773
Current smokers, n (%)	61 (47)	26 (46)	0.951

Values are presented as n, or as the mean ± SEM.

BMI, body mass index; HMW, high molecular weight; TC, total cholesterol; TG, triglyceride; HDL-C, high-density lipoprotein-cholesterol; LDL-C, low-density lipoprotein-cholesterol; SBP, systolic blood pressure; DBP, diastolic blood pressure.