Sex-Related Differences in the Risk Factor Profile and Medications of Patients With Atrial Fibrillation Recruited in J-TRACE

Hiroshi Inoue, MD¹; Takashi Nozawa, MD¹; Tadakazu Hirai, MD¹; Shinya Goto, MD²; Hideki Origasa, PhD³; Kazuyuki Shimada, MD⁴; Shinichiro Uchiyama, MD⁵; Takayuki Hirabayashi, MD⁶; Yukihiro Koretsune, MD⁷; Shiro Ono, MD⁸; Tooru Hasegawa, MD⁹; Yasuo Sasagawa, MD¹⁰; Yoshiaki Kaneko, MD¹¹; Yasuo Ikeda, MD¹² for the J-TRACE Investigators

Background: Clinical characteristics, including risk factors for thromboembolism, and medications differ between men and women with atrial fibrillation (AF) in Western countries. Whether such a difference exists for Japanese patients with AF is unclear, so data from J-TRACE were used to investigate this issue.

Methods and Results: A total of 2,892 patients (2,028 men, 864 women; 70.3 years old) with AF were analyzed for the respective prevalences of risk factors and medications. CHADS2 score was calculated to determine thromboembolic risk level. Women were older ($P<0.001$), and more frequently had heart failure ($P<0.001$), and hypertension ($P=0.051$) than men. The proportion of subjects aged 75 years or older was higher among women than among men ($P<0.001$). CHADS2 score was therefore significantly higher in women than in men (2.05 ± 1.29 vs 1.88 ± 1.33 , $P<0.001$). Sex-related differences were not observed for the prevalence of diabetes mellitus, myocardial infarction or ischemic stroke, nor did warfarin usage differ between men and women.

Conclusions: Sex-related differences were observed in the risk factor profile and medications of Japanese patients with AF. CHADS2 score was higher in women than in men. (*Circ J* 2010; **74**: 650–654)

Key Words: Atrial fibrillation; CHADS2 score; Clinical characteristics; Medications; Sex differences

Atrial fibrillation (AF) is a common cardiac arrhythmia seen in general practice as well as in the cardiology clinic. The prevalence of AF differs between men and women in Western countries,^{1–3} and also in Japan.^{4,5} Several studies have reported that there are sex-related differences in the clinical characteristics and medications of patients with AF.^{6–10} A prospective, cohort study indicated that the effects of AF on the risk of stroke were greater in women than in men after adjustment for age and comorbidity.⁹ Other studies also showed that AF is associated with an increase in cardiovascular events, including mortality and stroke, especially in women.^{7,11,12} Some risk stratification schemes consider women to be at high risk for ischemic stroke,^{13,14} while others do not.^{15,16} However, because the sex-related differences in risk factors for cardiovascular dis-

eases and medications of Japanese patients with AF have yet to be clarified, registry data for a large, nation-wide, multi-center, cooperative study, J-TRACE (The Japan Thrombosis Registry for Atrial Fibrillation, Coronary or Cerebrovascular Events),^{17,18} were analyzed to address this issue in the present study.

Methods

The details of J-TRACE have been reported elsewhere.^{17,18} Briefly, J-TRACE has a steering committee of 5 members and 41 regional coordinators selected from 10 regions of Japan (Appendix 1). Recruitment of patients to investigate risk factor profiles and current status of medications for risk factors and for prevention of cardiovascular events in patients with

Received November 4, 2009; revised manuscript received December 18, 2009; accepted December 21, 2009; released online February 20, 2010 Time for primary review: 19 days

¹Department of Internal Medicine, University of Toyama, Toyama, ²Department of Internal Medicine, Tokai University, Isehara, ³Department of Biostatistics, University of Toyama, Toyama, ⁴Department of Cardiology, Jichi Medical University, Shimotsuke, ⁵Department of Neurology, Tokyo Women's Medical University, Tokyo, ⁶Department of Cardiology, Sunagawa City Medical Center, Sunagawa, ⁷Institute for Clinical Research, Osaka National Hospital, Osaka, ⁸Department of Cardiology, Saiseikai Yamaguchi Hospital, Yamaguchi, ⁹Department of Cardiology, Hakodate Medical Association Hospital, Hakodate, ¹⁰Sasagawa Clinic, Niigata, ¹¹Medical & Biological Science, Gunma University, Maebashi and ¹²Department of Internal Medicine, Keio University, Tokyo, Japan

Mailing address: Hiroshi Inoue, MD, The Second Department of Internal Medicine, University of Toyama, 2630 Sugitani, Toyama 930-0194, Japan. E-mail: hiroshi@med.u-toyama.ac.jp

ISSN-1346-9843 doi:10.1253/circj.CJ-09-0802

All rights are reserved to the Japanese Circulation Society. For permissions, please e-mail: cj@j-circ.or.jp

Table 1. Clinical Characteristics of Japanese Patients With AF

	Men (n=2,028)	Women (n=864)	P value
Age (years)	69.4±9.4	72.6±8.5	<0.001
≥75 years (%)	32.0	44.5	<0.001
Chronic AF* (%)	68.8 (1,062/1,543)	66.1 (462/699)	0.199
BMI (kg/m ²)	23.8±3.2	23.4±4.1	<0.001
CHF (%)	17.0	27.1	<0.001
Hypertension (%)	57.2	61.1	0.051
DM (%)	19.1	16.7	0.125
Ischemic stroke (%)	29.4	26.3	0.089
VHD (%)	10.1	21.1	<0.001
MI (%)	7.6	5.9	0.096
HC (%)	25.1	35.5	<0.001
Drinker (%)	46.3	5.2	<0.001
Smoker (%)	21.2	4.3	<0.001
CHADS2 score	1.88±1.33	2.05±1.29	<0.001

Data are mean±SD or % of patients.

*In the myocardial infarction and stroke categories; subtypes of AF were not specifically determined.

AF, atrial fibrillation; BMI, body mass index; CHF, congestive heart failure; DM, diabetes mellitus; MI, myocardial infarction; VHD, valvular heart diseases including valve replacement; HC, hypercholesterolemia.

prior stroke, myocardial infarction (MI) or AF began in January 2005 and ceased in December 2006.

Study Population

Patients aged 20–90 years were eligible for enrollment if they had at least 1 of the 3 cardiovascular diseases (stroke, MI or AF). The study protocol was approved by an Institutional Review Board at each participating site and all patients gave informed consent. Those in the AF category, and those in the stroke and MI categories who also had AF, comprised the study subjects for this subanalysis of J-TRACE. Those in the recovery phase of acute MI or acute stroke were not eligible for enrollment in J-TRACE.

Baseline Characteristics

All subtypes of AF were included. AF was diagnosed electrocardiographically using standard diagnostic criteria. Risk factors and comorbidities were collected from the medical record as baseline data. Among them were hypertension, diabetes mellitus, hypercholesterolemia, valvular diseases, MI, ischemic stroke, congestive heart failure, smoking, and drinking. Regular use of medications, including anticoagulants, antiplatelet agents, and drugs for hypercholesterolemia, hypertension, and diabetes mellitus, was also determined from the medical record. Each patient's CHADS2 score¹⁵ was calculated to determine the level of cardioembolic risk: 1 point was given for advanced age (≥75 years), hypertension, congestive heart failure, or diabetes mellitus, and 2 points for prior stroke or transient ischemic attack.

Statistical Analysis

Continuous variables are shown as the mean±SD, and categorical variables as percentages. Continuous variables were compared by analysis of variance or Student's t-test, and categorical variables with the chi-square test, with $P<0.05$ considered significant.

Table 2. Distribution of CHADS2 Scores

CHADS2 score	Men	Women
0	15.9	11.1
1	28.4	25.0
2	23.5	29.6
3	19.2	20.6
4	10.5	10.0
5	2.2	3.4
6	0.3	0.3

Figures are % of patients.

$P<0.001$ between men and women.

Table 3. Age and CHADS2 Score

	Age			P value
	<65 years	65–74 years	≥75 years	
Men	1.24±1.12 (n=572)	1.63±1.22 (n=808)	2.74±1.17 (n=648)	<0.001
Women	1.38±1.16 (n=153)	1.57±1.14 (n=326)	2.72±1.12 (n=385)	<0.001

Data are mean±SD.

Results

Risk Factor Profile

A total of 2,892 patients (2,028 men, 864 women; mean age, 70.3 years) with AF comprised the study group. Numbers of patients and their mean age in the 3 categories were as follows: AF category, 1,543 men (68.9±9.6 years old) and 699 women (72.4±8.5); stroke category, 399 men (70.6±8.4) and 141 women (73.0±8.3); MI category, 86 men (71.7±8.1) and 24 women (75.3±8.0). Their clinical characteristics are summarized in **Table 1**. Some of the characteristics exhibited differences by sex. Women were older ($P<0.001$), and more frequently had congestive heart failure ($P<0.001$), hypertension ($P=0.051$), valvular diseases or valve replacement ($P<0.001$), and hypercholesterolemia ($P<0.001$) than the men, but drank ($P<0.001$) and smoked ($P<0.001$) less frequently than men. The proportion of subjects aged 75 years or older was higher and body mass index was slightly but significantly lower in women than in men ($P<0.001$, each case). The prevalences of chronic AF, diabetes mellitus, MI, and ischemic stroke did not differ between men and women.

The CHADS2 score was slightly but significantly higher in women than in men (**Table 1**, $P<0.001$) because of their higher prevalence of older age (≥75 years), hypertension, and congestive heart failure. The distribution of CHADS2 scores differed significantly between men and women (**Table 2**, $P<0.001$). It increased with age for both men and women, but did not differ between men and women in any age group (**Table 3**).

Medications

Medications are summarized in **Table 4**. Use of warfarin and antiplatelet agents did not differ between men and women. Reflecting the differences in prevalence of hypertension and hypercholesterolemia between men and women, drugs for the treatment of these diseases were used more frequently in women than in men ($P<0.001$, each case). In contrast, use of antidiabetic drugs was similar in men and women.

There were no apparent sex-related differences in the rate of use of warfarin or aspirin at any CHADS2 score (**Table 5**).

Table 4. Medications at Baseline

	Men	Women	P value
Warfarin	73.1	72.7	0.807
Antiplatelet agents	37.9	36.0	0.328
Aspirin	32.1	30.8	0.504
Ticlopidine	5.0	5.0	0.316
Cilostazol	2.0	1.3	0.191
Antihypertensives	71.8	78.8	<0.001
ACEI	17.4	14.8	0.087
ARB	28.4	32.2	0.039
β -blockers	21.4	21.3	0.927
Calcium antagonists	36.4	42.5	0.002
Diuretics	18.6	33.4	<0.001
Lipid-lowering drugs	16.7	26.4	<0.001
Statins	14.9	23.7	<0.001
Antidiabetic drugs	10.6	10.9	0.825
Oral	8.7	8.6	0.921
Insulin	1.4	2.2	0.111

Data are % of patients.

Only major drugs for treatment of comorbidities and prevention of thromboembolism are listed (see Uchiyama et al¹⁸ for more detailed information on medications in J-TRACE).

ACEI, angiotensin converting enzyme inhibitors; ARB, angiotensin II receptor blockers.

Warfarin usage differed significantly among CHADS2 scores in both men ($P<0.001$) and women ($P=0.001$). It increased gradually from approximately 60% to 80% as the score increased from 0 to 3 for both men and women; thereafter it reached a plateau, except in the case of women with a score of 6. Aspirin usage also differed significantly among CHADS2 scores in men ($P=0.008$), but not in women ($P=0.852$). It did not show any apparent score-dependent increase as observed in the case of warfarin usage.

Discussion

The major findings of the present study are as follows. First, there were sex-related differences in the risk factor profile and medications of patients with AF recruited in J-TRACE. Women were older and more frequently had hypertension, valvular diseases, congestive heart failure, and hypercholesterolemia than men. The prevalence of diabetes mellitus, ischemic stroke, and MI did not differ between men and women. Second, CHADS2 score was consequently slightly but significantly higher in women than in men with AF. This sex-related difference could be largely related to the higher proportion of women aged 75 years or older. Third, no sex-related differences in the use of warfarin or aspirin were observed at any CHADS2 score.

Risk Factor Profile of Patients With AF

Reports from Western countries^{6–10} suggest that sex-related differences could exist in the risk factors for cardiovascular diseases of patients with AF. In the present study, mean age was higher and the prevalence of hypertension also tended to be higher in women than in men, consistent with the previous reports;^{6–9} however, the prevalence of congestive heart failure was also higher in women than in men in the present study, a finding that is inconsistent with those reports from Western countries.^{6–9} Notably, the prevalence of diabetes mellitus and of a prior history of ischemic stroke were not

Table 5. Use of Warfarin and Aspirin at Each CHADS2 Score

CHADS2 score	Warfarin use (%)		Aspirin use (%)	
	Men	Women	Men	Women
0	57.9	61.4	32.3	28.1
1	68.9	66.2	33.9	31.4
2	74.8	73.8	30.6	32.0
3	84.9	77.0	26.9	30.3
4	81.1	87.2	34.4	29.1
5	77.3	79.3	31.3	27.6
6	83.3	66.7	83.3	66.7
P value	<0.001	0.001	0.008	0.852

consistent.^{6–9}

Cohort studies of the general population in Japan have indicated that the prevalences of hypertension and diabetes mellitus are higher in men than in women.^{19–21} The prevalence of cardiac diseases was not higher in women than in men with AF,^{4,5} so the higher prevalences of hypertension and congestive heart failure in women with AF found in the present study do not simply reflect the prevalence of these diseases in the general population of Japan. Valvular disease is a well-known risk factor for AF,²² especially for Japanese women.²³ Drinking and smoking could promote the development of AF,^{22–25} and were present more frequently in men than in women in the present study, as in the general population of Japan.^{4,5,19–21} The electrophysiological properties of the atria differ between men and women,²⁶ so greater comorbidity and age might be required for AF to develop in women than in men.

Thromboembolic Risk

A sex difference in CHADS2 score was found in the present study, a finding consistent with the ATRIA study.⁷ In the Euro Heart Survey the score might have been higher in women than in men, because mean age and the prevalences of hypertension, diabetes mellitus, and prior ischemic stroke were significantly higher in women than in men.⁹ In some studies the levels of biomarkers of a prothrombotic state were higher in women with AF than in men with AF.^{27,28} These findings could explain the inclusion of female sex as a risk factor in some schemes for predicting thromboembolic events in patients with AF.^{13,14} In fact, among patients with acute stroke, embolic infarction is observed more frequently in women than in men.²⁹ It is difficult to determine the reasons for the sex-related difference in thromboembolic risk; however, some components of the CHADS2 score were observed more frequently in women in the ATRIA study,⁷ Euro Heart Survey,⁹ and in the present study.

Medications

Registry studies in Western countries have indicated that warfarin usage does not differ between men and women.^{6,9} In the present study, the rate of warfarin usage did not differ between men and women as a whole nor did it differ between them at any CHADS2 score (Table 5). Warfarin usage is at present not necessarily less frequent in women than in men, as reported in earlier registry⁶ and community-based cohort³⁰ studies.

Use of aspirin and antidiabetic drugs was similar in men and women; however, drugs for hypertension and hypercholesterolemia were used more frequently by women than by men. The latter finding might reflect the sex-related differ-

ences in the prevalence of these diseases in the present study.

Study Limitations

First, enrollment of consecutive patients with stroke, MI, and AF was recommended, but may not necessarily have occurred at each participating site and this possible selection bias could have affected the present results. Second, data for subjects with AF were collected from 3 categories of J-TRACE^{17,18} possibly resulting in increased prevalences of ischemic stroke and MI. However, this might not necessarily have affected sex-related differences in the frequency of these diseases in the present study. Actually, when only patients of AF category were analyzed, the results did not differ in terms of sex-related differences in mean age, CHADS2 score, and prevalences of heart failure, hypertension, smoking, drinking habit and warfarin usage (data not shown). Third, the study design of the J-TRACE did not define the diagnostic criteria of comorbidities, including hypertension, hypercholesterolemia and others; however, data of comorbidities were collected from the medical record. If strict diagnostic criteria of comorbidities were used, the present results would not have changed greatly. Finally, the intensity of anticoagulation was not determined systematically, and follow-up data are not yet available.

Clinical Implications

Our findings indicate sex-related differences in the clinical risk factor profile of patients with AF, with the CHADS2 score slightly but significantly higher in women with AF than in men with AF in the clinical setting in Japan. Further follow-up studies are required to elucidate the effects of these sex-related differences on subsequent thromboembolic events.

Acknowledgment

This study was supported by the Japan Heart Foundation, Tokyo, Japan.

Disclosure

There is no conflict of interest to declare.

References

1. Feinberg WM, Blackshear JL, Laupacis A, Kronmal R, Hart RG. Prevalence, age distribution, and gender of patients with atrial fibrillation. *Arch Intern Med* 1995; **155**: 469–473.
2. Go AS, Hylek EM, Phillips KA, Chang YC, Henault LE, Selby JV, et al. Prevalence of diagnosed atrial fibrillation in adults: National implications for rhythm management and stroke prevention: The AnTicoagulation and Risk factors In Atrial fibrillation (ATRIA) study. *JAMA* 2001; **285**: 2370–2375.
3. Majeed A, Moser K, Carroll K. Trends in the prevalence and management of atrial fibrillation in general practice in England and Wales, 1994–1998: Analysis of data from the general practice research database. *Heart* 2001; **86**: 284–288.
4. Inoue H, Fujiki A, Origasa H, Ogawa S, Okumura K, Kubota I, et al. Prevalence of atrial fibrillation in the general population of Japan: An analysis based on periodic health examination. *Int J Cardiol* 2009; **137**: 102–107.
5. Iguchi Y, Kimura K, Aoki J, Kobayashi K, Terasawa Y, Sakai K, et al. Prevalence of atrial fibrillation in community-dwelling Japanese aged 40 years or older in Japan: Analysis of 41,436 non-employee residents in Kurashiki-City. *Circ J* 2008; **72**: 909–913.
6. Humphries KH, Kerr CR, Connolly SJ, Klein G, Boone JA, Green M, et al. New-onset atrial fibrillation: Sex differences in presentation, treatment, and outcome. *Circulation* 2001; **103**: 2365–2370.
7. Fang MC, Singer DE, Chang Y, Hylek EM, Henault LE, Jensvold NG, et al. Gender differences in the risk of ischemic stroke and peripheral embolism in atrial fibrillation: The AnTicoagulation and Risk factors In Atrial fibrillation (ATRIA) study. *Circulation* 2005; **112**: 1687–1691.
8. Gombert-Maitland M, Wenger NK, Feyzi J, Lengyel M, Volgman AS, Petersen P, et al. Anticoagulation in women with non-valvular atrial fibrillation in the stroke prevention using an oral thrombin inhibitor (SPORTIF) trials. *Eur Heart J* 2006; **27**: 1947–1953.
9. Dagnes N, Nieuwlaet R, Vardas PE, Andersen D, Levy S, Cobbe S, et al. Gender-related differences in presentation, treatment, and outcome of patients with atrial fibrillation in Europe: A report from the Euro Heart Survey on atrial fibrillation. *J Am Coll Cardiol* 2007; **49**: 572–577.
10. Kerr CR, Humphries K. Gender-related differences in atrial fibrillation. *J Am Coll Cardiol* 2005; **46**: 1307–1308.
11. Friberg J, Scharling H, Gadsboll N, Truelsen T, Jensen GB. Comparison of the impact of atrial fibrillation on the risk of stroke and cardiovascular death in women versus men (the Copenhagen City Heart Study). *Am J Cardiol* 2004; **94**: 889–894.
12. Stewart S, Hart CL, Hole DJ, McMurray JJV. A population-based study of long-term risks associated with atrial fibrillation: 20-year follow-up of the Renfrew/Paisley study. *Am J Med* 2002; **113**: 359–364.
13. Hart R, Pearce L, McBride R, Rothbart R, Asinger R. Factors associated with ischemic stroke during aspirin therapy in atrial fibrillation: Analysis of 2012 participants in the SPAF I-III clinical trials. *Stroke* 1999; **30**: 1223–1229.
14. Wang TJ, Massaro JM, Levy D, Vasa RS, Wolf PA, D'Agostino RB, et al. A risk score for predicting stroke or death in individuals with new-onset atrial fibrillation in the community: The Framingham Heart Study. *JAMA* 2003; **290**: 1049–1056.
15. Gage BF, Waterman AD, Shannon W, Boechler M, Rich MW, Radford MJ. Validation of clinical classification schemes for predicting stroke: Results from the National Register of Atrial Fibrillation. *JAMA* 2001; **285**: 2864–2870.
16. Atrial Fibrillation Investigators. Risk factors for stroke and efficacy of antithrombotic therapy in atrial fibrillation: Analysis of pooled data from five randomized controlled trials. *Arch Intern Med* 1994; **154**: 1449–1457.
17. Origasa H, Goto S, Uchiyama S, Shimada K, Ikeda Y, on behalf of the J-TRACE Investigators. The Japan Thrombosis Registry for Atrial Fibrillation, Coronary or Cerebrovascular Events (J-TRACE): A nation-wide, prospective large cohort study; the study design. *Circ J* 2008; **72**: 991–997.
18. Uchiyama S, Shibata Y, Hirabayashi T, Mihara B, Hamashige N, Kitagawa K, et al for the J-TRACE Investigators. Risk factor profiles in patients with stroke, myocardial infarction, and atrial fibrillation: The Japan Thrombosis Registry for Atrial Fibrillation, Coronary or Cerebrovascular Events (J-TRACE). *J Stroke Cerebrovasc Dis* (in press).
19. Ohsawa M, Okayama A, Okamura T, Itai K, Nakamura M, Tanno K, et al for the NIPPON DATA80 Research Group. Mortality risk attributable to atrial fibrillation in middle-aged and elderly people in the Japanese general population: Nineteen-year follow-up in NIPPON DATA80. *Circ J* 2007; **71**: 814–819.
20. Kubo M, Kiyohara Y, Kato I, Tanizaki Y, Arima H, Tanaka K, et al. Trends in the incidence, mortality, and survival rate of cardiovascular disease in a Japanese community: The Hisayama Study. *Stroke* 2003; **34**: 2349–2354.
21. Iso H, Sato S, Kitamura A, Imano H, Kiyama M, Yamagishi K, et al. Metabolic syndrome and the risk of ischemic heart disease and stroke among Japanese men and women. *Stroke* 2007; **38**: 1744–1751.
22. Benjamin EJ, Levy D, Vaziri SM, D'Agostino RB, Belanger AJ, Wolf PA. Independent risk factors for atrial fibrillation in a population-based cohort: The Framingham Heart Study. *JAMA* 1994; **271**: 840–844.
23. Fujishima M. Cardiac diseases as risk for cerebrovascular diseases. *Junkanki Senmon-i* 1998; **6**: 19–26 (in Japanese).
24. Djousse L, Levy D, Benjamin EJ, Blease S, Russ A, Larson MG, et al. Long-term alcohol consumption and the risk of atrial fibrillation in the Framingham study. *Am J Cardiol* 2004; **93**: 710–713.
25. Heeringa J, Kors JA, Hofman A, van Rooij FJA, Witteman JCM. Cigarette smoking and risk of atrial fibrillation: The Rotterdam Study. *Am Heart J* 2008; **156**: 1163–1169.
26. Tse HF, Oral H, Pelosi F, Knight BP, Strickberger SA, Morady F. Effect of gender on atrial electrophysiologic changes induced by rapid atrial pacing and elevation of atrial pressure. *J Cardiovasc Electrophysiol* 2001; **12**: 986–989.
27. Conway DS, Heeringa J, Van der Kuip DA, Chin BS, Hofman A, Witteman JC, et al. Atrial fibrillation and the prothrombotic state in the elderly: The Rotterdam study. *Stroke* 2003; **34**: 413–417.
28. Feinberg WM, Pearce LA, Hart RG, Cushman M, Cornell ES, Lip GY, et al. Markers of thrombin and platelet activity in patients with

atrial fibrillation: Correlation with stroke among 1531 participants in the Stroke Prevention in Atrial Fibrillation III study. *Stroke* 1999; **30**: 2547–2553.

29. Appellos P, Stegmayr B, Terent A. Sex differences in stroke epidemiology: A systematic review. *Stroke* 2009; **40**: 1082–1090.
30. Sudlow M, Thomson R, Thwaites B, Rodgers H, Kenny RA. Prevalence of atrial fibrillation and eligibility for anticoagulants in the community. *Lancet* 1998; **352**: 1167–1171.

Appendix 1

J-TRACE Steering Committee

Yasuo Ikeda, Keio University School of Medicine (Chair); Kazuyuki Shimada, Jichi Medical School; Shinichiro Uchiyama, Tokyo Women's Medical University; Shinya Goto, Tokai University School of Medicine; and Hideki Origasa, University of Toyama School of Medicine and Pharmaceutical Sciences

Secretariat

Hiroko Usami, BIOMEDIS INTERNATIONAL LTD

Regional Coordinators

Hokkaido Kiyohiro Houkin, Jyoji Nakagawara, Kazuaki Shimamoto, Hiroyuki Tsutsui

Tohoku Kunio Shirato, Hiroaki Shimokawa, Akifumi Suzuki, Yasuo Terayama

Kanetsu Yoshifusa Aizawa, Akira Imai, Masahiko Kurabayashi, Ban Mihara, Yasuo Sasagawa

Chiba/Saitama Nobuo Araki, Toshio Fukutake, Issei Komuro, Fumitaka Ohsuzu

Tokyo/Kanagawa Masahiko Aosaki, Kazuo Kimura, Norihiro Suzuki, Makoto Takagi, Shigeharu Takagi, Teruo Takano, Teruhisa Tanabe

Tokai Haruo Hirayama, Toyooki Murohara, Takeshi Nakano, Gen Sobue

Kansai Masatsugu Hori, Yukihiko Koretsune

Hokuriku Shunro Endo, Hiroshi Inoue, Kouji Kajinami

Chugoku/Shikoku Naohisa Hamashige, Shotai Kobayash, Masayasu Matsumoto, Masunori Matsuzaki

Kyushu Yoichiro Hashimoto, Hisao Ogawa, Yasushi Okada, Shuichi Okamatsu

Participating Investigators

Hokkaido Takeo Abumiya, Minoru Ajiki, Takeo Baba, Tomoo Furumoto, Shuuzaburo Fukuyama, Masatada Fukuoka, Takayuki Hirabayashi, Tooru Hasegawa, Kiyohiro Houkin, Katsuhisa Ishii, Satoshi Koyama, Kenji Kamiyama, Tetsuro Kohya, Satoshi Kuroda, Tsukasa Kubota, Takaaki Kato, Michifumi Kyuma, Takeo Murahashi, Tomoaki Matsumoto, Jyoji Nakagawara, Daigo Nagahara, Hiroshi Oimatsu, Toshiaki Osato, Hitoshi Ooiwa, Kazuhiro Sako, Motoi Sasaki, Toshihiro Shimizu, Yoshinobu Seo, Tetsuro Shoji, Tsukasa Satoh, Yasukuni Shikano, Mitsunori Shimazaki, Makoto Senoo, Yoshitoki Takagawa, Akihito Tsuchida, Shiho Takagawa, Hideki Takizawa, Shigemichi Tanaka, Hidekazu Takada, Shuichi Yonekura

Tohoku Morio Aihara, Mituaki Hatanaka, Yasuhiro Ishibashi, Hiroyuki Kuki, Takahiko Kikuchi, Yuichi Nozaki, Ayumu Ohnuma, Yukio Onodera, Hiroyuki Oosawa, Satoko Obara, Akifumi Suzuki, Nobuyuki Shiba, Hiroaki Takahashi, Kenichi Tamura, Hajime Yasuda

Kanetsu Kazunori Akaji, Takenori Akiyama, Tsuneo Fujita, Hiroshi Furushima, Mikio Fujimoto, Shintaro Gomi, Kenji Hiraga, Satoru Hirono, Masahiro Hirose, Akira Imai, Masahiro Inoue, Toshiya Iwasaki, Kunihiko Imai, Masaki Jinbo, Yoshiaki Kaneko, Kiminori Kato, Hiroshi Kamiyama, Bun-ichi Kato, Ban Mihara, Masatsugu Morikawa, Mitsunobu Murata, Tatsuru Mihara, Takahide Nagashima, Shibata/Kitakanbara J-TRACE team, Toyoshi Sasaki, Yasuo Sasagawa, Kaoru Suzuki, Keisuke Suzuki, Hiroshi Shimizu, Norisuke Satori, Yutaka Tomita, Yoshio Tanizaki, Toshinori Takahashi, Yasuhiko Yamauchi

Chiba/Saitama Nobuo Araki, Toshio Fukutake, Kouichi Honma, Yoshihiro Iijima, Shoichi Ito, Masatoshi Kusuhara, Yoichi Kuwabara, Yoshio Kobayashi, Kazutaka Matsui, Shinji Matsuda, Takashi Nakazato, Kyoichi Nomura, Toshio Nagai, Fumitaka Ohsuzu, Hideki Shige, Shigeki Tanaka, Kazuo Yamakawa

Tokyo/Kanagawa Hisanao Akiyama, Kuniya Asai, Eiiti Akiyama, Toshiaki Ebina, Mitsuki Endo, Tsutomu Endo, Eri Furukawa, Hirotaka Fujita, Kazuki Fukui, Shinya Fukazawa, Taiji Furukawa, Shinya Goto, Yoshinari Goseki, Haruhiko Hoshino, Kazuhiro Hara, Kei Hatori, Jynya Hosoda, Satoshi Hida, Masaharu Hirano, Kazunori Iwade, Noriaki Iwahashi, Aritomo Inoue, Shinobu Imai, Taizo Ishiyama, Kensuke Ishii, Sugao Ishiwata, Yuji Ikari, Kohei Iguchi, Yutaka Kitamura, Eitaro Kodani, Yasuhisa Kitagawa, Haruhisa Kato, Hirotaka Kato, Takashi Kiyonagi, Ikuyoshi Kusama, Kazuo Kimura, Noriyuki Kawaura, Tsukasa Kobayashi, Takahide Kohro, Kimio Kikushima, Atsumi Kume, Mitsuharu Kawamura, Kazuhiro Muramatsu, Kohei Matsushita, Yoko Morita, Yoshino Minoura, Jun Masuda, Yukiko Morita, Takayuki Mitsuhashi, Nobuhiko Maejima, Minako Murayama, Takamichi Miyamoto, Kagari Matsudaira, Manabu Miyagi, Hiroyoshi Morino, Riichiro Nakayama, Takehiko Nagao, Hiroshi Nishimura, Masashi Nakamaru, Tatsuya Nakachi, Tomoyori Nakatogawa, Takeshi Nakagawa, Masayuki Nakao, Akihiro Niwa, Hirotoshi Ohmura, Hiroyuki Ozaki, Syuuji Ono, Jun Okuda, Yasuo Ohkusu, Yuji Otsuka, Hisako Omori, Mie Shimura, Tomoaki Shimizu, Takahiro Shibata, Ryouma Shibue, Fumio Saito, Shigeru Nogawa, Kenichiro Saka, Toshihiko Saito, Toshiaki Sato, Yutaka Shiina, Naohisa Shindo, Shigeharu Takagi, Makoto Takagi, Kengo Tsukahara, Tatuya Takahashi, Itaru Takamisawa, Tetsu Takamizawa, Naoyuki Takahashi, Shigemasa Tani, Koichi Tamura, Shinichi Takahashi, Hidehito Takase, Takeshi Tadera, Hirokazu Tanaka, Teruhisa Tanabe, Nobuhiro Tanaka, Yasuyoshi Takei, Yoko Takiyama, Shinichiro Uchiyama, Jun Umemura, Hiromi Uno/Terashi, Mikio Usui, Ikuyoshi Watanabe, Masayuki Yotsukura, Hideto Yano, Takeshi Yamakawa, Minako Yoshida

Tokai Tetsuo Ando, Mikio Hirayama, Takashi Hara, Kazuhiro Hara, Nozomi Hishikawa, Mizuki Ito, Noboru Imai, Tetsuya Kitamura, Sukenari Koyabu, Noriko Kodama, Osamu Kawakami, Takashi Kameyama, Takahisa Kondo, Masaaki Kanashiro, Toyooki Murohara, Mamoru Nanasato, Mikiya Nakayabu, Hisayoshi Niwa, Masahiro Oya, Osamu Ohno, Makoto Sugiura, Hidetaka Watanabe

Kansai Ken Araki, Hideki Etani, Ryuzo Fukunaga, Shigetaka Furukado, Hisakazu Fuji, Katsuji Hashimoto, Hiroyuki Hashimoto, Kinji Ishikawa, Kazuo Kitagawa, Yukihiko Koretsune, Akio Kohama, Yoshiyuki Kijima, Shinsuke Nanto, Katsunori Nao, Yoshiyuki Nagai, Keiko Nagano, Osaka Police Hospital's Cardiology, Yutaka Okazaki, Masafumi Tagaya, Tsutomu Takahashi

Hokuriku Hironobu Akao, Shunro Endo, Nakaba Fujioka, Akira Fujiki, Tadakazu Hirai, Jun Harada, Hisanari Ishise, Hiroshi Inoue, Bunji Kaku, Michiya Kubo, Masanori Kyo, Kiyoo Mori, Koichi Mizumaki, Yutaka Nitta, Takashi Nozawa, Makoto Nonomura, Ryoko Sato, Shuaro Takashima, Yoshihiro Takeuchi, Yoshiharu Taguchi, Kortaro Tanaka, Tsukasa Takabatake, Tomio Taguchi

Chugoku/Shikoku Hitoshi Fukuda, Yuhji Furutani, Kaori Fujibe, Takashi Fujii, Naohisa Hamashige, Naohisa Hosomi, Masahiko Harada, Yuji Hisamatsu, Koji Hirashita, Hijiri Ito, Takahiro Iwami, Shoichi Kato, Juri Kitamura, Masateru Kohno, Mitsuhiro Kitani, Hiromi Koide, Masayasu Kimura, Satoshi Kataoka, Kyoko Kobayashi, Toshiro Miura, Shingo Mitaki, Yasuhiro Manabe, Eiichi Nomura, Koichi Noda, Atsushi Nagai, Kiyoshi Nishino, Manabu Nasu, Takanori Namba, Fumiaki Nakao, Shiro Ono, Eiichi Ohnuki, Kazunori Okada, Takayuki Okamura, Tomohiko Ohshita, Hiroaki Sugiura, Kozaburo Seki, Tsuyoshi Torii, Taketo Tanigawa, Hideo Terasawa, Tsuyoshi Ueyama, Masahiro Yamasaki, Kyounori Yasumoto, Toshihiko Yamagata

Kyushu Yutaka Akatsuka, Yoshihiro Fukumoto, Yasuo Fukuda, Takao Fujiki, Yuji Fukutome, Koji Hiyamuta, Rikuzo Hamada, Yasuo Hayashi, Yoichi Hokezu, Yoichiro Hashimoto, Yoshifumi Hirata, Tadashi Hamada, Takeshi Ideguchi, Takuroh Imamura, Masatoshi Koga, Junji Kawagoe, Sunao Kojima, Shigehiko Kumate, Ikuo Misumi, Junnko Mashiba, Takashi Matsuura, Hiroshi Nakane, Toshiyasu Ogata, Hirokuni Ohba, Shuichi Okamatsu, Hideki Oka, Yoshisato Shibata, Satoko Saito, Ichiro Shimada, Kazuhito Tsuruta, Kazuhiro Tashima, Yoshihide Taniwaki, Satoshi Terai, Takeshi Yamada, Hitoshi Yasumoto, Akira Yamada, Tohru Yamawaki