Inherited Cardiomyopathies

Jeffrey A. Towbin, MD

Cardiomyopathies (ie, diseases of the heart muscle) are major causes of morbidity and mortality. A significant percentage of patients with cardiomyopathies have genetic-based, inheritable disease and, over the past 2 decades the genetic causes of these disorders have been increasingly discovered. The genes causing these disorders when they are mutated appear to encode proteins that frame a “final common pathway” for that specific disorder, but the specifics of the phenotype, including age of onset, severity, and outcome is variable for reasons not yet understood. The “final common pathways” for the classified forms of cardiomyopathy include the sarcomere in the primarily diastolic dysfunction disorders hypertrophic cardiomyopathy and restrictive cardiomyopathy, the linkage of the sarcomere and sarcolemma in the systolic dysfunction disorder dilated cardiomyopathy, and the desmosome in arrhythmogenic cardiomyopathy. Left ventricular noncompaction cardiomyopathy (LVNC) is an overlap disorder and it appears that any of these “final common pathways” can be involved depending on the specific form of LVNC. The genetics and mechanisms responsible for these clinical phenotypes will be described. (*Circ J* 2014; **78**: 2347–2356)

Key Words: Cardiomyopathy; Genetics; Heart failure

Cardiomyopathies are major causes of morbidity and mortality and over the past 25 years, limited improvements in outcome have been reported.^{1,2} However, improvements in the understanding of the major forms of cardiomyopathy has occurred over that time, in large part because of advances in imaging, genetics and genomics.^{3,4} A new classification scheme for the cardiomyopathies was recently developed in which 5 forms of disease were formally classified: dilated cardiomyopathy (DCM), hypertrophic cardiomyopathy (HCM), restrictive cardiomyopathy (RCM), arrhythmogenic ventricular cardiomyopathy (AVC), and left ventricular non-compaction cardiomyopathy (LVNC).⁵ These are further classified into genetic/inherited forms and acquired/non-inherited forms. All forms occur at all ages except AVC, which is almost always identified in teenagers and young adults.

Our understanding of the genetic causes of cardiomyopathy has expanded significantly over the past decade. In many cases, the genetic and mechanistic causes of these disorders follow a disturbance in a particular, disease-specific “final common pathway”.⁶ For instance, HCM is now viewed as predominantly an inherited abnormality of contractile protein function (“disease of the sarcomere”). Progress in understanding the genetics of familial DCM has been complicated by its heterogeneous etiologies, but a genetic cause is thought to occur in approximately 50% of subjects, higher in children.^{3–5} The genes identified to date as causative appear to disturb the functional link between the cytoskeleton and sarcomere most commonly.⁴ Multiple genes have been identified for AVC, most resulting in disturbed desmosome/intercalated disk function.^{4,7} Although understanding the genetic basis for the development of RCM and LVNC has been more elusive, genes for both have been identified and

appear to include sarcomere dysfunction as a critical factor.⁸

In this review, the current genetic knowledge of each cardiomyopathy phenotype is covered and I will describe how disruption of these “final common pathways” leads to DCM (sarcolemma-sarcomere link), HCM (sarcomere), RCM (sarcomere), AVC (desmosome), and LVNC (sarcolemma and sarcomere).

Final Common Pathways

Over the past 2 decades, many of the genes responsible for the development of the different cardiomyopathies have been identified. In an initial attempt to target genes for genetic study, my group developed the “final common pathway” hypothesis, which stated that genes encoding proteins with similar functions or involved in the same pathway are responsible for a particular disease or syndrome phenotype.⁶ As a result, we and others identified structure-function similarities of proteins encoded by genes that, when disrupted, led to a somewhat predictable gross clinical phenotype. For instance, the genes identified as causative for inherited arrhythmia disorders tend to encode for ion channels: those for HCM encode for sarcomeric proteins, and AVCs by genes encoding cell-cell junction proteins, as examples (**Figure 1**). In addition, it appears that the protein disturbed by the mutated gene directly disrupts the normal function of the structures in which it is integrated (eg, the sarcomere in HCM when the mutated gene encodes a sarcomeric protein) most commonly but, in some instances, disrupt a binding partner protein, which causes downstream disturbance of the “final common pathway” (eg, a Z-disk protein disrupting the cell-cell junction via a maladaptive binding to desmin,

Received August 11, 2014; accepted August 11, 2014; released online September 2, 2014

The Heart Institute, Cincinnati Children's Hospital Medical Center, Cincinnati, OH, USA

Mailing address: Jeffrey A. Towbin, MD, Professor & Kindervelt-Samuel Kaplan Chair in Pediatric Cardiology and Cardiac Research, Cincinnati Children's Hospital, 3333 Burnet Avenue, Cincinnati, OH 45229, USA. E-mail: jeffrey.towbin@cchmc.org

ISSN-1346-9843 doi:10.1253/circj.CJ-14-0893

All rights are reserved to the Japanese Circulation Society. For permissions, please e-mail: cj@j-circ.or.jp

which negatively interacts with a desmosomal protein causing AVC) (Figure 1).

DCM

DCM is the most common cardiomyopathy, accounting for approximately 55% of cardiomyopathies.⁵ The annual incidence is 2–8/100,000 (0.57/100,000/year in children) with an estimated prevalence of 1/2,500 population.^{5,9} The percentage of cases with a genetic etiology is approximately 30–50% based on the presence of a family history.^{10–12} DCM is characterized by LV dilation and systolic dysfunction (a reduction in myocardial force generation) and is the most common indication

for cardiac transplantation (Figure 2).^{5,9–12} Clinical manifestations include heart failure (HF), thromboembolism, and sudden cardiac death (SCD). The age at onset includes newborn through late adulthood, although most patients are diagnosed between 20–50 years of age.¹²

The genetic forms of DCM usually result from mutated genes encoding 2 major subgroups of proteins: cytoskeletal and sarcomeric proteins.^{3,4,6,13} DCM can also present with muscular involvement and may be the presenting or primary clinical feature of several multisystem conditions, including Emery-Dreifuss muscular dystrophy (EDMD), Barth syndrome, myofibrillar myopathy, limb-girdle muscular dystrophy (LGMD), and Duchenne or Becker muscular dystrophy (DMD/BMD).¹⁴

Genetics of DCM

Inherited, familial DCM (FDCM) has been shown to occur in 30–50% of cases of DCM, with autosomal dominant inheritance being the predominant pattern of transmission; X-linked, autosomal recessive, and mitochondrial inheritance forms are less common.^{10–12,15,16} Over the past 20+ years, substantial progress has been made in understanding of the genetic etiology of FDCM (**Table 1**). Initial progress was made in the early 1990s studying families with X-linked forms of DCM, with the autosomal dominant forms unraveling shortly thereafter.

X-linked DCM, occurring in males in the teen years and early twenties with rapid progression from HF to either death from arrhythmia or transplantation, is distinguished by elevated serum creatine kinase muscle isoforms (CK-MM).^{16,17} Female carriers develop milder, slowly progressive DCM (5th decade). My group in 1993 identified the disease-causing gene as dystrophin and demonstrated significant reduction or absence of dystrophin protein in the heart.¹⁷ These findings were later confirmed by Muntoni et al¹⁸ and others,¹⁹ with mutations generally occurring in the 5' end of the gene. Dystrophin links the sarcomere to the sarcolemma and extracellular matrix (**Figure 3**).¹⁶ When mutated, dystrophin causes DMD/BMD, which are skeletal myopathies that present in males early in life with elevated CK-MM and lead to the early need for wheelchairs (DMD, before age 12 years; BMD, older than 16 years of age),^{20–22} with DCM usually occurring between 10 and 25 years of age.^{16–22}

The most common form of inherited DCM has autosomal dominant inheritance,^{6,13,15} presenting either as classic “pure” DCM or DCM associated with early-onset conduction system disease (CDDC). Genetic heterogeneity exists for autosomal dominant DCM, with more than 40 genes identified for DCM and CDDC,^{3,4,13,23,24} and most of these genes encoding cytoskeletal, sarcomeric, or Z-disk proteins, although mutations in a small number of ion channel-encoding and desmosome-encoding genes have also been identified (**Table 1**). These include genes primarily encoding cytoskeletal (δ -sarcoglycan (SGCD), β -sarcoglycan (SGCB), desmin (DES), lamin A/C (LMNA), vinculin), sarcomeric/myofibrillar (α -cardiac actin (ACTC), troponin T (TNNT2), troponin I (TNNI3), β -myosin heavy chain (MYH7), myosin binding protein C, and α -tropomyosin (TPM1)), and Z-disk proteins (muscle LIM protein (MLP)/cysteine and glycine-rich protein 3 (CSRP3), titin (TTN), telethonin/TCAP, α -actinin-2 (ACTN2), nebulin (NEBL), myopalladin (MYPN), ANKRD1/CARP, and ZASP/LIM-domain binding 3 (LDB3)) (**Table 1, Figure 3**). Ion channel-encoding genes identified to date include the cardiac sodium channel gene SCN5A and calcium homeostasis regulator phospholamban (PLN),^{23,24} and associated desmosome-encoding genes including desmoplakin (DSP), desmoglein-2 (DSG2), and desmocollin-2 (DSC2) have also been shown to result in a DCM phenotype (**Table 1, Figure 3**).²⁵ In the case of CDDC, LMNA is the most common cause, but other causes include DES and SCN5A (**Table 1, Figure 3**).^{26,27} Mechanistically, cytoskeletal proteins are thought to cause defects of force transmission, resulting in the DCM phenotype, whereas defects of force generation have been speculated to be associated with sarcomere protein-induced DCM.^{28,29} The altered desmosomal proteins appear to disrupt the links between the intercalated disk, Z-disk, and sarcomere.^{30,31} Purejav et al recently showed that a gene mutation may disrupt protein binding partners and, depending on this interaction, differential phenotypes and severity may occur.^{32,33}

Mutations in the sarcomere may produce variable forms of cardiomyopathy (**Tables 1–4, Figure 3**). As previously noted,

Table 1. Genetics of Dilated Cardiomyopathy

Gene	Gene name
ABCC9	ATP-binding cassette, subfamily C, member 9
ACTC	α -Cardiac actin
ACTN2	α -Actinin2
ANKRD1	Cardiac ankyrin repeat, domain 1
BAG3	BCL2-associated athanogene 3
CSRP3	Cysteine and glycine-rich protein 3
CTF1	Cardiotrophin 1
DES	Desmin
DYS	Dystrophin
DNAJC19	DnaJ (Hsp40) homolog, subfamily C, member 19
DSC2	Desmocollin 2
DSG2	Desmoglein 2
DSP	Desmoplakin
EMD	Emerin
EYA4	Eyes absent homolog 4
FHL2	Four and a half LIM domains 2
FKTN	Fukutin
FOXD4	Forkhead box D4
LAMA4	α 4-Laminin
LAMP2	Lysosomal-associated membrane protein 3
LDB3	LIM-domain binding 3
LMNA	Lamin A/C
MYBPC3	Myosin binding protein C
MYH6	β -Myosin heavy chain 6
MYH7	β -Myosin heavy chain 7
MYPN	Myopalladin
NEBL	Nebulette
NEXN	Nexilin (F actin-binding protein)
PLN	Phospholamban
PSEN1	Presenilin 1
PSEN2	Presenilin 2
RBM20	RNA binding motif protein 20
SCN5A	Voltage-gated sodium channel, α subunit
SDHA	Succinate dehydrogenase complex, subunit A, flavoprotein
SGCD	δ -Sarcoglycan
SYNE1	Spectrin repeat containing nuclear protein 1
SYNE2	Spectrin repeat containing nuclear protein 2
TAZ	Tafazzin
TCAP	Titin-cap (Telethonin)
TMPO	Thymopoietin
TNNC1	Cardiac troponin C, type 1
TNNI3	Cardiac troponin I, type 3
TNNT2	Cardiac troponin T, type 2
TPM1	α -Tropomyosin 1
TTN	Titin
TTR	Transthyretin
VCL	Vinculin

abnormalities in the force generation or transmission are thought to contribute to the development of DCM.^{28,29,34,35} α -Cardiac actin, a sarcomeric thin filament protein that interacts with tropomyosin and the troponin complex, links the sarcomere to the sarcolemma via its binding to the N-terminus of dystrophin. DCM-causing actin mutations disturb dystrophin-actin binding,^{36,37} causing force transmission abnormalities and DCM.

Figure 3. Cardiac myocyte cytoarchitecture. Schematic of key proteins of the extracellular matrix (ECM), sarcolemma (SL), sarcomere (SM), and nucleus, and some of their protein-protein interactions. Note the SL proteins include ion channels such as SCN5A, L-type calcium channels and others, as well as the dystrophin-associated binding proteins that interact with dystrophin and other cytoplasmic cytoskeletal proteins (dystroglycans, sarcoglycans, syntrophins, dystrobrevin, sarcospan, caveolin), and cadherins that bind with desmosomal proteins (desmocollin, plakophilin, desmoplakin, plakoglobin). The integral SL membrane proteins interact with the ECM via α -dystroglycan-laminin $\alpha 2$ connections. The SM includes thick and thin filament contractile proteins and Z-disk proteins. The amino-terminus of dystrophin binds actin and connects dystrophin with the SM intracellularly, the SL and ECM. The nucleus includes the important membrane proteins, lamin A/C and emerin. The intermediate filament protein, desmin, is another important and prominent linker protein. MLP, muscle LIM protein.

Further, actin interacts in the sarcomere with troponin T and MYH7; mutations in these genes cause DCM or HCM, depending on the position of the mutation³⁷ via development of force generation abnormalities. For MYH7, this probably occurs by perturbing the actin-myosin interaction or altering cross-bridge movement during contraction.^{34–37} TNNT2 mutations disrupt calcium-sensitive troponin C binding.^{34–37} Mutations in PLN have also been identified, which further supports calcium handling as a potentially important mechanism in DCM development.^{34–37} TPM1 mutations alter its surface charge, causing impaired interaction with actin and potentially changing its secondary structure, its binding to actin and position on the thin filament, and alternating actin-myosin interactions and myofilament Ca^{2+} sensitivity.

Another molecular level target for DCM is the Z-disk.^{38–40} Knoll et al identified mutations in MLP/CSRP3, which disturbs its interaction with TCAP³⁹ and its stretch sensor activity. Mutations in CSRP3 cause abnormalities in the T-tubule system and Z-disk architecture as well.⁴¹ Mutations in ACTN2, which binds CSRP3, is involved in crosslinking actin filaments, and shares a common actin-binding domain with dystrophin. In addition, ACTN2 mutations disrupt CSRP3 binding and stretch sensing.^{38–41} Mutations in ZASP/LBD3 cause DCM and LVNC, as well as myofibrillar myopathy.^{38–40,42} This protein, which interacts with ACTN2, the FATZ/calsarcin/myozenin protein family, the myotilin (myotilin, myopalladin, palladin) family, and TCAP that together form a complex with the car-

diac sodium channel, NEBL, and phosphoglucosylase 1, disrupts the actin cytoskeleton when mutated.⁴³ TTN, encoding a giant sarcomeric cytoskeletal protein, contributes to the maintenance of sarcomere organization and myofibrillar elasticity, and interacts with these proteins at the Z-disk/I-band transition zone and causes DCM or AVC when mutated.⁴⁴

Because of significant locus and allelic heterogeneity, genetic testing for DCM has been of limited utility, with pathogenic variants identified in 17–40% of cases using current 40 gene panels, with most genes contributing only a small percentage of pathogenic variants.^{4,13,23,24} Clear genotype-phenotype correlations are rare.

Muscle Is Muscle: Cardiomyopathy and Skeletal Myopathy Genes Overlap

Interestingly, nearly all of the genes identified for inherited DCM are also known to cause skeletal myopathy in humans and/or mouse models: dystrophin mutations cause DMD/BMD; SGCD causes LGMD2F; LMNA causes autosomal dominant EDMD and LGMD1B; ACTC and DES cause nemaline myopathy. In addition, taffazin (TAZ), α -dystrobrevin, LBD3, CSRP3, ACTN2, TTN, and SGCB mutations also cause skeletal myopathy, suggesting that cardiac and skeletal muscle function is interrelated; skeletal muscle fatigue seen in patients with DCM with and without HF may be related to primary skeletal muscle disease and not only cardiac dysfunction. It also sug-

Table 2. Genetics of Hypertrophic Cardiomyopathy

Gene	Gene name
ACTC	α -Cardiac actin
ACTN2	α -Actinin2
ANKRD1	Cardiac ankyrin repeat, domain 1
CAV3	Caveolin 3
COX15	COX 15 homolog, cytochrome C oxidase assembly protein
CRYAB	Crystallin α B
CSRP3	Cysteine and glycine-rich protein 3
GLA	α -Galactosidase
LAMP-2	Lysosomal-associated membrane protein 2
MYBPC3	Myosin binding protein C
MYH6	β -Myosin heavy chain 6
MYH7	β -Myosin heavy chain 7
MYL2	Myosin regulatory light chain 2, slow
MYL3	Myosin light chain 3, slow
MYLK2	Myosin light chain kinase 2
MYO6	Unconventional myosin VI
MYOZ2	Myozenin 2
NEXN	Nexilin (F actin-binding protein)
PLN	Phospholamban
PRKAG2	AMP-activated protein kinase, γ 2, non-catalytic subunit
TNNC1	Cardiac troponin C, type 1
TAZ	Tafazzin
TCAP	Titin-cap (Telethonin)
TNNI3	Cardiac troponin I, type 3
TNNT2	Cardiac troponin T, type 2
TPM1	α -Tropomyosin 1
TTN	Titin
TTR	Transthyretin
VCL	Vinculin

gests that striated muscle dysfunction results from disturbed “final common pathways” and both sets of muscles should be evaluated in DCM.

HCM

HCM is characterized by asymmetric or concentric wall thickening (Figure 2) in the absence of an underlying systemic condition or other cardiac disease.^{5,6,13,16,45} The major effects of this disorder on human health are its predilection to be inherited, its reputation as the most common cause of SCD in young, healthy individuals, and its potential to develop HF because of diastolic factors or development of systolic dysfunction.^{5,46}

Genetics of Familial HCM

HCM is among the most common inherited cardiac disorders, with a prevalence of 1/500 in young adults.⁵ During the past 20+ years, molecular genetic studies have given important insights into the pathogenesis of HCM and provided new perspectives for the diagnosis and management of affected patients. Genetic causes for HCM have been described, with over 20 genes discovered to date, most affecting the sarcomere; however, mutations in genes encoding proteins of the Z-disk or intracellular calcium modulators have also been identified (Table 2). Eight causative genes encode sarcomere proteins, with most (~80%) mutations identified in the MYH7 and

Table 3. Genetics of Restrictive Cardiomyopathy

Gene	Gene name
ACTC1	α -Cardiac actin
BAG3	BCL2-associated athanogene 3
DES	Desmin
GLA	α -Galactosidase
MYH7	β -Myosin heavy chain 7
MYL2	Myosin regulatory light chain 2, slow
MYL3	Myosin light chain 3, slow
MYPN	Myopalladin
TNNI3	Cardiac troponin I, type 3
TNNT2	Cardiac troponin T, type 2
TPM1	α -Tropomyosin 1
TTN	Titin
TTR	Transthyretin

Table 4. Genetics of Left Ventricular Noncompaction Cardiomyopathy

Gene	Gene name
ACTC1	α -Cardiac actin
CASQ2	Cardiac calsequestrin 2
DTNA	α -Dystrobrexin
DYS	Dystrophin
GLA	α -Galactosidase
LDB3	LIM-domain binding 3
LMNA	Lamin A/C
MYBPC3	Myosin binding protein C
MYH7	β -Myosin heavy chain 7
TAZ	Tafazzin
TNNT2	Cardiac troponin T, type 2

MYBPC3 genes (Figure 3).^{4,5,13,45} Mutations identified in sarcomeric genes are typically single nucleotide substitutions and in most instances the mutant protein is thought to incorporate into the sarcomere, exerting a poison peptide effect (dominant negative mutations), with the only exception being MYBPC3, where deletions or insertions leading to a frameshift (ie, a truncated protein with no function), resulting in haploinsufficiency occur.⁴⁷ In addition to defects in the sarcomere-encoding genes, HCM patients have been identified hosting mutations in Z-disk and other nonsarcomere-encoding genes.^{4,13,32,48} The giant protein titin and its interactive Z-disc proteins, including MLP, ZASP, telethonin, nexilin, myopalladin, myozenin-2, α -actinin 2, CARP, and vinculin, have been identified as causes of HCM when the respective gene is mutated⁴⁸ (Table 2, Figure 3). For Z-disk and calcium modulator genes, the specific mechanism has not been clearly elucidated.

Recent clinical guidelines for HCM recommend comprehensive testing for 5 genes (MYBPC3, MYH7, TNNI3, TNNT2, and TPM1), although current Nexgen sequencing approaches have facilitated much broader testing panels to be widely available from commercial laboratories¹⁹ (<http://www.genetests.org> and <http://www.ncbi.nlm.nih.gov/gtr>). Currently, genetic testing for HCM is primarily used to identify families with a detectable genetic cause of disease and to screen at-risk family members. In the clinic at the Heart Institute, mutations are seen in approximately 65–70% of all subjects. The utility of genetic test results in guiding clinical management is limited because of limited genotype-phenotype correlations. Although almost

Figure 4. Septal morphologies in hypertrophic cardiomyopathy (HCM) and the relationship to sarcomeric myofilament-encoding gene mutations. The 4 subgroups of septal morphology in HCM (sigmoidal, reverse curve, apical, neutral) are shown, together with the percentage of HCM cases represented by each morphology and the likelihood of identifying a mutation in a myofilament-encoding gene. Reproduced with permission from Bos JM et al.⁵²

1,000 variants for HCM have been identified to date,^{4,13,32,48,49} most are private and can, therefore, be detected only through comprehensive genetic testing.

Sarcomere variants are identified in up to 60–70% of patients with HCM who also have a family history, and in 40–60% with cases of sporadic HCM.^{13,49} Genotype-phenotype correlations for HCM, however, are incompletely defined, in part because of the private nature of the mutations (in other words, a unique variant). Further, suggestions that a specific mutation in a specific sarcomeric gene will lead to a specific phenotype and specific outcome have been questioned. The pattern and extent of left ventricular (LV) hypertrophy in patients with HCM, as well as outcomes, is heterogeneous even in first-degree relatives.⁵⁰ This heterogeneity has made it difficult to correlate genotype and phenotype, including outcomes. A systematic review and meta-analysis of genotype-phenotype correlations in 2,459 patients and 53 families with sarcomeric gene mutations reported in 18 publications showed that the presence of any sarcomere gene mutation was associated with younger age at presentation (38.4 vs. 46.0 years), family history of HCM (50.6% vs. 23.1%), family history of SCD (27.0% vs. 14.9%), and greater LV wall thickness (21.0 vs. 19.3 mm).⁵¹ The relationship between cardiac morphology, mutated gene, and outcome has also been evaluated.^{50,52–54} Bos et al studied 359 patients, with 34% being genotype-positive for a putative HCM-associated mutation in 1 or more genes. Echocardiographic reverse curve morphological subtype (**Figure 4**), age at diagnosis younger than 45 years, maximum LV wall thickness ≥ 20 mm, family history of HCM, and family history of SCD were identified as positive predictors of mutation identification; hypertension was a negative predictor.^{52,53} A score, based on the number of predictors of a positive genetic test result, predicted mutation identification in 6%, when only hypertension was present, to 80% when all 5 positive predictor markers were present.

RCM

RCM is the rarest form of cardiomyopathy, characterized by

dilated atria, increased ventricular stiffness with normal ventricular wall thickness and systolic function (**Figure 2**).^{5,8} SCD or a life-threatening cardiac event because of ventricular arrhythmias or heart block is common, as is HF; death occurs within a few years of diagnosis.^{5,8} Some reports suggest clinical overlap between RCM and HCM.^{55,56}

Genetics of RCM

RCM usually has autosomal dominant inheritance, but autosomal recessive, X-linked and mitochondrial-transmitted disease occurs. Most of the identified genes encode sarcomere or Z-disk proteins, such as TNNI3, TNNT2, MYH7, ACTC1, TPM1, MYL3, and MYL28.⁵⁷ (Table 3). Z-disk protein-encoding genes, including MYPN, TTN, and Bag3, have also been identified.^{4,5,8,13,33} Missense variants in DES have been identified in several families with desmin-related myopathy, which can present with RCM, with or without skeletal myopathy and/or atrioventricular block.⁸ Transthyretin (TTR) is associated with amyloid-related RCM.⁸

LV Noncompaction

LVNC is a heterogeneous myocardial disorder characterized by prominent trabeculae that are most evident in the LV apex, intratrabecular recesses, and LV myocardium with 2 distinct layers: compacted and noncompacted myocardium (**Figure 2**).^{5,13,16,58} Although LVNC primarily affects the LV, isolated RV and biventricular noncompaction also occurs.⁵⁸ Multiple forms of LVNC occur and include primary myocardial forms, a form associated with arrhythmias, and LVNC associated with congenital heart disease (CHD), including septal defects (atrial and ventricular), hypoplastic left heart syndrome (HLHS), and right heart obstructive abnormalities including pulmonic stenosis (PS) and Ebstein's anomaly (**Figure 5**), among others.⁵⁸ Outcomes appear to differ based on the specifics of the phenotype.

Genetics of LVNC

LVNC most commonly has X-linked recessive or autosomal

Figure 5. Echocardiographic features of the left ventricular noncompaction (LVNC) phenotype's heterogeneous clinical characteristics. (A) 4-chamber view demonstrates normal atrial and ventricular sizes and thicknesses, with a hypertrabeculated LV wall and apex (arrow). (B) 4-chamber view demonstrates normal atrial sizes with a dilated and hypertrabeculated LV wall and apex (arrow). (C) 4-chamber view demonstrates hypertrophic and hypertrabeculated ventricular walls and apex with a hypertrophic septum (arrow). (D) 4-chamber view demonstrates normal to small ventricular sizes with dilated atria bilaterally (arrows). The LV is hypertrabeculated. (E) 4-chamber view demonstrates normal atrial sizes with severely hypertrabeculated left and right ventricular walls and apices (arrows). (F) 4-chamber view demonstrates normal atrial sizes with tricuspid atresia (no tricuspid valve) and a hypertrabeculated LV wall and apex (arrow). (G) 4-chamber view of Ebstein's anomaly with LVNC, which demonstrates a normal left atrial size with a large right atrium, displaced tricuspid valve towards the right ventricle (closed arrowhead), and a severely hypertrabeculated LV wall and apex (open arrowhead).

dominant inheritance, with reports suggesting 70% having autosomal dominant and 30% X-linked inheritance, at least in childhood.^{58–60} Autosomal recessive and mitochondrial inheritance also occurs.⁵⁸ When LVNC is associated with CHD, the congenital cardiac defect may be heterogeneous in families, but this form of LVNC is transmitted as an autosomal dominant trait along with the CHD; affected members may have no CHD at initial evaluation because the cardiac defects include “minor” forms of CHD (small VSDs, ASDs, patent ductus arteriosus), which have spontaneously normalized or penetrance may be reduced, while others may have severe forms of CHD (HLHS, Ebstein's, etc).^{58,61} Ichida et al reported that 44% of her patients had inherited LVNC, with 70% having autosomal dominant and 30% X-linked inheritance.⁵⁹ In addition, patients with

chromosomal abnormalities are associated with LVNC.

The first genetic cause of isolated LVNC was initially described by Bleyl et al, when they identified mutations in the X-linked TAZ gene, the gene also responsible for Barth syndrome, in patients and carrier females (Table 3).⁶² Multiple genes causing autosomal dominant LVNC have now been identified, including mutations in genes causing CHD with LVNC. In patients with HLHS and LVNC, the cytoskeletal gene α -dystrobrevin has been identified, while mutations in Nkx-2.5 in children with LVNC and ASD and MYH7 in patients with LVNC and Ebstein's anomaly have also been reported (Table 3).^{58,63–66} In LVNC without CHD, mutations in the Z-line protein-encoding ZASP/LDB3 gene and the sarcomere-encoding genes (MYH7, ACTC, TNNT2, MYBPC3, TMP1,

Table 5. Genetics of Arrhythmogenic Ventricular Cardiomyopathy

Gene	Gene name
CTNNA3	α T-catenin
DES	Desmin
DNAJC19	DnaJ (Hsp40) homolog, subfamily C, member 19
DSC2	Desmocollin 2
DSG2	Desmoglin 2
DSP	Desmoplakin
JUP	Junctional plakoglobin
LMNA	Lamin A/C
PKP2	Plakophilin 2
PKP4	Plakophilin 4
RYR2	Ryanodine receptor 2
STRN	Striatin
TGF β 3	Transforming growth factor- β 3
TMEM43	Transmembrane 43
TTN	Titin

and TNIN3) appear to account for 20% or more of cases.^{58,67,68} Hoedemaekers et al additionally demonstrated an association of LVNC with genetic variants in 2 calcium handling genes, as well as TAZ and LMNA,⁶⁸ while Probst et al further showed that sarcomere gene mutations are important in LVNC, showing a prevalence of 29%, with MYH7 and MYBPC3 most frequently mutated (13% and 8%, respectively).⁶⁷ Dellefave et al also identified sarcomere mutations in LVNC, including patients presenting with HF in infancy with and without compound heterozygosity.⁶⁹ In addition to sarcomere-encoding genes and the cytoskeleton, mutations in the sodium channel gene, SCN5A, is associated with LVNC and rhythm disturbance.⁶³ Diagnostic testing in patients with LVNC appears to have a detection rate of clinically significant variants in 35–40% of individuals, with sarcomere-encoding genes most commonly found to be mutated.^{4,70}

Arrhythmogenic Cardiomyopathies

AVC, previously referred to as arrhythmogenic right ventricular dysplasia or cardiomyopathy and recently re-named in a consensus statement,⁵ constitutes a hereditary cardiomyopathy usually with an autosomal dominant inheritance pattern, although a recessive cardiocutaneous disorder also occurs.⁷¹ The RV was initially thought to solely be affected in AVC, and therefore any associated disease of the LV was considered exclusionary for the diagnosis in the original “Task Force criteria” for diagnosis but, more recently, it has been recognized that the LV can also be involved and actually is often associated with severe disease and a worse prognosis.⁷² Classically, however, the RV is dilated with fibrofatty infiltration with no or only minimal LV involvement.^{5,7} Clinically, progressive disease is characterized by systolic impairment and biventricular dilation (with or without a ventricular aneurysm) and clinical features of HF (Figure 2). In most parts of the world, phenotypic expression is more common in men than in women (2–3:1).⁷³ AVC commonly manifests during late childhood or adolescence, but can also emerge in the elderly, and has a prevalence of 1/2,000, which can be higher in certain geographical regions such as the Veneto region or the Greek island Naxos.⁷⁴ AVC is recognized as a leading cause of SCD in young adults ≤ 35 years of age and may account for up to 10% of cardiovas-

cular deaths in the <65 age group.⁵

Genetics of AVC

Analyses of first- and second-degree relatives of patients with AVC suggest that up to 50% of AVC cases are familial.^{5,7,75} AVC is most commonly inherited as an autosomal dominant trait with incomplete penetrance,^{5,7,75} although 2 autosomal recessive forms have been described.⁷¹ To date, 15 genes have been reported to cause AVC (Table 5).^{4,5,7,13,76} My group and others demonstrated that compound and digenic heterozygosity is involved in AVC pathogenesis in up to 20% of cases and leads to more severe disease.^{77,78} As penetrance is incomplete, genetically affected relatives often demonstrate variable and mild (or no) phenotype and the prevalence of familial disease is often underestimated in clinical practice.^{13,73}

The first gene to be identified in AVC was junctional plakoglobin (JUP) when a homozygous 2-nucleotide deletion in JUP was reported in patients with Naxos disease.^{7,71,73} This gene encodes the plakoglobin protein, a key protein of the desmosome in the intercalated disk. In autosomal dominant AVC, the desmosomal protein-encoding DSP gene was identified as disease-causing.^{7,71,73}

Three groups of desmosomal proteins are known: (1) transmembrane desmosomal cadherins, including DSC2 and DSG2; (2) DSP, a plakophilin family protein that attaches directly to intermediate filaments (desmin in the myocardium); and (3) linker proteins such as armadillo family proteins, including JUP and plakophilin 2 (PKP2) that mediate interactions between the desmosomal cadherin tails and DSP.⁷¹ In approximately 80% of patients with confirmed pathogenic mutations, PKP2, DSP and DSG2 are altered.⁷⁴ Besides desmosomal gene mutations, mutations in genes encoding proteins that interact with desmosomal proteins were found as well, including: (1) the transforming growth factor β 3 that conveys cytokine-stimulating fibrosis and modulates cell adhesion and growth; (2) the transmembrane protein 43 (TMEM43), which functions as a PPAR- γ response element, an adipogenic transcription factor; (3) DES, which binds desmoplakin; and (4) TTN, which bridges the sarcomere along its longitudinal axis and forms a continuous filament along the myofibril (Table 5).^{4,7,13,73,79,80}

Genetically determined disruption of the integrity of the intercalated disk is a key factor promoting the development of AVC and SCD. Recent data indicates that loss of desmosomal integrity can substantially affect gap junctions, sodium channel function and electrical propagation, thereby promoting ventricular arrhythmias in the absence of overt structural damage^{81,82} and thus providing an overlapping phenotype (cardiomyopathy plus arrhythmias) because of disruption of 2 “final common pathways” (desmosome and ion channel).^{80,83}

Conclusions

The cardiomyopathies appear to occur caused by disruption of “final common pathways”. These disruptions may be purely genetic causes, such as mutations in a single gene that results in a dysfunctional protein and leads to a domino effect of downstream protein interaction abnormalities and ultimately a phenotype. In other situations, multiple mutations in the same gene (compound heterozygosity) or in different genes (digenic heterozygosity) may lead to a phenotype that may be classic, more severe, or even overlap with other disease forms. In some cases, different intersecting pathways may become disturbed, resulting in complex phenotypes. Further, acquired causes may play a role by causing disruption of these functional pathways. For instance, DCM results from disruption of the “final common

pathway” linking the sarcomere and sarcolemma, and mutations in the affected genes are responsible for cardiac and skeletal muscle dysfunction. The mechanisms of disease, which include disruption of the linkage because of protein-protein interaction abnormalities that occur with dysfunctional proteins, as well as the interplay of other factors such as mechanical stress and stretch, are being elucidated in detail with the development of animal models of the human disease. Many of the genes identified are now clinically available in fee-for-service laboratories. Novel therapies have resulted from the improved understanding of this clinical phenotype, as noted. Similarly, the genetic basis and mechanistic understanding of HCM as a disturbance of sarcomere function has occurred over the past 2 decades and genetic tests are clinically available. The development of novel targeted therapies has been somewhat slow in coming but is expected to develop in the near future. In the case of LVNC, the “new kid on the block”, genetic understanding is in the early phase. The future of cardiomyopathy care is poised to shift in the next decade because of these new developments, as well as the growing science of stem cell therapy. Because children have “pure” disease states, unfettered by comorbidities, the dream of “cures” of muscle disease (cardiac, skeletal muscle) will likely be realized more fully in this population. We must move towards that goal.

References

- Lee CS, Chien CV, Bidwell JT, Gelow JM, Denfeld QE, Creber RM, et al. Comorbidity profiles and inpatient outcomes during hospitalization for heart failure: An analysis of the U.S. Nationwide inpatient sample. *BMC Cardiovasc Disord* 2014; **14**: 73.
- Shamszad P, Hall M, Rossano JW, Denfield SW, Knudson JD, Penny DJ, et al. Characteristics and outcomes of heart failure-related intensive care unit admissions in children with cardiomyopathy. *J Card Fail* 2013; **19**: 672–677.
- Cahill TJ, Ashrafian H, Watkins H. Genetic cardiomyopathies causing heart failure. *Circ Res* 2013; **113**: 660–675.
- Teekakirikul P, Kelly MA, Rehm HL, Lakdawala NK, Funke BH. Inherited cardiomyopathies: Molecular genetics and clinical genetic testing in the postgenomic era. *J Mol Diagn* 2013; **15**: 158–167.
- Maron BJ, Towbin JA, Thiene G, Antzevitch C, Corrado D, Arnett D, et al. Contemporary definitions and classification of the cardiomyopathies: An American Heart Association Scientific Statement from the Council on Clinical Cardiology, Heart Failure and Transplantation Committee; Quality of Care and Outcomes Research and Functional Genomics and Translational Biology Interdisciplinary Working Groups; and Council on Epidemiology and Prevention. *Circulation* 2006; **113**: 1807–1816.
- Bowles NE, Bowles KR, Towbin JA. The “final common pathway” hypothesis and inherited cardiovascular disease: The role of cytoskeletal proteins in dilated cardiomyopathy. *Herz* 2000; **25**: 168–175.
- Ellinor PT, MacRae CA, Thierfelder L. Arrhythmogenic right ventricular cardiomyopathy. *Heart Fail Clin* 2010; **6**: 161–177.
- Sen-Chowdhry S, Syrris P, McKenna WJ. Genetics of restrictive cardiomyopathy. *Heart Fail Clin* 2010; **6**: 179–186.
- Towbin JA, Lowe AM, Colan SD, Sleeper LA, Orav EJ, Clunie S, et al. Incidence, causes and outcomes of dilated cardiomyopathy in children. *JAMA* 2006; **296**: 1867–1876.
- Fatkin D; members of the CSANZ Cardiac Genetic Diseases Council Writing Group. Guidelines for the diagnosis and management of familial dilated cardiomyopathy. *Heart Lung Circ* 2011; **20**: 691–693.
- Givertz MM, Mann DL. Epidemiology and natural history of recovery of left ventricular function in recent onset dilated cardiomyopathies. *Curr Heart Fail Rep* 2013; **10**: 321–330.
- Dec GW, Fuster V. Idiopathic dilated cardiomyopathy. *N Engl J Med* 1994; **331**: 1564–1575.
- Ackerman MJ, Priori SG, Willems S, Berul C, Brugada R, Calkins H, et al. HRS/EHRA expert consensus statement on the state of genetic testing for the channelopathies and cardiomyopathies: This document was developed as a partnership between the Heart Rhythm Society (HRS) and the European Heart Rhythm Association (EHRA). *Heart Rhythm* 2011; **8**: 1308–1339.
- Finsterer J, Stöllberger C, Wahbi K. Cardiomyopathy in neurological disorders. *Cardiovasc Pathol* 2013; **22**: 389–400.
- Hershberger RE, Lindenfeld J, Mestroni L, Seidman CE, Taylor MR, Towbin JA; Heart Failure Society of America. Genetic evaluation of cardiomyopathy: A Heart Failure Society of America practice guideline. *J Card Fail* 2009; **15**: 83–97.
- Towbin JA, Bowles NE. The failing heart. *Nature* 2002; **415**: 227–233.
- Towbin JA, Hejtmancik JF, Brink P, Gelb B, Zhu XM, Chamberlain JS, et al. X-linked dilated cardiomyopathy (XLCM): Molecular genetic evidence of linkage to the Duchenne muscular dystrophy gene at the Xp21 locus. *Circulation* 1993; **87**: 1854–1865.
- Muntoni F, Cau M, Ganau A, Congiu R, Arvedi G, Mateddu A, et al. Brief report: Deletion of the dystrophin muscle-specific promoter region associated with X-linked dilated cardiomyopathy. *N Engl J Med* 1993; **329**: 921–925.
- Diegoli M, Grasso M, Favalli V, Serio A, Gambarin FI, Klersy C, et al. Diagnostic work-up and risk stratification in X-linked dilated cardiomyopathies caused by dystrophin defects. *J Am Coll Cardiol* 2011; **58**: 925–934.
- Rahimov F, Kunkel LM. The cell biology of disease: Cellular and molecular mechanisms underlying muscular dystrophy. *J Cell Biol* 2013; **201**: 499–510.
- Constantin B. Dystrophin complex functions as a scaffold for signaling proteins. *Biochim Biophys Acta* 2014; **1838**: 635–642.
- Chelly J, Desguerre I. Progressive muscular dystrophies. *Handb Clin Neurol* 2013; **113**: 1343–1366.
- Morales A, Hershberger RE. Genetic evaluation of dilated cardiomyopathy. *Curr Cardiol Rep* 2013; **15**: 375, doi:10.1007/s11886-013-0375-1.
- Mestroni L, Taylor MR. Genetics and genetic testing of dilated cardiomyopathy: A new perspective. *Discov Med* 2013; **15**: 43–49.
- Garcia-Pavia P, Syrris P, Salas C, Evans A, Mirelis JG, Cobo-Marcos M, et al. Desmosomal protein gene mutations in patients with idiopathic dilated cardiomyopathy undergoing cardiac transplantation: A clinicopathological study. *Heart* 2011; **97**: 1744–1752.
- Wolf CM, Wang L, Alcalai R, Pizad A, Burgon PG, Ahmad F, et al. Lamin A/C haploinsufficiency causes dilated cardiomyopathy and apoptosis-triggered cardiac conduction system disease. *J Mol Cell Cardiol* 2008; **44**: 293–303.
- Taylor MR, Slavov D, Ku L, Di Lenarda A, Sinagra G, Carniel E, et al. Prevalence of desmin mutations in dilated cardiomyopathy. *Circulation* 2007; **115**: 1244–1251.
- Debold EP, Schmitt JP, Patlak JB, Beck SE, Moore JR, Seidman JG, et al. Hypertrophic and dilated cardiomyopathy mutations differentially affect the molecular force generation of mouse alpha-cardiac myosin in the laser trap assay. *Am J Physiol Heart Circ Physiol* 2007; **293**: H284–H291.
- Ehler E, Perriard JC. Cardiomyocyte cytoskeleton and myofibrillogenesis in healthy and diseased heart. *Heart Fail Rev* 2000; **5**: 259–269.
- Basso C, Czarnowska E, Della Barbera M, Bauce B, Beffagna G, Wlodarska EK, et al. Ultrastructural evidence of intercalated disc remodelling in arrhythmogenic right ventricular cardiomyopathy: An electron microscopy investigation on endomyocardial biopsies. *Eur Heart J* 2006; **27**: 1847–1854.
- Yang Z, Bowles NE, Scherer SE, Taylor MD, Kearney DL, Ge S, et al. Desmosomal dysfunction due to mutations in desmoplakin causes arrhythmogenic right ventricular dysplasia/cardiomyopathy. *Circ Res* 2006; **99**: 646–655.
- Purevjav E, Arimura T, Augustin S, Huby AC, Takagi K, Nunoda S, et al. Molecular basis for clinical heterogeneity in inherited cardiomyopathies due to myopalladin mutations. *Hum Mol Genet* 2012; **21**: 2039–2053.
- Purevjav E, Varela J, Morgado M, Kearney DL, Li H, Taylor MD, et al. Nebulette mutations are associated with dilated cardiomyopathy and endocardial fibroelastosis. *J Am Coll Cardiol* 2010; **56**: 1493–1502.
- Fatkin D, Graham RM. Molecular mechanisms of inherited cardiomyopathies. *Physiol Ref* 2002; **82**: 945–980.
- McNally EM, Golbus JR, Puckelwartz MJ. Genetic mutations and mechanisms in dilated cardiomyopathy. *J Clin Invest* 2013; **123**: 19–26.
- Olson TM, Michels VV, Thibodeau SN, Tai YS, Keating MT. Actin mutations in dilated cardiomyopathy, a heritable form of heart failure. *Science* 1998; **280**: 750–752.
- Kamisago M, Sharma SD, DePalma SR, Solomon S, Sharma P, McDonough B, et al. Mutations in sarcomere protein genes as a cause of dilated cardiomyopathy. *N Engl J Med* 2000; **343**: 1688–1696.
- Pyle WG, Solaro RJ. At the crossroads of myocardial signaling: The role of Z-discs in intracellular signaling and cardiac function. *Circ Res* 2004; **94**: 296–305.

39. Knoll R, Hoshijima M, Hoffman HM, Person V, Lorenzen-Schmidt I, Bang ML, et al. The cardiac mechanical stretch sensor machinery involves a Z disc complex that is defective in a subset of human dilated cardiomyopathy. *Cell* 2002; **111**: 943–955.
40. Vatta M, Sinagra G, Brunelli L, Faulkner G. Remodeling of dystrophin and sarcomeric Z-band occurs in pediatric cardiomyopathies: A unifying mechanism for force transmission defect. *J Cardiovasc Med (Hagerstown)* 2009; **10**: 149–156.
41. Mohapatra B, Jimenez S, Lin JH, Bowles KR, Coveler KJ, Marx JG, et al. Mutations in the muscle LIM protein and α -actinin-2 genes in dilated cardiomyopathy and endocardial fibroelastosis. *Mol Genet Metab* 2003; **80**: 207–215.
42. Lin X, Ruiz J, Bajraktari I, Ohman R, Banerjee S, Gribble K, et al. Z-disc-associated, alternatively spliced, PDZ motif-containing protein (ZASP) mutations in the actin-binding domain cause disruption of skeletal muscle actin filaments in myofibrillar myopathy. *J Biol Chem* 2014; **289**: 13615–13626.
43. Martinelli VC, Kyle WB, Kojic S, Vitulo N, Li Z, Belgrano A, et al. ZASP interacts with the mechanosensing protein Ankrd2 and p53 in the signalling network of striated muscle. *PLoS One* 2014; **9**: e92259, doi:10.1371/journal.pone.0092259.
44. LeWinter MM, Granzier HL. Cardiac titin and heart disease. *J Cardiovasc Pharmacol* 2014; **63**: 207–212.
45. Maron BJ, Maron MS. Hypertrophic cardiomyopathy. *Lancet* 2013; **381**: 242–255.
46. O'Mahony C, Elliott PM. Prevention of sudden cardiac death in hypertrophic cardiomyopathy. *Heart* 2014; **100**: 254–260.
47. Marston S, Copeland O, Jacques A, Livesey K, Tsang V, McKenna WJ, et al. Evidence from human myectomy samples that MYBPC3 mutations cause hypertrophic cardiomyopathy through haploinsufficiency. *Circ Res* 2009; **105**: 219–222.
48. Lopes LR, Zekavati A, Syrris P, Hubank M, Giambartolomei C, Dalageorgou C, et al. Genetic complexity in hypertrophic cardiomyopathy revealed by high-throughput sequencing. *J Med Genet* 2013; **50**: 228–239.
49. Ho CY. Genetics and clinical destiny: Improving care in hypertrophic cardiomyopathy. *Circulation* 2010; **122**: 2430–2440.
50. Bos JM, Will ML, Gersh BJ, Kruisselbrink TM, Ommen SR, Ackerman MJ. Characterization of a phenotype-based genetic test prediction score for unrelated patients with hypertrophic cardiomyopathy. *Mayo Clin Proc* 2014; **89**: 727–737.
51. Lopes LR, Rahman MS, Elliott PM. A systematic review and meta-analysis of genotype-phenotype associations in patients with hypertrophic cardiomyopathy caused by sarcomeric protein mutations. *Heart* 2013; **99**: 1800–1811.
52. Bos JM, Towbin JA, Ackerman MJ. Diagnostic, prognostic, and therapeutic implications of genetic testing for hypertrophic cardiomyopathy. *J Am Coll Cardiol* 2009; **54**: 201–211.
53. Theis JL, Bos JM, Bartleson VB, Will ML, Binder J, Vatta M, et al. Echocardiographic-determined septal morphology in Z-disc hypertrophic cardiomyopathy. *Biochem Biophys Res Commun* 2006; **351**: 896–902.
54. Geske JB, Bos JM, Gersh BJ, Eidem BW, Ackerman MJ. Deformation patterns in genotyped patients with hypertrophic cardiomyopathy. *Eur Heart J Cardiovasc Imaging* 2014; **15**: 456–465.
55. Rivenes SM, Kearney DL, Smith EO, Towbin JA, Denfield SW. Sudden death and cardiovascular collapse in children with restrictive cardiomyopathy. *Circulation* 2000; **102**: 876–882.
56. Walsh MA, Grenier MA, Jefferies JL, Towbin JA, Lorts A, Czosek RJ. Conduction abnormalities in pediatric patients with restrictive cardiomyopathy. *Circ Heart Fail* 2012; **5**: 267–273.
57. Caleshu C, Sakhuja R, Nussbaum RL, Schiller NB, Ursell PC, Eng C, et al. Furthering the link between the sarcomere and primary cardiomyopathies: Restrictive cardiomyopathy associated with multiple mutations in genes previously associated with hypertrophic or dilated cardiomyopathy. *Am J Med Genet A* 2011; **155A**: 2229–2235.
58. Towbin JA. Left ventricular noncompaction: A new form of heart failure. *Heart Fail Clin* 2010; **6**: 453–469.
59. Ichida F, Hamamichi Y, Miyawaki T, Ono Y, Kamiya T, Akagi T, et al. Clinical features of isolated noncompaction of the ventricular myocardium: Long-term clinical course, hemodynamic properties, and genetic background. *J Am Coll Cardiol* 1999; **34**: 233–240.
60. Sasse-Klaassen S, Gerull B, Oechslin E, Jenni R, Thierfelder L. Isolated noncompaction of the left ventricular myocardium in the adult is an autosomal dominant disorder in the majority of patients. *Am J Med Genet A* 2003; **119A**: 162–167.
61. Stahli BE, Gebhard C, Biaggi P, Klaassen S, Valsangiacomo Buechel E, Attenhofer Jost CH, et al. Left ventricular non-compaction: Prevalence in congenital heart disease. *Int J Cardiol* 2013; **167**: 2477–2481.
62. Bleyl SB, Mumford BR, Brown-Harrison MC, Pagotto LT, Carey JC, Pyscher TJ, et al. Xq28-linked noncompaction of the left ventricular myocardium: Prenatal diagnosis and pathologic analysis of affected individuals. *Am J Med Genet* 1997; **72**: 257–265.
63. Udeoji DU, Philip KJ, Morrissey RP, Phan A, Schwarz ER. Left ventricular noncompaction cardiomyopathy: Updated review. *Ther Adv Cardiovasc Dis* 2013; **7**: 260–273.
64. Ouyang P, Saarel E, Bai Y, Luo C, Lv Q, Xu Y, et al. A de novo mutation in NKX2.5 associated with atrial septal defects, ventricular noncompaction, syncope and sudden death. *Clin Chim Acta* 2011; **412**: 170–175.
65. Postma AV, van Engelen K, van de Meerakker J, Rahman T, Probst S, Baars MJ, et al. Mutations in the sarcomere gene MYH7 in Ebstein anomaly. *Circ Cardiovasc Genet* 2011; **4**: 43–50.
66. Vermeer AM, van Engelen K, Postma AV, Baars MJ, Christiaans I, De Haaij S, et al. Ebstein anomaly associated with left ventricular noncompaction: An autosomal dominant condition that can be caused by mutations in MYH7. *Am J Med Genet C Semin Med Genet* 2013; **163C**: 178–184.
67. Probst S, Oechslin E, Schuler P, Greutmann M, Boyé P, Knirsch W, et al. Sarcomere gene mutations in isolated left ventricular noncompaction cardiomyopathy do not predict clinical phenotype. *Circ Cardiovasc Genet* 2011; **4**: 367–374.
68. Hoedemaekers YM, Caliskan K, Michels M, Frohn-Mulder I, van der Smagt JJ, Pfeifferkorn JE, et al. The importance of genetic counseling, DNA diagnostics, and cardiologic family screening in left ventricular noncompaction cardiomyopathy. *Circ Cardiovasc Genet* 2010; **3**: 232–239.
69. Dellefave LM, Pytel P, Mewborn S, Mora B, Guris DL, Fedson S, et al. Sarcomere mutations in cardiomyopathy with left ventricular hypertrabeculation. *Circ Cardiovasc Genet* 2009; **2**: 442–449.
70. Shieh JT. Implications of genetic testing in noncompaction/hypertrabeculation. *Am J Med Genet C Semin Med Genet* 2013; **163C**: 206–211.
71. Protonotarios N, Tsatsopoulou A. Naxos disease and Carvajal syndrome: Cardiac cutaneous disorders that highlight the pathogenesis and broaden the spectrum of arrhythmogenic right ventricular cardiomyopathy. *Cardiovasc Pathol* 2004; **13**: 185–194.
72. Marcus FI, McKenna WJ, Sherrill D, Basso C, Bauce B, Bluemke DA, et al. Diagnosis of arrhythmogenic right ventricular cardiomyopathy/dysplasia: Proposed modification of the task force criteria. *Circulation* 2010; **121**: 1533–1541.
73. Azaouagh A, Churzidse S, Konorza T, Erbel R. Arrhythmogenic right ventricular cardiomyopathy/dysplasia: A review and update. *Clin Res Cardiol* 2011; **100**: 383–394.
74. Herren T, Gerber PA, Duru F. Arrhythmogenic right ventricular cardiomyopathy/dysplasia: A not so rare “disease of the desmosome” with multiple clinical presentations. *Clin Res Cardiol* 2009; **98**: 141–158.
75. Hamid MS, Norman M, Quraishi A, Firoozi S, Thaman R, Gimeno JR, et al. Prospective evaluation of relatives for familial arrhythmogenic right ventricular cardiomyopathy/dysplasia reveals a need to broaden diagnostic criteria. *J Am Coll Cardiol* 2002; **40**: 1445–1450.
76. Te Rijdt WP, Jongbloed JD, de Boer RA, Thiene G, Basso C, van den Berg MP, et al. Clinical utility gene card for: Arrhythmogenic right ventricular cardiomyopathy (ARVC). *Eur J Hum Genet* 2014; **22**, doi:10.1038/ejhg.2013.124.
77. Xu T, Yang Z, Vatta M, Rampazzo A, Beffagna G, Pilichou K, et al. Compound and digenic heterozygosity contributes to arrhythmogenic right ventricular cardiomyopathy. *J Am Coll Cardiol* 2010; **55**: 587–597.
78. Bauce B, Nava A, Beffagna G, Basso C, Lorenzon A, Smaniotta G, et al. Multiple mutations in desmosomal proteins encoding genes in arrhythmogenic right ventricular cardiomyopathy/dysplasia. *Heart Rhythm* 2010; **7**: 22–29.
79. Patel DM, Green KJ. Desmosomes in the heart: A review of clinical and mechanistic analyses. *Cell Commun Adhes* 2014; **21**: 109–128.
80. Vatta M, Marcus F, Towbin JA. Arrhythmogenic right ventricular cardiomyopathy: A ‘final common pathway’ that defines clinical phenotype. *Eur Heart J* 2007; **28**: 529–530.
81. Delmar M, McKenna WJ. The cardiac desmosome and arrhythmogenic cardiomyopathies: From gene to disease. *Circ Res* 2010; **107**: 700–714.
82. Noorman M, Hakim S, Kessler E, Groeneweg JA, Cox MG, Asimaki A, et al. Remodeling of the cardiac sodium channel, connexin43, and plakoglobin at the intercalated disk in patients with arrhythmogenic cardiomyopathy. *Heart Rhythm* 2013; **10**: 412–419.
83. Towbin JA, Lorts A. Arrhythmias and dilated cardiomyopathy common pathogenetic pathways? *J Am Coll Cardiol* 2011; **57**: 2169–2171.