3. Tanaka A, Koga S, Hiramatsu Y. Donepezil-induced adverse side effects of cardiac rhythm: 2 cases report of atrioventricular block and Torsade de pointes. *Intern Med* 2009; **48**: 1219–1223.
4. Takaya T, Okamoto M, Yodoi K, Hata K, Kijima Y, Nakajima H, et al. Torsades de pointes with QT prolongation related to donepezil use. *J Cardiol* 2009; **54**: 507–511.
5. Hadano Y, Ogawa H, Wakeyama T, Iwami T, Kimura M, Mochizuki M, et al. Donepezil-induced Torsades de pointes without QT prolongation. *J Cardiol Cases* 2013; **8**: e69–e71.
6. Kitt J, Irons R, Al-Obaidi M, Missouri C. A case of donepezil-related Torsades de pointes. *BMJ Case Rep* 2015; **2015**: bcr2015211900.
7. Gurbuz AS, Ozturk S, Acar E, Efe SC, Akgun T, Kilicgedik A, et al. Acquired long QT syndrome and Torsades de pointes related to donepezil use in a patient with Alzheimer disease. *Egypt Heart J* 2016; **68**: 197–199.
8. Vogel SM, Mican LM, Smith TL. Donepezil-induced QTc prolongation: A case report. *Ment Health Clin* 2019; **9**: 128–132.
9. Kuwahata S, Takenaka T, Motoya T, Masuda K, Yonezawa H, Shinchi S, et al. Effect of QT prolongation in patients taking cholinesterase inhibitors (donepezil) for Alzheimer's disease. *Circ Rep* 2021; **3**: 115–121.
10. Hagiwara-Nagasawa M, Kambayashi R, Goto A, Nunoi Y, Izumi-Nakaseko H, Chiba K, et al. Analysis of electropharmacological and proarrhythmic effects of donepezil using the halothane-anesthetized intact dogs and the conscious chronic atrioventricular block ones. *Naunyn Schmiedebergs Arch Pharmacol* 2021; **394**: 581–589.
11. Sugiyama A. Sensitive and reliable proarrhythmia in vivo animal models for predicting drug-induced Torsades de pointes in patients with remodelled hearts. *Br J Pharmacol* 2008; **154**: 1528–1537.
12. Johannesen L, Vicente J, Mason JW, Sanabria C, Waite-Labott K, Hong M, et al. Differentiating drug-induced multichannel block on the electrocardiogram: Randomized study of dofetilide, quinidine, ranolazine, and verapamil. *Clin Pharmacol Ther* 2014; **96**: 549–558.
13. Hagiwara-Nagasawa M, Kambayashi R, Goto A, Nunoi Y, Izumi-Nakaseko H, Takei Y, et al. Cardiohemodynamic and arrhythmogenic effects of the anti-atrial fibrillatory compound vanoxerine in halothane-anesthetized dogs. *Cardiovasc Toxicol* 2021; **21**: 206–215.
14. Antzelevitch C, Shimizu W, Yan GX, Sicouri S, Weissenburger J, Nesterenko VV, et al. The M cell: Its contribution to the ECG and to normal and abnormal electrical function of the heart. *J Cardiovasc Electrophysiol* 1999; **10**: 1124–1152.

Ryuichi Kambayashi, BSc

Ai Goto, BSc

Hiroko Izumi-Nakaseko, PhD

Akio Matsumoto, MD, PhD

Atsushi Sugiyama, MD, PhD

Department of Pharmacology (R.K., A.G., H.I.-N., A.S.),

Department of Aging Pharmacology (A.M., A.S.),

Faculty of Medicine, Toho University, Tokyo, Japan

Circulation Reports

Circ Rep 2021; **3**: 556–557

doi:10.1253/circrep.CR-21-0086

Measurement of Early and Late Repolarization Periods in Addition to QT Interval to Help Predict the Torsadogenic Risk of Donepezil Based on Reverse Translational Animal Research on Its Proarrhythmic Potential — Reply —

We read with great interest the Letter to the Editor from Kambayashi et al. We thank them for their insightful comments regarding our recent article.¹ In a previous study, these authors analyzed the effect of donepezil on the QT interval by separately measuring the early (J-T_{peak}) and late (T_{peak}-T_{end}) repolarization periods in canine models. Intravenous administration of 1 mg/kg donepezil hydrochloride significantly prolonged the corrected J-T_{peak}, in addition to modestly prolonging T_{peak}-T_{end}.² Kambayashi et al suggested that, in donepezil-treated patients, simultaneous measurement of the early and late repolarization periods, along with the QT interval, may predict the onset

of drug-associated Torsade de pointes.

In our study, we investigated the correlation between patients' background characteristics or blood biochemical findings and the corrected QT (QTc) interval in patients taking donepezil. On univariate analysis, QTc was associated with hemoglobin, serum calcium concentration, and the estimated glomerular filtration rate (eGFR). On multivariate analysis, serum potassium concentration and eGFR were significantly associated with QTc.¹ However, as Kambayashi et al point out, it was difficult to predict which of the patients taking donepezil would have QTc prolongation.

The T_{peak}-T_{end} interval serves as an index of transmural dispersion of repolarization.³ Prolongation of the T_{peak}-T_{end} interval was previously demonstrated in patients with hypertrophic cardiomyopathy,^{4,5} and has been reported to be a risk factor for ventricular fibrillation in patients with Brugada syndrome.⁶ In addition, prolongation of the T_{peak}-T_{end} interval was associated with an increased risk of sudden cardiac death.⁷

We agree with the suggestion of Kambayashi et al to measure and evaluate J-T_{peak} and T_{peak}-T_{end} to predict fatal arrhythmias in patients taking donepezil. However, there are some debatable points that need to be resolved in the

Received July 2, 2021; accepted July 3, 2021; J-STAGE Advance Publication released online August 6, 2021 Time for primary review: 1 day

Mailing address: So Kuwahata, MD, PhD, Department of Internal and Cardiovascular Medicine, Tarumizu Chuo Hospital, Tarumizu Municipal Medical Center, 1-140 Kinko-cho, Tarumizu 891-2124, Japan. E-mail: sokuwahata@ybb.ne.jp

All rights are reserved to the Japanese Circulation Society. For permissions, please e-mail: cr@j-circ.or.jp

ISSN-2434-0790

future. First, the previous study Kambayashi et al refer to was an animal experiment in canines, and it is unclear whether the results can be extrapolated to humans.² Second, their study involved plasma concentrations that were 30-fold higher than the clinically effective concentrations.² In that study, Kambayashi et al said that intravenous administration of donepezil hydrochloride at concentrations as high as 1 mg/kg may have reached toxicological concentrations that could induce respiratory arrest in the anesthetized dogs.² Further clinical studies within the effective plasma concentration range for humans are needed to prove this result in a clinical setting. Third, it is difficult to measure the QT interval separately for the early and late periods; it may not be easy for everyone to do. T-wave morphology has substantial changes, such as flattened or notched T-waves.⁸ In the study of Kambayashi et al, when the end of the T wave was unclear, the end was estimated using the monophasic action potential signal as a guide.² An ambiguous definition of the peak of the T-wave can cause large variations in measurements.⁸ Therefore, the objectivity of the measured values may not be maintained and may not be suitable for screening that needs to be performed within a short time. Automatic measurements of J-T_{peak} and T_{peak}-T_{end} intervals have already been reported,⁸ but are not yet commonly used. We hope that electrocardiograms will be equipped with automatic measurement of J-T_{peak} and T_{peak}-T_{end} intervals, and that screening will become widespread.

Acknowledgments

None.

Sources of Funding

This research received no grant from any funding agency in the public, commercial, or not-for-profit sectors.

Disclosures

The authors declare that there are no conflicts of interest.

References

1. Kuwahata S, Takenaka T, Motoya T, Masuda K, Yonezawa H, Shintchi S, et al. Effect of QT prolongation in patients taking cholinesterase inhibitors (donepezil) for Alzheimer's disease. *Circ Rep* 2021; **3**: 115–121.
2. Hagiwara-Nagasawa M, Kambayashi R, Goto A, Nunoi Y, Izumi-Nakaseko H, Chiba K, et al. Analysis of electropharmacological and proarrhythmic effects of donepezil using the halothane-anesthetized intact dogs and the conscious chronic atrioventricular block ones. *Naunyn-Schmiedeberg Arch Pharmacol* 2021; **394**: 581–589.
3. Yan GX, Antzelevitch C. Cellular basis for the normal T wave and the electrocardiographic manifestations of the long-QT syndrome. *Circulation* 1998; **98**: 1928–1936.
4. Shimizu M, Ino H, Okeie K, Yamaguchi M, Nagata M, Hayashi K, et al. T-peak to T-end interval may be a better predictor of high-risk patients with hypertrophic cardiomyopathy associated with a cardiac troponin I mutation than QT dispersion. *Clin Cardiol* 2002; **25**: 335–339.
5. Riza Demir A, Celik Ö, Sevinç S, Uygur B, Kahraman S, Yilmaz E, et al. The relationship between myocardial fibrosis detected by cardiac magnetic resonance and Tp-e interval, 5-year sudden cardiac death risk score in hypertrophic cardiomyopathy patients. *Ann Noninvasive Electrocardiol* 2019; **24**: e12672.
6. Castro Hevia J, Antzelevitch C, Tornés Bázaga F, Dorantes Sánchez M, Dorticós Balea F, Zayas Molina R, et al. Tpeak-Tend and Tpeak-Tend dispersion as risk factors for ventricular tachycardia/ventricular fibrillation in patients with the Brugada syndrome. *J Am Coll Cardiol* 2006; **47**: 1828–1834.
7. Panikkath R, Reinier K, Uy-Evanado A, Teodorescu C, Hattenhauer J, Mariani R, et al. Prolonged Tpeak-to-tend interval on the resting ECG is associated with increased risk of sudden cardiac death. *Circ Arrhythm Electrophysiol* 2011; **4**: 441–447.
8. Johannesen L, Vicente J, Hosseini M, Strauss DG. Automated algorithm for J-Tpeak and Tpeak-Tend assessment of drug-induced proarrhythmia risk. *PLoS One* 2016; **11**: e0166925.

So Kuwahata, MD, PhD

Toshiro Motoya, PhD

Toshihiro Takenaka, MD, PhD

Department of Internal and Cardiovascular Medicine (S.K., T.T.);

Pharmaceutical Department (T.M.),

Tarumizu Chuo Hospital,

Tarumizu Municipal Medical Center,

Tarumizu, Japan