New Products from the Reaction of Acetyllycoctonine with *N*-Bromosuccinimide (NBS)

Xiang-Li SHEN and Feng-Peng WANG*

Department of Chemistry of Medicinal Natural Products, West China College of Pharmacy, Sichuan University; No. 17, Duan 3, Renmin Nan Road, Chengdu 610041, P. R. China. Received April 8, 2004; accepted May 24, 2004

Six new artificial products 11–16 were obtained from the reaction of acetyllycoctonine (10) with *N*-bromosuccinimide (NBS). The structures of these compounds were established on the basis of spectral data. It is emphasized that the varieties and yields of the products in this reaction depended greatly upon reaction conditions and the types of substrate.

Key words C₁₉-diterpenoid alkaloid; acetyllycoctonine; oxidation; *N*-deethylation; imine

Although many papers on oxidation involving the nitrogen atom of the C₁₉-diterpenoid alkaloids with various oxidizers such as KMnO₄, OsO₄, CrO₃, Ag₂O, *m*-CPBA, etc. have been reported,¹⁾ there have been no prior reports on preparation of the imines starting from these alkaloids with NBS. In the previous works on modification of C₁₉-diterpenoid alkaloids, we reported the reaction of the aconitine-type alkaloids 3-acetylpseudoaconine (1),^{2,3)} 3,13-diacetylunaconitine (2),³⁾ 3-acetylunaconitine (3),^{4,5)} (4),⁶⁾ (5), and (6),⁷⁾ with *N*-bromosuccinimide (NBS) to give the *N*-deethyl and/or the imine derivatives. In addition, it was found that the varieties and yields of the resultants were greatly depended upon reaction conditions.³⁾ Only a paper reported by Marion and colleagues described that the lycoctonine-type alkaloid delcosine (7) exposed to NBS (80 °C, 68 h) afforded the lactam (8).⁸⁾ To observe application of this method, we turned our attention to the lycoctonine-type C₁₉-diterpenoid alkaloids which differ from the aconitine-type ones by an oxygenated group at C-7. After NBS oxidation of lycoctonine (9) to give the complex products, its acetyl derivative acetyllycoctonine (10) was used for this oxidation reaction. Meanwhile, we have tried hard to isolate others by-products than the major ones to clearly understand the reaction process. In this paper, we wish to report the isolation and identification of the new products produced by the NBS oxidation of 10.

Results and Discussion

Treatment of acetyllycoctonine (10) derived from lycoctonine (9) from the roots of *Delphinium potaninii*⁹⁾ in *t*-BuOH with NBS at 50 °C for 2 h followed by a general work-up gave a residue which was chromatographed on silica gel column repeatedly to afford the major product 11 (33.5%), in addition to the minor ones 12 (2.9%), 13 (10.1%), 14 (2.5%), 15 (2.4%), and 16 (1.5%). They were obtained as amorphous powder and their molecular formulae were also established based on HR-MS and ¹³C-NMR data. A comparison to the ¹H- and ¹³C-NMR spectra of 11 (Table 1 and Experimental) with those of acetyllycoctonine (10) showed that the additional signals at δ_H 9.19 (s) and δ_C 174.8 (d) for an iminium moiety and an obvious downfield shift effect from δ 1.02 to δ 1.53 belong to the *N*-ethyl group. Another chemical shift of the carbons attached to the nitrogen atoms C-17, C-19, C-21 also shifted downfield. Thus, surprisingly, its structure was assigned as 11, which is very stable to the acid such as 5%

HCl or base (25% NH₄OH). Compound 12 as a hydrolyzed derivative of 11 was established easily on the basis of comparison of their ¹H- (¹³C)-NMR spectra.

The ¹H- (¹³C)-NMR spectra of compounds 13, 14, and 15 showed absence of the *N*-ethyl group and presence of the characteristic signals at δ_H 7.72 (br s), δ_C 167.5 (d); δ_H 4.04 (br s); δ_H 6.25 (d, *J*=4.4 Hz), δ_C 173.7 (s) for an imine group, an NH group and a lactam moiety, respectively. From these observations, their structures were determined. Finally, the exhibition of an amide group (δ_C 171.3 s, 22.2 t) in the ¹³C-NMR spectrum of 16 led to establishment of its structure.

The formation of compounds 11–16 can be explained by

* To whom correspondence should be addressed. e-mail: wfp@wcums.edu.cn

Table 1. ^{13}C -NMR Data of Compounds **10**,¹¹⁾ **11**, **12**, **13**, **14**, **15**, and **16**

Carbon	10	11	12	13	14	15	16
1	84.0	79.1	80.1	81.9	82.5	83.6	81.0
2	26.1	17.7	18.2	20.0	25.0	25.1	25.0
3	31.9	23.4	24.3	24.5	28.8	28.5	30.8
4	37.2	45.4	48.1	46.1	37.0	47.1	36.1
5	43.3	40.8	41.7	42.9	48.3	49.2	50.1
6	90.9	86.2	82.3	90.6	91.0	91.7	90.6
7	88.5	86.8	86.8	86.5	84.0	85.3	83.8
8	77.5	76.1	77.1	77.0	Missed	76.2	77.4
9	50.4	41.6	42.3	42.9	43.6	42.8	43.3
10	38.1	42.6	44.5	45.3	45.1	44.6	45.7
11	49.0	50.6	51.6	50.5	48.5	48.5	47.9
12	28.7	29.7	30.7	30.4	29.4	28.6	27.9
13	46.1	36.4	37.2	38.0	38.5	37.7	38.2
14	84.0	82.7	83.4	83.9	84.2	81.7	83.8
15	33.7	32.5	32.8	33.1	33.4	33.1	33.6
16	82.6	81.6	82.3	80.7	82.9	81.4	82.5
17	64.6	69.7	69.7	64.2	60.4	59.2	61.7
18	69.1	64.1	64.7	65.7	69.0	65.6	68.7
19	52.4	174.8	178.6	167.5	48.1	173.3	45.1
21	51.0	57.4	57.5	—	—	—	171.3
22	14.1	12.3	13.1	—	—	—	22.2
1-OCH ₃	55.7	55.3	56.0	56.3	55.7	55.7	55.2
6-OCH ₃	57.8	56.7	57.6	57.7	57.7	57.8	58.3
14-OCH ₃	58.0	57.8	58.6	58.6	57.9	58.5	56.3
16-OCH ₃	56.3	55.4	56.2	56.4	56.2	56.4	57.8
18-OAc	170.9	169.3	—	170.4	170.8	170.2	170.7
	20.8	19.9	—	20.6	20.7	20.7	20.7

the mechanism depicted in Fig. 1. Compound **14** was produced by the intermediate A from **10** through a one-step process (route a) with loss of ethylene and HBr. **14** was subjected first in a nucleophilic addition followed by the α -elimination of HBr to form **13**. Another elimination of the intermediate A (route b) afforded two iso-imminium salts **11** or **12**, and C; the former gave **13** by an Cope elimination, and the latter was attacked by OH^- to form the *N,O*-mixed acetal D followed by further oxidation to give **16**. Similarly, the lactam **15** could be formed through the intermediate E from **14** via F and G.

It is worthy to note that the major product is the iminium salt (**11**) instead of the imine or *N*-deethyl derivative, as compounds **1**–**6** belong to the aconitine-type alkaloids when treatment of acetyllycoctonine (**10**) under optimized conditions (6 mmol NBS, 50 °C, 2 h).¹⁰⁾ This indicated these reactions involving the nitrogen atom were greatly affected by not only the 7-oxygenated substituents such as OH group but also the substituent at C-3³⁾ or C-18 in addition to the reaction conditions.³⁾ This is also a new method for preparation the imines of lycoctonine-type alkaloids, but, it is still necessary to promote the yield of the desired products. The results described here should also prove helpful in further research on scope and limitation of the NBS oxidation using various analogues of lycoctonine.

Experimental

General Experimental Procedures ^1H - and ^{13}C -NMR spectra were recorded on a Varian INOVA 400/54 or a Bruker AC-E 200 spectrometer, in CDCl_3 with TMS as internal standard. MS were measured on Finnigan LCO and a Micromass Auto Spec Ultima-Tof spectrometer.

Reaction of Acetyllycoctonine (10) with NBS To a solution of acetyllycoctonine (**10**) (970 mg, 1.90 mmol) in *t*-BuOH (170 ml), which was derived from lycoctonine (**9**) isolated from the roots of *Delphinium potaninii*,⁷⁾

Fig. 1. A Plausible Mechanism from **10** to Compounds **11**, **12**, **13**, **14**, **15**, and **16**

NBS (1063 mg, 6.00 mmol) was added and the solution was heated at 50 °C for 2 h. Evaporation, basifying (25% NH_4OH , 50 ml), extraction (CHCl_3 , 40 ml $\times$ 3), drying (anhydrous Na_2SO_4), removal of solvent and column chromatography (silica gel H, CHCl_3 – CH_3OH /98 : 2–75 : 25) afforded the pure compounds **13** (98 mg, 10.1%), **15** (23 mg, 2.4%), and fractions A, B, and C. Fraction A was subjected column chromatography (silica gel H, petroleum ether–acetone/5 : 2) to afford compound **16** (15 mg, 1.5%). Similarly, separa-

tions using silica gel column chromatography of fraction B (silica gel H, CHCl_3 – CH_3OH /97:3) and fraction C [silica gel H, CHCl_3 – CH_3OH /9:1 (saturated by 25% NH_4OH)] afforded the compounds **14** (25 mg, 2.5%), **11** (325 mg, 33.5%), and **12** (28 mg, 2.9%), respectively.

Compound 11: White amorphous powder. $\text{C}_{27}\text{H}_{42}\text{NO}_8$, HR-EI-MS m/z 508.2784 (Calcd for $\text{C}_{27}\text{H}_{42}\text{NO}_8$, 508.2832); mp 129–130.5 °C; $[\alpha]_{\text{D}}^{20} +48.3^\circ$ ($c=0.785$, CHCl_3); $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 1.53 (3H, t, $J=7.2$ Hz, $N\text{-CH}_2\text{CH}_3$), 2.16 (3H, s, OAc), 3.18, 3.38, 3.43, 3.46 (each 3H, s, $\text{OCH}_3\times 4$), 3.70 (1H, t, $J=4.4$ Hz, H-14 β), 4.72, 4.86 (each 1H, ABq, $J=11.6$ Hz, H₂-18), 9.19 (1H, s, H-19); $^{13}\text{C-NMR}$ (50 MHz, CDCl_3): see Table 1; EI-MS: m/z 509 ($\text{M}+1$, 100).

Compound 12: White amorphous powder. $\text{C}_{25}\text{H}_{40}\text{NO}_7$, HR-EI-MS m/z 466.2743 (Calcd for $\text{C}_{25}\text{H}_{40}\text{NO}_7$, 466.2726); mp 132–133 °C; $[\alpha]_{\text{D}}^{20} +62.1^\circ$ ($c=0.580$, CHCl_3); $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 1.51 (3H, t, $J=7.2$ Hz, $N\text{-CH}_2\text{CH}_3$), 3.15, 3.38, 3.43, 3.49 (each 3H, s, $\text{OCH}_3\times 4$), 3.70 (1H, t, $J=4.4$ Hz, H-14 β), 9.42 (1H, s, H-19); $^{13}\text{C-NMR}$ (100 MHz, CDCl_3): see Table 1; EI-MS: m/z 466 (M^+ , 5), 436 (M-30, 55), 406 (M-60, 100).

Compound 13: White amorphous powder. $\text{C}_{25}\text{H}_{37}\text{NO}_8$, HR-EI-MS m/z 479.2501 (Calcd for $\text{C}_{25}\text{H}_{37}\text{NO}_8$, 479.2519); mp 136–137 °C; $[\alpha]_{\text{D}}^{20} +103.8^\circ$ ($c=0.530$, CHCl_3); $^1\text{H-NMR}$ (200 MHz, CDCl_3) δ : 2.09 (3H, s, OAc), 3.13, 3.33, 3.34, 3.39 (each 3H, s, $\text{OCH}_3\times 4$), 3.64 (1H, t, $J=4.4$ Hz, H-14 β), 4.21, 4.31 (each 1H, ABq, $J=11.4$ Hz, H₂-18), 7.72 (1H, brs, H-19); $^{13}\text{C-NMR}$ (50 MHz, CDCl_3): see Table 1; EI-MS: m/z 479 (M^+ , 100), 464 (M-15, 95), 448 (M-31, 90).

Compound 14: White amorphous powder. $\text{C}_{25}\text{H}_{39}\text{NO}_8$, HR-EI-MS m/z 481.2688 (Calcd for $\text{C}_{25}\text{H}_{39}\text{NO}_8$, 481.2676); mp 63–64.5 °C; $[\alpha]_{\text{D}}^{20} +55.7^\circ$ ($c=0.415$, CHCl_3); $^1\text{H-NMR}$ (200 MHz, CDCl_3) δ : 2.07 (3H, s, OAc), 3.28, 3.32, 3.37, 3.42 (each 3H, s, $\text{OCH}_3\times 4$), 3.60 (1H, t, $J=4.6$ Hz, H-14 β), 3.88 (2H, s, H₂-18); $^{13}\text{C-NMR}$ (50 MHz, CDCl_3): see Table 1; EI-MS: m/z 481 (M^+ , 5), 466 (M-15, 65), 450 (M-31, 100).

Compound 15: White amorphous powder. $\text{C}_{25}\text{H}_{37}\text{NO}_9$, HR-EI-MS m/z 495.2471 (Calcd for $\text{C}_{25}\text{H}_{37}\text{NO}_9$, 495.2468); mp 97–98.5 °C; $[\alpha]_{\text{D}}^{20} +64.9^\circ$ ($c=0.535$, CHCl_3); $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 2.08 (3H, s, OAc), 3.26, 3.34, 3.40, 3.43 (each 3H, s, $\text{OCH}_3\times 4$), 3.65 (1H, t, $J=4.4$ Hz, H-14 β),

4.29, 4.58 (each 1H, ABq, $J=12.0$ Hz, H₂-18), 6.25 (1H, d, $J=4.4$ Hz, NH); $^{13}\text{C-NMR}$ (50 MHz, CDCl_3): see Table 1; EI-MS: m/z 495 (M^+ , 15), 480 (M-15, 85), 464 (M-31, 20).

Compound 16: White amorphous powder. $\text{C}_{27}\text{H}_{41}\text{NO}_9$, HR-EI-MS m/z 523.2757 (Calcd for $\text{C}_{27}\text{H}_{41}\text{NO}_9$, 523.2781); mp 89–91 °C; $[\alpha]_{\text{D}}^{20} +33.2^\circ$ ($c=0.845$, CHCl_3); $^1\text{H-NMR}$ (200 MHz, CDCl_3) δ : 2.05 (3H, s, $N\text{-COCH}_3$), 2.14 (3H, s, OAc), 3.20, 3.32, 3.37, 3.39 (each 3H, s, $\text{OCH}_3\times 4$), 3.61 (1H, t, $J=4.6$ Hz, H-14 β), 3.93, 4.02 (each 1H, ABq, $J=11.2$ Hz, H₂-18); $^{13}\text{C-NMR}$ (50 MHz, CDCl_3): see Table 1; EI-MS: m/z 523 (M^+ , 10), 505 (M-18, 10), 476 (M-47, 100).

Acknowledgment This work was supported by the National Nature Science Foundation of China (No. 3007088).

References

- 1) Wang F. P., Liang X. T., "The Alkaloids: Chemistry and Pharmacology," ed. by Cordell G. A., Academic Press, New York, 1992, pp. 151–247.
- 2) Wang F. P., Fan J. Z., Li Z. B., Li B. G., *Chin. Chem. Lett.*, **10**, 375–378 (1999).
- 3) Wang F. P., Li Z. B., Yang J. S., Li B. G., *Chin. Chem. Lett.*, **10**, 453–456 (1999).
- 4) Xu L., Chen Q. H., Wang F. P., *Tetrahedron*, **58**, 4267–4271 (2002).
- 5) Chen Q. H., Xu L., Wang F. P., *Tetrahedron*, **58**, 9431–9444 (2002).
- 6) Chen Q. H., Xu L., Wang F. P., *Heterocycles*, **57**, 2357–2363 (2002).
- 7) Chen Q. H., Xu L., Wang F. P., *Chin. Chem. Lett.*, **14**, 147–150 (2003).
- 8) Anet R., Clayton D. W., Marion L., *Can. J. Chem.*, **35**, 397–408 (1957).
- 9) Sheng X. L., Chen D. L., Jian X. X., Wang F. P., *West China J. Pharm. Sci.*, **16**, 4–7 (2001).
- 10) Wang F. P., unpublished data.
- 11) Pelletier S. W., Mody N. V., Sawhney R. S., Bhattacharyya J., *Heterocycles*, **7**, 327–339 (1977).