Phenyl Polypropanoids from *Lindelofia stylosa*

Muhammad Iqbal CHOUDHARY,^{*,a} Afshan BEGUM,^a Ahmed ABBASKHAN,^a Asma AJAZ,^a SHAFIQUE-UR-REHMAN,^b and ATTA-UR-RAHMAN^a

^aH.E.J. Research Institute of Chemistry, International Center for Chemical Sciences, University of Karachi; Karachi-75270, Pakistan; and ^bDepartment of Botany, University of Azad Jammu and Kashmir; Muzafarabad-13100, Pakistan.

Received May 2, 2005; accepted June 30, 2005

A phytochemical investigation on the aerial parts of *Lindelofia stylosa* has resulted in the isolation of seven phenyl propanoids 1–7. This includes three analogs of lithospermic acid 1–3, along with rosmarinic acid (4) and its ester derivatives 5–7. Compound 1 was identified as a new natural product. These compounds were studied for their antioxidant properties.

Key words *Lindelofia stylosa*; rosmarinic acid; rosmarinic acid derivative; lithospermic acid; antioxidant activity

About ten species of genus *Lindelofia* (Boraginaceae) are found in Central and West Asia and in Himalayan regions of Pakistan. *Lindelofia stylosa* (KAR. & KIR.) is a perennial herb reported to contain pyrrolizidine alkaloids.^{1–8} In continuation of our studies on the biologically active phytochemicals, we investigated the whole herb of *L. stylosa* of Pakistani origin. The present work resulted in the isolation of compounds 1–7. The structures of the isolated compounds were elucidated mainly with the help of NMR spectroscopic techniques.

Three members of the phenyl propanoid class (compounds 1–3) were isolated from the EtOAc fraction of the plant which were identified to be the derivatives of lithospermic acid.

New compound 1 was isolated as a brown yellow gum. The compound showed an M^+ at m/z 579.1493 in HR-FAB-MS (-ve), in agreement with the formula $C_{30}H_{28}O_{12}$ (Calcd 579.1502). Its IR spectrum showed the presence of an α,β -unsaturated carbonyl system at 1722 cm^{-1} . The overall spectral data of compound 1 closely resembled the known compound dimethyl lithospermate (2),⁹ the only difference between the two compounds being the appearance of signals for $O-CH_2CH_3$ moiety in the NMR spectra of 1. The $O-CH_2CH_3$ protons appeared as an AB quartet at δ 4.14 ($J=14.2\text{ Hz}$, $J=7.0\text{ Hz}$), and a triplet at δ 1.19 ($J=7.0\text{ Hz}$), which corresponded to the carbons resonated at δ 62.4 and 14.3, respectively. This indicated that compound 1 is an ethyl ester of lithospermic acid.

Compound 2 exhibited an M^+ at m/z 565.1343 in the HR-FAB-MS (-ve), which corresponded to the formula $C_{29}H_{26}O_{12}$ (Calcd 565.1346). The spectral data of compound 2 was found to be identical with the reported dimethyl lithospermate, which was first isolated from *Salvia miltiorrhiza*.⁹

Compound 3 showed an M^+ at m/z 745.1766 in the HR-FAB-MS (-ve), suggestive of the formula $C_{38}H_{34}O_{16}$ (Calcd 745.1768). The overall spectral data of compound 3 was identified as a known methyl ester derivative of lithospermic acid B, which was isolated from *Salvia przewalskii*.¹⁰

The comparison of ^1H - and ^{13}C -NMR data and physical data with those reported in the literature, indicated compounds 4–7 were esters of rosmarinic acid (4),^{11,12} *O*-methyl rosmarinic acid (5),¹³ ethyl rosmarinic acid (6)¹⁴ and butyl derivative (7).¹⁵ Their isolation from *L. stylosa* is reported here for the first time.

The isolated compounds 1–7 were found to possess radical scavenging properties in DPPH radical assay, among which, compounds 3 and 4 have exhibited the strongest activities. The rosmarinic acid (4) was found to be as active as the standard (3-*t*-butyl-4-hydroxyanisole).

Compound 4 has been reported in previous studies to be potent antioxidant.^{16,17} It increases the prostaglandin E₂ production and reduces the production of leukotriene B₄ in human polymorphonuclear leukocytes.¹⁸ It is also an inhibitor of complement C3-convertase^{19,20} as well as an inhibitor of lipid peroxidation.²¹

Our results have indicated that the derivatives of lithospermic acid, possessing a C-19 lactate moiety, are more active than those without such substituents, such as compounds 1 and 2. The study of antioxidant potentials of compounds 1–7 was carried out by using Fe^{2+} -chelating and superoxide scavenging assays. The results demonstrated the selectivity of compounds 3–5 in these two assays, while the others were inactive. Interestingly, compound 3 has shown activity which is comparable to the standard propyl gallate used in the two assays. Our results demonstrated that the rosmarinic acid ester derivatives 4–5 and compound 3 possess significant antioxidant activities in various assays (see Table 1).

Experimental

General Experimental Procedures Melting points were determined on a Yanaco apparatus. UV spectra were measured on a Shimadzu UV240 ma-

Fig. 1. Structures of Compounds 1–7

* To whom correspondence should be addressed. e-mail: hej@cyber.net.pk

Table 1. Activities of Compounds 1–7 (IC₅₀ mM) in Various Antioxidant Assays

Compound	DPPH radical scavenging activity	Superoxide anion scavenging activity	Iron chelating assay
Ethyl lithospermate (1)	0.1028	— ^{b)}	—
Dimethyl lithospermate (2)	0.1197	—	—
Methyl lithospermate B (3)	0.0506	0.113	0.053
Rosmarinic acid (4)	0.0801	0.282	0.034
Methyl rosmarinate (5)	0.1456	0.443	0.092
Ethyl rosmarinate (6)	0.0412	—	—
Butyl rosmarinate (7)	0.2706	—	—
3- <i>t</i> -Butyl-4-hydroxyanisole ^{a)}	0.0442	—	—
Propyl gallate ^{a)}	0.03	0.106	0.064

a) Standard reference compound. b) Inactive.

chine in MeOH solutions. IR spectra were recorded as KBr discs on a JASCO A-302 spectrometer. ¹H- (500 MHz) and ¹³C-NMR (125 MHz) spectra were recorded in CD₃OD solutions on a Bruker AV-500 machine with tetramethylsilane (TMS) as an internal standard and the data is given in δ (ppm). 2D NMR spectra were taken on a Bruker AMX 500 NMR spectrometer. Electron impact mass spectra (EI-MS) were taken at 70 eV on a Finnigan MAT-112 or MAT-312 instrument and major ions are presented as *m/z* (%). Fast bombardment (FAB)-MS were measured as glycerol matrix on a JEOL HX110 Mass spectrometer. TLC purification was carried out on pre-coated silica gel cards (E. Merck) and the spots were observed first under UV (254 nm) and then sprayed with cerium(IV)sulfate reagent and heated until coloration developed. Recycling preparative HPLC (RPHPLC) was used for final purification (JAI LC-908W, Japan Analytical Industry Co. Ltd.) with a column YMC ODS H-80 or L-80 (YMC, Japan).

Plant Material Whole plants of *Lindelia stylosa* (KAR. & KIR.) were collected from Britswarr Gali, Leepa Valley, Azad Kashmir, Pakistan, in September–October 2002 by Prof. Shafiq-ur-Rehman, Department of Botany, University of Azad Jammu & Kashmir (voucher specimen No AJKU-786165).

Extraction and Isolation Air-dried and powdered *Lindelia stylosa* (3 kg) was extracted thrice (each for one week) with MeOH (6 l) at room temperature. The resulting MeOH extract (ca. 150 g) was partitioned between hexane, EtOAc, *n*-butanol and water to obtain 15, 24, 50 and 40 g of extracts, respectively. The EtOAc fraction was chromatographed on a silica gel column and eluted with hexane, hexane–EtOAc (1:1, 2×500 ml), EtOAc (100%, 2×750 ml), EtOAc–MeOH (2:1, 2×500 ml) and EtOAc–MeOH (1:1, 2×750 ml). Fraction eluted at EtOAc–MeOH (2:1, 3.5 g) was again chromatographed on a silica gel column with acetone–CHCl₃ (20 to 80%, 2×300 ml) to obtain seven fractions, among which fractions 3 and 4 (350 mg) were combined and subjected to purification on a recycling preparative HPLC (RPHPLC) to obtain compound 3 (40 mg, 1.33×10^{−3}%, *t_R* 52 min), using H₂O–MeOH (1:1); H-80 column with flow rate of 4 ml/min. The fraction eluted at MeOH–EtOAc; 1:1 (5.0 g) from silica gel column chromatography was passed through a polyamide column and eluted with CHCl₃ with increasing proportions of methanol.

Among these, two fractions, 7–12 and 16–20, obtained by solvent gradient of 10–25% MeOH–CHCl₃ were combined and subjected to RPHPLC purification. The use of ODS chromatography (L-80 column) on HPLC of fr. 7–12 (310 mg) using H₂O–MeOH (1:1) as a mobile phase at a flow rate of 4 ml resulted into two unresolved fractions A and B. These fractions were subjected again to RPHPLC with modified separation conditions using a H-80 column, H₂O–MeOH (1:2) and flow rate of 3 ml per min, for isolation of compound 5 *t_R* time 30 min (120 mg, 4.0×10^{−3}%). Compounds 6 (110 mg, 3.66×10^{−3}%) and 7 (20 mg, 6.66×10^{−4}%) were isolated under the similar RPHPLC conditions as those of compound 5 with retention time of 38 and 56 min, respectively. Both of these compounds were obtained from fraction B. Fractions 16–20 from polyamide column (162 mg) when subjected to RPHPLC with H₂O–MeOH (1:2) as a mobile phase, H-80 column and a flow rate of 3 ml per min, resulted in the isolation of compounds 2 (15 mg, 5.0×10^{−4}%) and 1 (20 mg, 6.66×10^{−4}%).

A part of the *n*-butanolic fraction (15 g) on treatment with HP-20 resin followed by polyamide column chromatography on 15–25% MeOH–CHCl₃ afforded a fraction (3 g) which was treated with Sephadex LH-20 and finally with RPHPLC on an L-80 column using H₂O–MeOH (1:1) as a mobile

phase to yield rosmarinic acid (4) (180 mg, 6.0×10^{−3}%) and some unidentified compounds.

Ethyl Lithospermate (1): C₃₀H₂₈O₁₂; Brown yellow gum, [α]_D²³ +65.6° (*c*=0.25, MeOH); UV (MeOH) λ_{max} (log ϵ 309 (3.46), 290 (3.46), 254 (3.46), 224 (3.56), 197 (3.84), 194 (5.04) nm; IR (KBr) ν_{max} 3300, 2933, 1722, 1608, 1517, 1444, 1265, 1166 cm^{−1}; HR FAB-MS (−ve) *m/z* 579.1493 (*M*[−]1); EI-MS *m/z* 321 (6.7%), 280 (27.6%), 123 (100%), 77 (34.7%); ¹H-NMR δ (CD₃OD, 500 MHz): δ 7.71 (d, *J*=15.9 Hz, H-7), 7.19 (d, *J*=8.4 Hz, H-6), 6.82 (d, *J*=8.4 Hz, H-5), 6.5–7.0 (overlap., H-27), 6.5–7.0 (overlap., H-23), 6.5–7.0 (overlap., H-17), 6.5–7.0 (overlap., H-26), 6.5–7.0 (overlap., H-16), 6.5–6.9 (overlap., H-13), 6.28 (d, *J*=15.9 Hz, H-8), 5.16 (m, H-10), 5.88 (d, *J*=4.8 Hz, H-21), 4.41 (d, *J*=4.8 Hz, H-20), 4.14 (q, *J*=14.2, 7.0 Hz, OCH₂), 3.68 (s, OMe), 3.02 (m, H-11), 1.19 (t, *J*=7.0 Hz, OCH₂CH₃). ¹³C-NMR: δ (CD₃OD, 125 MHz): δ 173.6 (C-19), 171.6 (C-18), 168.0 (C-9), 148.8 (C-3), 146.8 (C-14), 146.6 (C-24), 146.2 (C-4), 145.4 (C-25), 145.4 (C-15), 144.1 (C-7), 133.3 (C-22), 128.7 (C-12), 127.0 (C-2), 124.4 (C-1), 121.9 (C-17), 121.8 (C-6), 118.4 (C-27), 118.3 (C-5), 117.5 (C-13), 116.4 (C-8), 116.4 (C-26), 116.3 (C-16), 113.4 (C-23), 88.5 (C-21), 74.8 (C-10), 62.4 (OCH₂), 57.3 (C-20), 53.2 (OMe), 37.8 (C-11), 14.3 (OCH₂CH₃).

DPPH Radical Scavenging Assay The assay was performed according to the method developed by Lee *et al.*²² Five microliters of each sample was dissolved in DMSO and mixed with 95 μ l of DPPH in ethanol. The concentration of DPPH was maintained at 300 mM with variable concentrations of sample. The mixture was dispersed in a 96-well plate and incubated at 37 °C for 30 min. The absorbance at 515 nm was measured by microtitre plate reader (Spectramax plus 384 Molecular Device, U.S.A.), and percent radical scavenging activity was determined in comparison with the DMSO-treated control (3-*t*-butyl-4-hydroxyanisole).

Superoxide Anion Scavenging Assay The reaction mixture was prepared by mixing 280 μ M β -nicotinamide adenine dinucleotide reduced form (NADH), 80 μ M nitroble tetrazolium (NBT), 8 μ M phenazine methosulphate (PMS) and various concentrations of test samples in 200 μ l of 0.1 M phosphate buffer (pH 7.5). The NBT, NADH and PMS solutions were prepared in the same buffer. Test samples were dissolved in DMSO. The reaction was performed in 96-well microtitre plates at room temperature and absorbance was measured at 560 nm.²³

Measurement of Fe²⁺ Chelating Ability The Fe²⁺ chelating ability was determined according to the modified method of Decker and Welch.²⁴ The concentrations of Fe²⁺ ion were measured from the formation of ferrous ion-ferrozine complex. The pure compounds (31.25 to 1 mM) were mixed with 0.4 mM FeCl₂ and 1 mM ferrozine (ratio 10:1:2). The mixture was shaken and left at room temperature for 10 mins. The absorbance of the resulting mixtures were measured at 562 nm by microtitre plate reader. A lower absorbance of the reaction mixture indicated a higher Fe²⁺-chelating ability.

Acknowledgments The authors are grateful to the Luminar and Gold Electronics (LG), Pakistan, for financial support to Afshan Begum. They also gratefully acknowledge the useful suggestions of Prof. Y. Tsuda, Visiting Professor at the H.E.J. Research Institute of Chemistry (Kanazawa University, Kanazawa, Japan).

References

- Asibal C. F., Glinski J. A., Gelbanum L. T., Zalkow L. H., *J. Nat. Prod.*, **52**, 109 (1989).
- Zalkow L. H., Bonetti S. J., Gelbanum L. T., Gordon M. M., Patill B. B., Shani A., Van Derveer D., *J. Nat. Prod.*, **42**, 603 (1979).
- Men'shikov G. P., *J. Gen. Chem., USSR* (Engl. Transl.), **11**, 209 (1941).
- Men'shikov G. P., *J. Gen. Chem., USSR* (Engl. Transl.), **15**, 225 (1945).
- Men'shikov G. P., *J. Gen. Chem., USSR* (Engl. Transl.), **17**, 343 (1947).
- Men'shikov G. P., *J. Gen. Chem., USSR* (Engl. Transl.), **18**, 1736 (1948).
- Labenskii A. S., Men'shikov G. P., *J. Gen. Chem., USSR* (Engl. Transl.), **18**, 1836 (1948).
- Tsyrl'nikova L. G., Peresleni E. M., Sheinker Yu. N., *Zhur. Obshch. Khim.*, **32**, 2705–2709 (1962).
- Kang H. S., Chung H. Y., Jung J. H., Kang S. S., Choi J. S., *Arch. Pharm. Res.*, **20**, 496–500 (1997).
- Zhijun Wu., Mingan O., Chongren Y., Yunnan. *Zhiwu. Yanjiu.*, **21**,

- 512—516 (1999).
- 11) Kelley C. J., Mahajan J. R., Brooks C., Newbert L. A., Breneman W. R., Carmack M., *J. Org. Chem.*, **40**, 1804—1815 (1975).
- 12) Kelley C. J., Harruff R. C., Carmack M., *J. Org. Chem.*, **41**, 449—455 (1976).
- 13) Kuo Y. H., Lee S. M., Lai J. S., *J. Chin. C. Soc.*, **47**, 241—246 (2000).
- 14) Hou Z. F., Tu Y. Q., Li Y., *J. Chin. C. Soc.*, **49**, 255—258 (2002).
- 15) Haung H., Chao Q. R., Tan R. X., Sun H. D., Wang D. C., Ma J., Zhao S. X., *Planta Med.*, **65**, 92—93 (1999).
- 16) Lamaison J. L., Petitjean-Freytet C., Carnat A., *Ann. Pharm. Fr.*, **48**, 103—108 (1990).
- 17) Psotova J., Lasovsky J., Vicar J., *Biomed. Papers*, **147**, 147—153 (2003).
- 18) Al-Sereiti M. R., Abu-Amer K. M., Sen P., *Indian J. Exp. Biol.*, **37**, 124—130 (1999).
- 19) Engleberger W., Hadding U., Etschenberg E., Graf E., Leyck S., Winkelmann J., Parnham M. J., *Int. J. Immunopharmacol.*, **10**, 729—737 (1988).
- 20) Peake P. W., Pussell B. A., Martyn P., Timmermans V., Charlesworth J. A., *Int. J. Immunopharmacol.*, **13**, 853—857 (1991).
- 21) Naomi O., Akiko Y., Midori N., Toshikazu Y., *Carcinogenesis*, **25**, 549—557 (2004).
- 22) Lee S. K., Zakaria H., Chung H., Luvengi L., Gamez E. J. C., Mehta R. J., Kinghorn D., Pezzuto J. M., *Combinatorial Chemistry and High Throughput Screening*, **1**, 35—46 (1998).
- 23) Gaulejac N. S. C., Glories Y., Vivas N., *Food Research International*, **32**, 327—333 (1999).
- 24) Decker E. A., Welch B., *J. Agric. Food Chem.*, **38**, 674—677 (1990).