New Neolignans and Lignans from Vietnamese Medicinal Plant *Machilus odoratissima* NEES

PHAN Minh Giang,^{a,b} PHAN Tong Son,^a Katsuyoshi MATSUNAMI,^b and Hideaki OTSUKA^{*b}

^a Faculty of Chemistry, College of Natural Science, Vietnam National University; 19 Le Thanh Tong Street, Hanoi, Vietnam; and ^b Graduate School of Biomedical Sciences, Hiroshima University; 1–2–3 Kasumi, Minami-ku, Hiroshima 734–8553, Japan. Received September 29, 2005; accepted November 24, 2005

Four new natural neolignans and lignans, which were given the trivial names odoratisols A–D (1–4), together with (–)-licarin A, kachirachirol B, obovatifol, and machilin-I were isolated from the air-dried bark of the Vietnamese medicinal plant *Machilus odoratissima* NEES (Lauraceae). Their absolute structures were determined on the basis of spectroscopic analyses including circular dichroism spectra.

Key words *Machilus odoratissima*; Lauraceae; neolignan; lignan; odoratisol; absolute structure

Machilus is a genus in the Lauraceae family which includes twelve species distributed throughout Vietnam. They are *Machilus odoratissima* NEES, *M. thunbergii* SIEB. et ZUCC., *M. velutina* CHAMP. ex. BENTH., *M. oreophila* HANCE, *M. robustus* J. J. SON., *M. bonii* LEC., *M. coriacea* A. CHEV., *M. thunbergii* var. *condorensis* LEC., *M. parviflora* MEISSN., *M. platycarpa* CHUN., *M. macrophylla* HEMSLEY, and *M. cochinchinensis* LEC.^{1–3} *Machilus odoratissima* NEES (Vietnamese name *Khao nham*) is a timber tree growing up to a height of 8–10 m. The tree bark is used in the folk medicine as antiseptic and anti-inflammatory remedies. The leaves are used to treat snake bite and burn wounds.⁴ We carried out the first systematic study on *M. odoratissima* dealing with the isolation and structural elucidation of four new and four known neolignans and lignans.

The bark of *M. odoratissima* was air-dried in the shadow, powdered, and extracted with MeOH at room temperature. The MeOH extract was partitioned between H₂O and solvents of increasing polarities to afford *n*-hexane-, CH₂Cl₂-, EtOAc-, and 1-BuOH-soluble fractions. Column-chromatographic separation of the *n*-hexane- and CH₂Cl₂-soluble fractions led to the isolation of four new neolignans and lignans, named odoratisol A–D (1–4) and four known ones, (–)-licarin A, kachirachirol B, obovatifol, and machilin-I, which were determined by comparing their physical ([α]_D) and spectroscopic data with the literature values.^{5–9}

Odoratisol A (1) was isolated as an oil and its molecular formula was deduced to be C₂₁H₂₄O₅ from negative-ion high-resolution (HR)-FAB-MS. The IR spectrum indicated the presence of hydroxyl groups (3450 cm^{–1}) and aromatic rings (1609, 1517, 1458 cm^{–1}). The ¹H- (Table 1) and ¹³C-NMR (Table 2) spectroscopic data indicated that 1 had a planar structure of 5-methoxydehydrodiisoeugenol. The *trans* geometry of 1'-propenyl group was determined on the basis of the large coupling constant between H-7' and H-8' (*J*=15.6 Hz). The *trans* relationship of H-7 and H-8 was established based on ¹H-NMR spectroscopic data characteristic of *trans*-7-aryl-8-methyl-7,8-dihydro-benzofuranoid-type neolignans [δ _H 5.0 (1H, d, *J*=9.5 Hz, H-7), 3.37 (1H, dq, *J*=9.5, 6.6 Hz, H-8), 1.32 (3H, d, *J*=6.6 Hz, H-9)].^{10,11} The positions of the methoxyl groups at δ _H 3.81 (3H, s) and 3.82 (6H, s) were assigned using nuclear Overhauser effect spectroscopy (NOESY) (Fig. 2). 5-Methoxydehydrodiisoeugenol was reported previously from *Myristica fragrans*,¹² however,

its stereochemistries at C-7 and C-8 have not been determined at that time. In this study the circular dichroism (CD) spectrum was used to determine the absolute configuration of 1 to be the 7*S*,8*S* stereoisomer of 5-methoxydehydrodiisoeugenol from a positive Cotton effect at 242 nm (+4.35) and a negative one at 269 nm (–5.93), which were similar to those exhibited by (–)-licarin A.⁶ This configuration agreed well with the same sign of the optical rotations of 1 ([α]_D²⁵ –35.1°) and (–)-licarin A ([α]_D²⁵ –44.0°).⁶ Thus 1 was isolated for the first time from Nature and its absolute stereostructure was concluded to be (7*S*,8*S*)-7,8-dihydro-7-(4-hy-

Fig. 1. Absolute Structures of Compounds 1–4, (–)-Licarin A, Verrucosin, Austrobailignan-7, and Futokadsurin B

* To whom correspondence should be addressed. e-mail: hotsuka@hiroshima-u.ac.jp

Table 1. ^1H -NMR Spectroscopic Data of **1**–**4** (δ in ppm, J in Hz in Parentheses, 400 MHz, CDCl_3)

H	1	2	3	4
2	6.59 s	6.84 d (2.0)	6.97 d (1.7)	6.95 d (1.7)
5		6.78 d (8.5)	6.84 d (8.3)	6.83 d (8.3)
6	6.59 s		6.90 dd (8.3, 1.7)	6.87 dd (8.3, 1.7)
7	5.0 d (9.5)	4.75 d (3.2)	4.31 d (9.3)	4.27 d (9.3)
8	3.37 dq (9.5, 6.6)	4.27 dq (3.2, 6.4)	1.70 br dq (9.3, 6.6)	1.68 br dq (9.3, 6.6)
9	1.32 d (6.6)	1.09 d (6.4)	0.98 d (6.6)	0.95 d (6.6)
2'	6.72 s	6.91 d (2.0)	6.78 d (1.7)	6.70 d (1.7)
5'		6.88 d (8.3)	6.80 d (8.0)	6.71 d (8.0)
6'	6.69 s	6.84 dd (8.3, 2.0)	6.74 dd (8.0, 1.7)	6.79 dd (8.0, 1.7)
7'	6.29 d (15.6)	6.28 d (15.6)	5.03 d (8.8)	5.01 d (8.8)
8'	6.03 dq (15.6, 6.3)	6.08 dq (15.6, 6.6)	2.14 br dq (8.8, 6.8)	2.13 br dq (8.8, 7.1)
9'	1.79 d (6.3)	1.80 d (6.6)	0.58 d (6.8)	0.59 d (7.1)
MeO-3	3.82 s	3.819 s	3.82 s	3.82 s
MeO-5	3.82 s			
MeO-3'	3.81 s	3.822 s	3.77 s	
OCH_2O				5.85 s

All assignments were made on the basis of heteronuclear single quantum correlation (HSQC) and NOESY experiments.

Table 2. ^{13}C -NMR Spectroscopic Data of **1**–**4** (δ in ppm, 100 MHz, CDCl_3)

C	1	2	3	4
1	133.2	133.8	133.2	132.5
2	103.6	109.0	109.5	109.3
3	147.1	146.5	146.6	146.55
4	134.9	144.9	145.2	145.3
5	147.1	113.9	114.2	114.3
6	103.6	119.9	119.3	119.5
7	94.0	82.4	87.4	82.8
8	45.7	73.7	47.8	47.8
9	17.6	13.4	14.9	15.0
1'	132.3	131.9	132.8	135.2
2'	109.5	109.5	109.8	120.2
3'	144.2	145.6	146.2	147.4
4'	146.6	151.6	144.6	146.59
5'	136.4	119.1	113.9	107.7
6'	113.3	118.8	119.9	107.5
7'	131.0	130.5	83.1	87.5
8'	123.5	125.0	46.0	45.9
9'	18.3	18.5	15.0	14.9
MeO-3	56.4	55.8	55.86	55.9
MeO-5	56.4			
MeO-3'	56.0	55.9	55.9	
OCH_2O				100.9

Fig. 2. NOESY Correlations of Compounds **1**, **3**, and **4**

droxy-3,5-dimethoxyphenyl)-3'-methoxy-8-methyl-1'-*trans*-propenylbenzofuran.

Odoratisol B (**2**) was isolated as an oil and had the molecular formula $\text{C}_{20}\text{H}_{24}\text{O}_5$ based on negative-ion HR-FAB-MS. The IR spectrum indicated the presence of hydroxyl groups (3448 cm^{-1}) and aromatic rings (1603 , 1511 , 1458 cm^{-1}). The ^1H - (Table 1) and ^{13}C -NMR (Table 2) spectroscopic data of **2** were superimposable with those of machilin C,¹³⁾ including the *erythro* stereochemistry between H-7 and H-8 as shown by a small coupling constant ($J=3.2\text{ Hz}$) between them. However, the optical rotation of **2** ($[\alpha]_{\text{D}}^{25} +18.6^\circ$) was almost of the same in value but reverse in sign in comparison with that of machilin C ($[\alpha]_{\text{D}}^{25} -16.5^\circ$) showing the need to determine the absolute stereochemistry at two stereogenic elements at C-7 and C-8. The CD spectrum of **2** showed positive Cotton effect at 258 nm ($+0.38$) established the configu-

rations at C-7 and C-8 are *7S* and *8R* as in the cases of analogous neolignans of *erythro* series.¹⁴⁾ On the basis of these data, the structure of **2** was concluded to be *7S,8R-erythro-4'-hydroxy-3,2'-dimethoxy-4'-trans-propenyl-neolignan*.

Odoratisol C (**3**) was isolated as an oil, $[\alpha]_{\text{D}}^{25} -26.0^\circ$, and its molecular formula was characterized to be $\text{C}_{20}\text{H}_{24}\text{O}_5$ in negative-ion HR-FAB-MS. The IR spectrum indicated the presence of hydroxyl groups (3450 cm^{-1}) and aromatic rings (1607 , 1514 , 1457 cm^{-1}). The ^1H - (Table 1) and ^{13}C -NMR (Table 2) spectroscopic data showed the structural resemblance of **3** and verrucosin,¹⁵⁾ which, however, displayed a positive optical rotation ($[\alpha]_{\text{D}} +14.8^\circ$). The *trans* H-7/H-8, *trans* H-8/H-8', and *cis* H-7'/H-8' relative stereochemistries of the tetrahydrofuran ring were determined by comparing

the ^1H -NMR data [δ_{H} 0.98 (3H, d, $J=6.6$ Hz, H-9), 1.70 (1H, br dq, $J=9.3$, 6.6 Hz, H-8), 4.31 (1H, d, $J=9.3$ Hz, H-7); 0.58 (3H, d, $J=6.8$ Hz, H-9'), 2.14 (1H, br dq, $J=8.8$, 6.8 Hz, H-8'), 5.03 (1H, d, $J=8.8$ Hz, H-7')] with those reported in literature for 7,8-*trans*-8,8'-*trans*-7',8'-*cis*-configured tetrahydrofuran-type lignans.^{15–17} The *trans* H-7/H-8 and *cis* H-7'/H-8' configurations were in agreement with the upfield shift ($\Delta\delta_{\text{H}} -0.4$) of methyl proton signals on going from C-9 to C-9' due to the anisotropic effect of the aromatic group in the case of the *cis*-configuration of the aryl group at C-7' and methyl substituent at C-8'. The stereochemical assignments were supported by the NOESY spectrum (Fig. 2) of **3**, which showed NOEs between H₃-9 (δ_{H} 0.98) and H-7 (δ_{H} 4.31), between H₃-9 and H-8' (δ_{H} 2.14), and between H-7 and H-8', between H₃-9' (δ_{H} 0.58) and H-2' (δ_{H} 6.78).^{17,18} The locations of two 4-hydroxy-3-methoxyphenyl moieties were also confirmed by NOEs between H-6 [δ_{H} 6.90 (dd, $J=8.3$, 1.7 Hz)] and H-7 and between H-2' [δ_{H} 6.78 (d, $J=1.7$ Hz)] and H-7'. Since **3** had the same relative stereochemistry but opposite optical rotation in comparison with those of verrucosin, the absolute configurations at the C-7, C-8, C-7', and C-8' were deduced to be opposite to those of verrucosin. Thus the absolute structure of **3** was determined to be (7*R*,8*R*,7'*S*,8'*R*)-4-hydroxy-3-methoxy-4'-hydroxy-3'-methoxy-7,7'-epoxylignan.

Odoratisol D (**4**) was isolated as an oil and its molecular formula was determined $\text{C}_{20}\text{H}_{22}\text{O}_5$ by means of negative-ion HR-FAB-MS. The IR spectrum indicated the presence of hydroxyl groups (3450 cm^{-1}) and aromatic rings (1608 , 1517 , 1442 cm^{-1}). The ^1H -NMR (Table 1) spectrum showed the presence of a 4-hydroxy-3-methoxyphenyl and a monosubstituted 3,4-methylenedioxyphenyl systems, which contained an aromatic methoxyl group [δ_{H} 3.82 (3H, s)] and a methylenedioxy group [δ_{H} 5.85 (2H, s)], and a 2,5-disubstituted 3,4-dimethyltetrahydrofuran ring [δ_{H} 0.95 (3H, d, $J=6.6$ Hz, H-9), 1.68 (1H, br dq, $J=9.3$, 6.6 Hz, H-8), 4.27 (1H, d, $J=9.3$ Hz, H-7); 0.59 (3H, d, $J=7.1$ Hz, H-9'), 2.13 (1H, br dq, $J=8.8$, 7.1 Hz, H-8'), 5.01 (1H, d, $J=8.8$ Hz, H-7')]. The ^1H - and ^{13}C -NMR (Table 2) spectroscopic data of **4** resembled those of futokadsurin B¹⁸) except for the lack of an additional methoxyl group. The *trans* H-7/H-8, *trans* H-8/H-8', and *cis* H-7'/H-8' relative stereochemistry of the tetrahydrofuran ring were conclusive on the basis of the comparison of the ^1H -NMR data with those reported in literature.^{16–19} Upfield shift ($\Delta\delta_{\text{H}} -0.36$) of methyl proton signals on going from C-9 to C-9' agreed with *trans* H-7/H-8 and *cis* H-7'/H-8' configurations. NOEs observed between H₃-9 (δ_{H} 0.95) and H-7 (δ_{H} 4.27), between H₃-9 and H-8' (δ_{H} 2.13), and between H-7 and H-8', but not between H₃-9' (δ_{H} 0.59) and H-7' (δ_{H} 5.01) supported the stereochemical assignments.¹⁷ Thus **4** was concluded to have the same relative stereochemistry as that of **3**. NOESY spectrum (Fig. 2) of **4** showed the correlations between the methoxyl group (δ_{H} 3.82) and H-2, between H-2 (δ_{H} 6.95) and H-7 (δ_{H} 4.27), between H-6 (δ_{H} 6.87) and H-7, between H-2' (δ_{H} 6.70) and H-7' (δ_{H} 5.01), and between H-6' (δ_{H} 6.79) and H-7' confirmed the assignments of the position of the 4-hydroxy-3-methoxyphenyl moiety at C-7 and the 3',4'-methylenedioxyphenyl moiety at C-7' as shown in Fig. 2. To establish the absolute structure of **4** the CD spectra of **3** and **4** were measured and compared. Similar CD curves of **3** and **4** were seen, namely, **4** displayed

the Cotton effects at 213 nm (-1.25), 240 nm ($+0.95$), and 285 nm (-0.40) assuring the absolute structure of **4** to be (7*R*,8*R*,7'*S*,8'*R*)-4-hydroxy-3-methoxy-3',4'-methylenedioxy-7,7'-epoxylignan.

Experimental

General Procedure Optical rotations were measured on a JASCO P-1030 polarimeter. FT-IR spectra were recorded on a Horiba FT-710 spectrophotometer. ^1H - (400 MHz) and ^{13}C -NMR (100 MHz) spectra were recorded using a JEOL JNM- α 400 NMR spectrometer with tetramethylsilane as an internal standard. Negative-ion HR-FAB-MS were measured on a JEOL SX-102 mass spectrometer with PEG-400 as a calibration matrix. HPLC was carried out with a JASCO PU-1580 pump and an UV-2075 Plus detector (set at 210 nm) on YMC ODS columns (150 $\times$ 4.6 mm i.d. in analytical and 150 $\times$ 20 mm i.d. in preparative scales) at the corresponding flow rates of 0.5 and 5 ml/min. Silica (Si) gel 60 (0.063–0.200 mm, Merck, Germany) and reversed-phase octadecyl Si (ODS) gel (YMC, Japan) were used for open-column chromatography. TLC was carried out on Merck TLC plates (Si gel 60 F₂₅₄), and detected by spraying with 10% H_2SO_4 in 50% EtOH, followed by heating on a hot plate at 200 °C.

Plant Material The air-dried bark (2.0 kg) of *M. odoratissima* was collected in Province Thai Nguyen, Vietnam, and identified by Dr Nguyen Hoanh Coi of the Military Institute of Drug Controls (Hanoi, Vietnam), in June 2000. A voucher specimen (no. HCTN 2000-6) is deposited in the Laboratory of Chemistry of Natural Products, Faculty of Chemistry, Vietnam National University, Hanoi, Vietnam.

Extraction and Isolation of 1–10 The powdered air-dried bark of *M. odoratissima* (2.0 kg) was extracted with MeOH by percolation at room temperature (3 times, for 3 d each). After concentration by evaporation under reduced pressure, the resultant MeOH extract was suspended in H_2O and sequentially extracted with *n*-hexane, CH_2Cl_2 , EtOAc, and 1-BuOH. The *n*-hexane-soluble fraction (5.9 g) was separated on a Si gel open column using mixtures of *n*-hexane in EtOAc (10:1, 4:1, 2:1, 1:1). Five pooled fractions were collected on the basis of TLC patterns. Fraction 2 (1.8 g), fraction 3 (0.5 g), and fraction 4 (0.4 g) underwent the same treatment, first separation on an ODS gel open column (MeOH– H_2O , 3:1, 4:1), then purification on ODS preparative HPLC (MeOH– H_2O , 3:1) to give odoratisol A (**1**, 12.2 mg), odoratisol B (**2**, 7.0 mg), odoratisol C (**3**, 27.9 mg), odoratisol D (**4**, 40.2 mg), (–)-licarin A (0.48 g), machilin-I (13.4 mg), kachirachirol B (16.1 mg), and obovatifol (17.8 mg). Similar procedure was used to separate the CH_2Cl_2 -soluble fraction yielding **1** (7.9 mg), (–)-licarin A (55.8 mg), and kachirachirol B (56.4 mg).

Odoratisol A (**1**): Oil, $[\alpha]_{\text{D}}^{25} -35.1^\circ$ ($c=1.22$, CHCl_3). UV λ_{max} (MeOH) nm (log ϵ): 270 (4.18), 219 (4.43). IR ν_{max} (film) cm^{-1} : 3450, 2957, 2926, 2854, 1609, 1517, 1493, 1458, 1375, 1078. CD (MeOH): $\Delta\epsilon$ (nm): -2.14 (218), $+4.35$ (242), -5.93 (269) ($c=2.0\times 10^{-5}\text{ M}$). ^1H - and ^{13}C -NMR: see Tables 1 and 2. Negative-ion HR-FAB-MS: m/z 355.1544 $[\text{M}-\text{H}]^-$ (Calcd for $\text{C}_{21}\text{H}_{23}\text{O}_5$: 355.1545).

Odoratisol B (**2**): Oil, $[\alpha]_{\text{D}}^{25} +18.6^\circ$ ($c=0.70$, CHCl_3). UV λ_{max} (MeOH) nm (log ϵ): 258 (3.68), 218 (3.87). IR ν_{max} (film) cm^{-1} : 3448, 2957, 2925, 2854, 1603, 1511, 1458, 1377, 1061. CD (MeOH): $\Delta\epsilon$ (nm): -0.74 (219), $+0.38$ (258), $+0.18$ (350) ($c=4.2\times 10^{-5}\text{ M}$). ^1H - and ^{13}C -NMR: see Tables 1 and 2. Negative-ion HR-FAB-MS: m/z 343.1523 $[\text{M}-\text{H}]^-$ (Calcd for $\text{C}_{20}\text{H}_{23}\text{O}_5$: 343.1545).

Odoratisol C (**3**): Oil, $[\alpha]_{\text{D}}^{25} -26.0^\circ$ ($c=2.79$, CHCl_3). UV λ_{max} (MeOH) nm (log ϵ): 280 (3.71), 232 (4.05). IR ν_{max} (film) cm^{-1} : 3450, 2958, 2926, 2873, 1608, 1517, 1488, 1442, 1377, 1036. CD (MeOH): $\Delta\epsilon$ (nm): -1.25 (213), $+0.95$ (240), -0.40 (285) ($c=8.8\times 10^{-5}\text{ M}$). ^1H - and ^{13}C -NMR: see Tables 1 and 2. Negative-ion HR-FAB-MS: m/z 343.1542 $[\text{M}-\text{H}]^-$ (Calcd for $\text{C}_{20}\text{H}_{23}\text{O}_5$: 343.1545).

Odoratisol D (**4**): Oil, $[\alpha]_{\text{D}}^{25} -12.8^\circ$ ($c=4.0$, CHCl_3). UV λ_{max} (MeOH) nm (log ϵ): 282 (3.80), 234 (3.97). IR ν_{max} (film) cm^{-1} : 3450, 2959, 2927, 2873, 1608, 1517, 1488, 1442, 1377, 1036. CD (MeOH): $\Delta\epsilon$ (nm): -1.25 (213), $+0.95$ (240), -0.40 (285) ($c=8.8\times 10^{-5}\text{ M}$). ^1H - and ^{13}C -NMR: see Tables 1 and 2. Negative-ion HR-FAB-MS: m/z 341.1401 $[\text{M}-\text{H}]^-$ (Calcd for $\text{C}_{20}\text{H}_{21}\text{O}_5$: 341.1389).

Acknowledgments This work was supported by a Grant-in-Aid from the Japan Society for the Promotion of Science (JSPS). P.M.G. is grateful to acknowledge the JSPS for a Postdoctoral Research Fellowship at Hiroshima University and the International Foundation for Science (Stockholm, Sweden) for a research grant. We thank the Research Center of the Graduate School of Biomedical Sciences, Hiroshima University, Japan, for the meas-

urements on its 400 MHz NMR instrument.

References

- 1) Pham H. H., "Illustrated Flora of Vietnam," Tom 1, Fascile 1, Published by the Author, Montreal, 1991, pp. 488—492.
- 2) Le K. K., "Common Flora of Vietnam," Vol. 2, Science and Technique, Hanoi, 1971, pp. 265—266.
- 3) Do T. L., "Medicinal Plants and Remedies of Vietnam," Science and Technique, Hanoi, 1995, pp. 666—668.
- 4) Nguyen D. M., "Anti-infective Drugs from Plants of Vietnam," Medicine, Ho Chi Minh city, 1995, pp. 105—109.
- 5) Aiba C. J., Campos Corrêa R. G., Gottlieb O. R., *Phytochemistry*, **12**, 1163—1164 (1973).
- 6) Nascimento I. R., Lopes L. M. X., Davin L. B., Lewis N. G., *Tetrahedron*, **56**, 9181—9193 (2000).
- 7) Ito K., Ichino K., Iida J., Lai J., *Phytochemistry*, **23**, 2643—2645 (1984).
- 8) Tsai I. L., Hsieh C. F., Duh C. Y., *Phytochemistry*, **48**, 1371—1375 (1998).
- 9) Shimomura H., Sashida Y., Oohara M., *Phytochemistry*, **27**, 634—636 (1988).
- 10) Benevides P. J. C., Sartorelli P., Kato M. J., *Phytochemistry*, **52**, 339—343 (1999).
- 11) Nascimento I. R., Lopes L. M. X., *Phytochemistry*, **52**, 345—350 (1999).
- 12) Forrest J. E., Heacock R. A., Forrest T. P., *J. Chem. Soc. Perkin Trans. I*, **1974**, 205—209 (1974).
- 13) Shimomura H., Sashida Y., Oohara M., *Phytochemistry*, **26**, 1513—1515 (1987).
- 14) Matsuda N., Kikuchi M., *Chem. Pharm. Bull.*, **44**, 1676—1679 (1996).
- 15) Hattori M., Hada S., Kawata Y., Tezuka Y., Kikuchi T., Namba T., *Chem. Pharm. Bull.*, **35**, 3315—3322 (1987).
- 16) Kraft C., Jenett-Siems K., Köhler I., Tofern-Reblin B., Siems K., Bienzle U., Eich E., *Phytochemistry*, **60**, 167—173 (2002).
- 17) Urzúa A., Freyer A. J., Shamma M., *Phytochemistry*, **26**, 1509—1511 (1987).
- 18) Konishi T., Konoshima T., Daikonya A., Kitanaka S., *Chem. Pharm. Bull.*, **53**, 121—124 (2005).
- 19) Hada S., Hattori M., Tezuka Y., Kikuchi T., Namba T., *Phytochemistry*, **27**, 563—568 (1988).