A Novel Cyclopentene Derivative and a Polyhydroxylated Steroid from a South China Sea Gorgonian *Menella* sp.

Xing-Yun CHAI,^a Jian-Fan SUN,^a Li-Ying TANG,^b Xian-Wen YANG,^a Yun-Qiu LI,^a Hui HUANG,^{a,c}
Xue-Feng ZHOU,^a Bin YANG,^a and Yonghong LIU^{*,a}

^aKey Laboratory of Marine Bio-resources Sustainable Utilization/Guangdong Key Laboratory of Marine Materia Medica/Research Center for Marine Microbes, South China Sea Institute of Oceanology, Chinese Academy of Sciences; Guangzhou 510301, China; ^bInstitute of Traditional Chinese Materia Medica, Chinese Academy of Medical Sciences; Beijing 100700, P.R. China; and ^cNational Experiment Station of Tropical Marine Biology; Sanya 572000, China.

Received February 5, 2010; accepted July 13, 2010; published online July 29, 2010

A chemical investigation on a South China Sea gorgonian, *Menella* sp. resulted in the isolation and elucidation of menellin A (**1**), a highly oxygenated racemate with C₈ skeleton, and a polyhydroxylated steroid, menellsteroid C (**2**), along with eight known compounds (**3**–**10**). The structures of the new compounds were elucidated by means of MS, 1D and 2D NMR spectra, and the relative stereochemistry of **1** was determined by X-ray single-crystal diffraction analysis. In addition, compound **7** was isolated as a new natural product. Compounds **1**–**3** and **7** were selected to test the anti-inflammatory inhibition against lipopolysaccharide (LPS)-induced nitric oxide (NO) production in RAW264.7 macrophages. **1** and **3** exhibited modest inhibitory effects with IC₅₀ of 71.3, 33.9 μM, respectively, compared to the positive control aminoguanidine (IC₅₀ 25.0 μM).

Key words *Menella* sp.; gorgonian; menellin A; menellsteroid C; X-ray diffraction; anti-inflammatory inhibition

Gorgonians, with trivial names of Sea Fan or Sea Whip *etc.*, are one of the important sources recognized the new and significant biological activity of metabolites and are arousing the interest of chemists all over the world. There are a large number of papers every year on secondary metabolite research of the gorgonians.^{1–4)} However, one of the prolific gorgonian in the South China Sea, the *Menella* sp., has been given comparatively less attention, with the exception of four papers that described the guaiane sesquiterpenoids, polyhydroxylated steroids and other known compounds.^{5–8)}

As a part of our systematic research on the structurally novel and significantly bioactive metabolites from South China Sea invertebrates, we undertook a chemical investigation on an unidentified gorgonian species *Menella* sp., collected on Hainan Island. This resulted in the isolation and structure elucidation of a novel constituent, we named menellin A (**1**), and a polyhydroxylated steroid menellsteroid C (**2**), together with eight known compounds (**3**–**10**), including a new natural product (**7**). The structures of these new compounds were elucidated by spectrum analysis of MS, 1D and 2D NMR. The relative stereochemistry of **1** was determined by X-ray single-crystal diffraction analysis.

Results and Discussion

Menellin A (**1**) was obtained as a colorless quadrate crystal (CHCl₃–MeOH). Its molecular formula was determined as C₁₁H₁₄O₈ by an [M–H][–] ion peak at *m/z* 273.0618 (Calcd 273.0616) in the high-resolution-electrospray ionization (HR-ESI)-MS, indicating 5 degrees of unsaturation. The IR spectrum exhibited the characteristic absorption for a hydroxyl (3485 cm^{–1}) and carboxyls [1736, 1706 (br) cm^{–1}]. The ¹H- and ¹³C-NMR spectra (Table 1) showed the signals for an olefin [δ_{H} 6.82 (t, 1.5); δ_{C} 142.4, 137.0], three methoxys [$\delta_{\text{H}}/\delta_{\text{C}}$ 3.78/53.1 (9), 3.45/59.6 (10), 3.77/52.5 (11)], three carboxyl carbons [δ 173.6 (C-8), 172.6 (C-7), 165.7 (C-6)], an oxygenated methine at δ 4.65/94.9, and a methine at δ 4.21/58.9, as well as an oxygenated quaternary carbon at δ 86.7. These data implied **1** to be a highly oxy-

genated compound with a single ring. These assignments and the planar structure were achieved by heteronuclear single quantum coherence (HSQC) and heteronuclear multiple bond coherence (HMBC) experiments. The HMBC correlations from two methoxys (OMe-9, 11) to two carboxyls (C-8, 6) respectively suggested the appearance of two methyl esters in the structure of **1**. An HMBC correlation from H-4 (δ 6.82) to C-6 established the connection of one methyl ester with the olefin. The HMBC correlations from additional methoxy proton to C-3, from H-3 to the quaternary olefin carbon (C-5), C-8 and C-2, as well as from H-1 to C-7 and C-8 showed the planar structure of **1** to be a derivative of cyclopentene with three carboxyl groups attached at the ring (Fig. 1).

The relative stereochemistry of **1** was finally determined by X-ray single-crystal diffraction analysis after the failure by nuclear Overhauser effect spectroscopy (NOESY) spectrum, where no useful NOESY correlation was observed between the H-1 with H-3, or between OMe-9 with OMe-10. However, the X-ray diffraction result showed that the **1** is a natural racemate in consideration of its space group (P-1), as P-1 is a racemic space group, one with 1*S*, 2*S*, 3*S* configurations (**1**) and another with all *R* configurations. This conclusion was further confirmed by the almost zero optical rota-

Table 1. The NMR Data of Compound **1**

No.	δ_{C}	δ_{H}
1	58.9	4.21 d (1.5)
2	86.7	—
3	94.9	4.65 br s
4	142.4	6.82 t (1.5)
5	137.0	—
6	165.7	—
7	172.6	—
8	173.6	—
9	53.1	3.78
10	59.6	3.45
11	52.5	3.77

* To whom correspondence should be addressed. e-mail: yonghongliu@scsio.ac.cn.

Fig. 1. Structures of Compounds from *Menella* sp. (1–10) and (+)-Kjellmanone (11)

tion value.

Menellsteroid C (2) was obtained as white powder. Its molecular formula was determined as $C_{27}H_{48}O_4$ by its ESI-MS ion $[M+Na]^+$ at m/z 459 and NMR data (Table 2). The 1H - and ^{13}C -NMR spectra showed characteristic signals of five methyls, including two tertiary ones (C-18, 19) and three secondary methyls at δ_H/δ_C 0.99/19.2 (C-21), 0.89/22.9 (C-26) and 0.89/23.2 (C-27), which in collaboration with its typical C-3 oxygenated proton signal at δ_H 4.02 (m, H-3) and one at δ_H 3.46 (brs, H-6) due to both C-5 and C-6 hydroxylated in its 1H -NMR, suggested 2 a polyhydroxylated cholestan-type steroid. A detailed comparison of its NMR data with those of menellsteroid A (3)^{7,9)} showed both are isomers with high structural similarity (Fig. 1), which was supported and confirmed by the HMBC spectra (Fig. 2). The differences be-

Fig. 2. Key HMBC and NOESY Correlations of 1 and 2

Table 2. NMR Data (CD_3OD , 500/125 MHz) for 2, 3 and 7, δ in ppm and J in Hz

No.	2		3	7	
	δ_H	δ_C	δ_C	δ_H	δ_C
1	2.07 t (11.5), 1.79 m	35.4	33.2	4.07 dd (10.5, 5.5)	71.7
2	1.78, 1.53 each m	32.0	31.6	2.05, 2.15 each m	42.7
3	4.02 m	68.2	68.3	3.95 m	65.6
4	1.57 m, 2.09 br d (11.5)	42.0	41.1	1.57 m, 1.80 br d (11.5)	36.7
5	—	77.5	77.5	—	82.2
6	3.46 br s	76.6	76.4	—	214.8
7	1.79, 1.53 m	35.2	36.4	2.79 t (12.5), 2.15 m	41.6
8	1.78 m	30.4	28.4	1.69 m	41.5
9	1.46 m	53.0	49.5	1.47 m	46.6
10	—	41.0	40.0	—	46.7
11	3.85 dt (5.0, 10.5)	69.5	69.5	1.86, 1.27 each m	23.2
12	1.24 m, 2.28 dd (12.5, 5.0)	52.8	50.5	1.99, 1.32 each m	40.7
13	—	44.3	43.1	—	43.6
14	1.19 m	57.6	59.3	1.17 m	58.0
15	1.39 br d (11.5), 1.18 m	25.0	25.0	1.40 d (11.5), 1.18 m	25.1
16	1.54, 1.30 each m	29.4	29.1	1.55, 1.30 each m	29.2
17	1.18 m	56.6	58.3	1.18 m	58.0
18	0.74 s	13.5	14.9	0.71 s	12.5
19	1.29 s	17.6	20.3	0.78 s	9.5
20	1.40 m	37.1	37.3	1.40 m	37.1
21	0.99 d (6.5)	19.2	19.3	0.98 d (6.5)	19.2
22	1.05 m	37.3	37.3	1.05 m	36.7
23	1.60 m	25.3	25.3	1.61 m	25.3
24	1.17 m	40.7	40.7	1.18 m	40.7
25	1.54 m	29.2	29.2	1.54 m	29.2
26	0.89 d (6.5)	22.9	23.0	0.89 d (6.5)	23.0
27	0.89 d (6.5)	23.2	23.2	0.90 d (6.5)	23.2

Fig. 3. Single-Crystal X-Ray Structure of **1**

tween them are the downfield chemical shift of 3.5 ppm C-9, and 2.3 ppm for C-12, and the upfield shift to δ 3.85 (H-11) compared to that of δ 4.14 in **3**, suggesting that **2** differs from **3** only at the probable stereochemistry of C₁₁-OH. This conclusion was supported by a comparison of their ¹³C-NMR and confirmed by the NOESY correlation between H-11 with H-19. Also, the H-11 β configuration could be deduced by its coupling pattern (dt, 5.0, 10.5 Hz) according to the literature.⁷⁾ Thus, compound **2** was determined to be cholestan-3 β ,5 α ,6 β ,11 α -tetrol, and called menellsteroid C.

Compound **7** was obtained as a white solid, with its molecular formula of C₂₇H₄₆O₄ by its ESI-MS. The NMR data (Table 2) showed the typical signals of a polyoxygenated cholestan-type steroid: two tertiary methyls, three secondary methyls, and three oxygenated carbons. Its NMR data was closely similar to that of compound **5**,¹⁰⁾ including the coupling pattern of H-1 (4.07, dd, 5.0, 11.5 Hz), except for the appearance of an additional ketone signal at δ 214.8 in **7**, and the disappearance of the characteristic proton signal around δ 3.50, which suggests the hydroxyl group at C-6 in compound **5** was oxygenated to a ketone in **7**. All the NMR data was assigned by HSQC spectrum and was in agreement with references.^{10,11)} Thus, **7** was identified as 1 β ,3 β ,5 α -trihydroxycholestan-6-one. Since it was isolated as a natural product for the first time, we propose its trivial name menellsteroid D.

The other seven known compounds were identified as menellsteroid A (**3**),^{7,9)} cholestan-3 β ,5 α ,6 β -triol (**4**),⁹⁾ cholestan-1 β ,3 β ,5 α ,6 β -tetrol (**5**),¹⁰⁾ nephalsterol (**6**),¹²⁾ and cholestan-3 β -en-6-one (**8**),¹³⁾ and two briarane diterpenoids juncecellolides B (**9**) and D (**10**),¹⁴⁾ by comparison of their NMR data with those of references.

Compound **1** is of special interest, as it is the first example with a highly oxygenated cyclopentene ring discovered from a gorgonian. Furthermore, **1** was a rarely reported and naturally occurring racemate. It was determined by the X-ray diffraction with a P-1 space group, and confirmed by its optical rotation value. It is structurally similar to (+)-kjellmanone (**11**),^{15,16)} a cyclopentenone derivative from the marine alga *Sargassum kjellmanianum*. Compound **7** was isolated as a natural product for the first time, and compounds **9** and **10** were the first two briarane diterpenoids found from *Menella* sp. to date.

Compounds **1**–**3** and **7** were chosen to test the anti-inflammatory against lipopolysaccharide (LPS)-induced nitric oxide (NO) production in RAW264.7 macrophages. The results showed that **1** and **3** exhibited modest inhibition effects

with IC₅₀ of 71.3 and 33.9 μ M, respectively, compared to the positive control aminoguanidine with IC₅₀ 25.0 μ M. In addition, compounds **1**–**10** were evaluated by the anti-oxidant capacity in a radical 2, 2-diphenyl-(2,4,6-trinitrophenyl)hydrazyl (DPPH) free-radical assay, but no obvious scavenging effect was observed.

Experimental

General Experimental Procedures Melting points were determined on an X-5 digital micro melting point apparatus (Beijing Tech Instrument Co., Ltd., China) and are uncorrected. Optical rotations were recorded on a JASCO P-1020 digital polarimeter. IR spectra were measured on JASCO FT/IR-480 plus spectrometers. NMR spectra were recorded using Bruker 500 MHz NMR spectrometers. HR-ESI-MS spectra were recorded on an Applied Biosystems Mariner 5140 spectrometer. The column chromatographies were applied on the Buchi Sepacore (C-615/605) system. All solvents used were of analytical grade (Tianjing Fuyu Chemical and Industry Factory, China). Silica gel and preparative TLC plates (20 $\times$ 20 $\times$ 0.04 cm) (Qingdao Mar. Chem. Ind. Co., Ltd., China), Sephadex LH-20 gel (Pharmacia, Sweden) and C₁₈ reverse-phased silica gel (150–200 mesh, Merck, Germany) were used for column chromatography.

Animal Material Fresh gorgonians were collected in April 2009 (7–10 m depth) off Meishan Island, Hainan province, China. The specimen was identified by Professor Hui Huang, the South China Sea Institute of Oceanology, Academia Sinica. A voucher specimen (No. M090402) was deposited in the Key Laboratory of Marine Bio-resources Sustainable Utilization, South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou, P.R. China.

Extraction and Isolation The fresh gorgonian (ca. 5 kg) were exhaustively extracted with 95%, 70% EtOH and CHCl₃-MeOH (1:1) at room temperature, and the solvent was evaporated *in vacuo*. The residue was partitioned in H₂O and extracted with petroleum ether (PE), EtOAc, and *n*-butanol, respectively, to provide the PE fraction (16 g), EtOAc fraction (9.5 g) and the *n*-butanol fraction (11 g). The EtOAc fraction was chromatographed on silica gel and eluted with PE-acetone (50:1), followed by a gradient of CHCl₃-MeOH (100:0, 30:1, 10:1, 1:1) to yield 8 fractions (Frs. A–G). Fr. B (2.1 g) was re-separated by Sephadex LH-20 (MeOH), followed by silica gel column chromatography (CC) and eluted with PE-acetone (5:3) to get 6 subfractions (Fr. Ba–f). A further isolation of Fr. Bc yielded **10** (4.6 mg) by preparative thin layer chromatography (p-TLC, two plates) with PE-EtOAc (1:1) as developer. The p-TLC (PE-acetone 5:4, one plate) of Fr. Bb provided **9** (1.8 mg). Fr. E (800 mg) was chromatographed by silica gel CC and eluted with PE-EtOAc (1:3), followed by a recrystallization (CHCl₃-MeOH 1:1) to obtain **1** (13.7 mg).

The PE fraction was chromatographed on silica gel and eluted with a gradient of PE-EtOAc (100:0, 50:1, 10:1, 2:1, 0:1) to yield 14 fractions (Frs. 1–14). Fr. 12 (2.5 g) was subjected to silica gel CC with elution by PE-EtOAc (3:1) to get 4 fractions (Frs. 12A–D), and twice to p-TLC of Fr. 12C (200 mg) (CHCl₃-acetone 10:1 and PE-EtOAc 1:1 as developer, respectively) which yielded **8** (2.2 mg). Fr. 14 (4 g) was resubjected to silica gel CC and eluted with a gradient of PE-EtOAc (3:1–1:4) to 9 fractions (Frs. 14A–I). A purification by Sephadex LH-20 (MeOH) to Fr. 14E (420 mg) produced **4** (32.0 mg) and 4 other portions (Frs. 14E₁–E₄). Frs. 14E₃ and E₄ were subjected to p-TLC (CHCl₃-EtOAc 1:5 and PE-acetone 2:3, respectively) to obtain **7** (3.9 mg) (from Fr. 14E₃), and **3** (16.1 mg) and **6** (6.2 mg). Fr. 14C (600 mg) was applied to silica gel CC and eluted with

CHCl₃–MeOH (10 : 1) to yield 7 subfractions (1–7). Fr. 14C₆ (40 mg) was applied to p-TLC (3 plates) and developed by CHCl₃–EtOH (5 : 1) to get **2** (4.2 mg) and **5** (8.5 mg).

Menellin A (**1**): Colorless quadrate crystal (CHCl₃–MeOH); mp 173.2–175.1 °C; $[\alpha]_D^{25} +0.0002$ ($c=0.13$, MeOH); UV $\lambda_{\max}$ (MeOH) nm (log ϵ): 222 (3.98) nm; IR (KBr) $\nu_{\max}$ cm⁻¹: 3485, 3010, 2955, 1736, 1706 (br), 1645, 1436, 1258, 1146, 960; ¹H- and ¹³C-NMR (CD₃OD, 500/125 MHz), see Table 1. ESI-MS m/z 275 [M+H]⁺; HR-ESI-MS m/z 273.0618 [M–H][–] (Calcd for C₁₁H₁₃O₈; 273.0616).

Menellsteroid B (**2**): White powder; $[\alpha]_D^{25} -18.5$ ($c=0.12$, MeOH); IR (KBr) $\nu_{\max}$ cm⁻¹: 3480, 2905, 2871, 1458, 1376, 1200; ¹H- and ¹³C-NMR (CD₃OD, 500/125 MHz), see Table 2. ESI-MS m/z 459 [M+Na]⁺; HR-ESI-MS m/z 437.3552 [M+H]⁺ (Calcd for C₂₇H₄₉O₄; 437.3553).

1 β ,3 β ,5 α -Trihydroxycholestan-6-one (Proposed Menellsteroid D) (**7**): White solid; ¹H- and ¹³C-NMR (CD₃OD, 500/125 MHz), see Table 2. ESI-MS m/z 457 [M+Na]⁺.

X-Ray Crystallographic Analysis of Menellin A (1) C₁₁H₁₄O₈, $M=274.22$, triclinic, space group $P-1$, $a=7.9389(5)$ Å, $b=9.1825(6)$ Å, $c=9.7595(6)$ Å. $V=625.73(7)$ Å³, $Z=2$, $D_{\text{calcd}}=1.455$ Mg/m³, crystal dimensions $0.46\times0.43\times0.42$ mm³, $\mu=0.126$ mm⁻¹. 5216 Reflections measured, 2659 reflections independent ($R_{\text{int}}=0.0154$), $R=0.0362$, $R_w=0.1068$, X-ray diffraction experiments for this compound were carried out on a Bruker Smart 1000 CCD diffractometer at 273 K using MoK α radiation ($\lambda=0.71070$ Å). Absorption corrections were done by semi-empirical from equivalents. The structure was resolved using direct methods and refined with full-matrix least-squares on F^2 . Crystallographic data for the structure of **1** have been deposited in the Cambridge Crystallographic Data Centre (deposition number CCDC 754564). Copies of the data can be obtained free of charge via www.ccdc.cam.ac.uk/conts/retrieving.html [or from the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB21EZ, U.K.; fax: t44 1223 336 033 or deposit@ccdc.cam.ac.uk].

Biological Assay The procedure reported by Yang *et al.*¹⁷⁾ was used for the measurement of anti-inflammatory inhibition against LPS-induced NO production in RAW 246.7 in macrophages, and the DPPH free-radical scavenging inhibitory activity was measured as described by Lee *et al.*¹⁸⁾

Acknowledgements This study was supported by Grants from the National Natural Science Foundation of China (Grant numbers: 40706046, 30973679, and 20902094), the National Key Basic Research Program of China (973)'s Project (2010CB833800), Knowledge Innovation Program of the Chinese Academy of Science (LYQY200703, SQ200904, and KSCX2-

YW-G-073), the Scientific Research Foundation for the Returned Overseas Chinese Scholars, State Education Ministry, and LMB (091002) Foundation. Our thanks are due to Mr. Jiang-Hui Tang for his assistance in collection of the gorgonian, Mr. Xiao-Long Feng (Sun Yat-Sen University) for the analysis of single-crystal X-ray diffraction, and Dr. Xiao-Qi Zhang (Jinan University) for his kind help in the measurement of physical data.

References

- 1) Zhang W., Guo Y. W., *Chin. J. Nat. Med.*, **1**, 69–75 (2003).
- 2) Berrue F., Kerr R. G., *Nat. Prod. Rep.*, **26**, 681–718 (2009).
- 3) Blunt J. W., Copp B. R., Hu W. P., Munro M. H. G., Northcote P. T., Prinsep M. R., *Nat. Prod. Rep.*, **25**, 35–94 (2008).
- 4) Blunt J. W., Copp B. R., Hu W. P., Munro M. H. G., Northcote P. T., Prinsep M. R., *Nat. Prod. Rep.*, **26**, 170–244 (2009).
- 5) Deng S. Z., Li F. Y., Peng S. S., Rao Z. G., Wu H. M., Xu J. H., *Chin. J. Appl. Chem.*, **14**, 80–82 (1997).
- 6) Zhang W., Guo Y. W., Mollo E., Cimino G., *Helv. Chim. Acta*, **87**, 2919–2925 (2004).
- 7) Zhang W., Huang H., Ding Y., Gavagnin M., Mollo E., Cimino G., Guo Y. W., *Helv. Chim. Acta*, **89**, 813–820 (2006).
- 8) Li L., Wang C. Y., Huang H., Mollo E., Cimino G., Guo Y. W., *Helv. Chim. Acta*, **91**, 111–117 (2008).
- 9) Qiu Y. Q., Qi S. H., Zhang S., Yang J., Xiao Z. H., *Pharmazie*, **61**, 645–647 (2006).
- 10) Kobayashi M., Hayashi T., Hayashi K., Tanabe M., Nakagawa T., Mitsuhashi H., *Chem. Pharm. Bull.*, **31**, 1848–1855 (1983).
- 11) Endo Y., Fukasawa H., Hashimoto Y., Shudo K., *Chem. Pharm. Bull.*, **42**, 462–469 (1994).
- 12) Wang G. Y. S., Li F. Y., Zeng L. M., *Chem. J. Chin. Univ.*, **13**, 623–627 (1992).
- 13) Zhan Y. C., Chen L., Sun Y., Zhang N., Pei Y. H., *J. Shenyang Pharm. Univ.*, **23**, 358–360 (2006).
- 14) Shin J., Park M., Fenical W., *Tetrahedron*, **45**, 1633–1638 (1989).
- 15) Nakayama M., Fukuoka Y., Nozaki H., Matsuo A., Hayashi S., *Chem. Lett.*, **1980**, 1243–1246 (1980).
- 16) Qin M. L., Li X. M., Yin S. W., Wang C. Y., Wang B. G., *Marine Sci.*, **31**, 47–50 (2007).
- 17) Yang Y. W., Zeng H. W., Liu X. H., Li S. M., Xu W., Sheng Y. H., Zhang C., Zhang W. D., *J. Pharm. Pharmacol.*, **60**, 937–941 (2008).
- 18) Lee Y. Y., Jang D. S., Jin J. L., Yun-Choi H. S., *Planta Med.*, **71**, 776–777 (2005).