Bioactive Constituents from Chinese Natural Medicines. XXXVI.¹⁾ Four New Acylated Phenylethanoid Oligoglycosides, Kankanosides J₁, J₂, K₁, and K₂, from Stems of *Cistanche tubulosa*

Yingni PAN,^{a,b} Toshio MORIKAWA,^a Kiyofumi NINOMIYA,^a Katsuya IMURA,^a Dan YUAN,^b Masayuki YOSHIKAWA,^{*,a,c} and Osamu MURAOKA^{*,a}

^a Pharmaceutical Research and Technology Institute, Kinki University; 3–4–1 Kowakae, Higashi-osaka, Osaka 577–8502, Japan; ^b School of Traditional Chinese Medicine, Shenyang Pharmaceutical University; 103 Wenhua Rd., Shenyang 110016, People's Republic of China; and ^c Kyoto Pharmaceutical University; Misasagi, Yamashina-ku, Kyoto 607–8412, Japan. Received November 28, 2009; accepted February 2, 2010; published online February 4, 2010

Four new acylated phenylethanoid oligoglycosides, kankanosides J₁ (1), J₂ (2), K₁ (3), and K₂ (4), were isolated from stems of *Cistanche tubulosa* (Orobanchaceae) together with isocampneoside I (5). Their structures were elucidated on the basis of chemical and physicochemical evidence. Among them, 3–5 were found to inhibit D-galactosamine-induced cytotoxicity in primary cultured mouse hepatocytes.

Key words *Cistanche tubulosa*; kankanoside; phenylethanoid glycoside; Orobanchaceae; hepatoprotective activity

During the course of our studies on bioactive constituents from Chinese natural medicines,^{1–3)} we found that methanolic extract of dried stems of *Cistanche tubulosa* (SCHRENK) R. WIGHT (Orobanchaceae) showed vasorelaxant⁴⁾ and hepatoprotective activities.¹⁾ From the dried stems of *C. tubulosa*, five iridoids, kankanosides A–D and kankanol, a monoterpene glycoside, kankanoside E, two phenylethanoid oligoglycosides, kankanosides F and G, and an acylated oligosugar, kankanose, were isolated together with 30 known constituents.^{4,5)} Recently, we additionally isolated 19 phenylethanoid oligoglycosides including kankanosides H₁, H₂, and I¹⁾ and two acylated oligosugars from fresh stems of *C. tubulosa*.¹⁾ Furthermore, principal phenylethanoid glycosides, echinacoside, acteoside, and isoacteoside, were found to inhibit increase in serum aspartate aminotransferase (sAST) and alanine aminotransferase (sALT) levels in liver injured mice induced by D-galactosamine (D-GalN)/lipopolysaccharide at doses of 25–100 mg/kg *per os* (*p.o.*), and structural requirements of phenylethanoid glycosides for the hepatoprotective activity were elucidated.¹⁾ As a continuing study on constituents from the fresh stems of *C. tubulosa*, we further isolated four new acylated phenylethanoid oligoglycosides, kankanosides J₁ (1), J₂ (2), K₁ (3), and K₂ (4). This paper deals with isolation and structure elucidation of 1–4.

Fresh stems of *C. tubulosa* (cultivated in Urumuqi, Xinjiang Province, China) were extracted with methanol under reflux to yield a methanolic extract (8.36% from the fresh stems). From the methanolic extract, H₂O- and MeOH-eluted fractions (5.63% and 2.73%, respectively) were obtained by Diaion HP-20 column chromatography (H₂O→MeOH) as was described previously.¹⁾ By the intensive chromatographies on the MeOH-eluted fraction, four new phenylethanoid oligoglycosides, kankanosides J₁ (1, 0.0002%), J₂ (2, 0.0002%), K₁ (3, 0.0002%), and K₂ (4, 0.0005%) together with isocampneoside I⁶⁾ (5, 0.0006%) were isolated.

Structures of Kankanosides J₁ (1) and J₂ (2) Kankanoside J₁ (1) was obtained as a white powder with negative optical rotation ([α]_D²⁵ –6.5° in MeOH). The IR spectrum of 1 showed absorption bands at 3414, 1734, 1719, 1701, 1638, 1508, 1159, 1067, and 1046 cm^{–1} ascribable to hydroxyls, ester carbonyls, ether functions, and aromatic

rings. The positive- and negative-ion FAB-MS spectra of 1 showed quasimolecular ion peaks at *m/z* 719 (M+Na)⁺ and *m/z* 695 (M–H)[–], and the molecular formula was determined as C₃₂H₄₀O₁₇ by high-resolution positive-ion FAB-MS measurement. The ¹H- and ¹³C-NMR spectra of 1 (CD₃OD, Tables 1, 2), which were assigned by various NMR experiments,⁷⁾ showed signals assignable to a methoxy group [δ 3.21 (3H, s, 7-OCH₃)], a methylene and a methine bearing an oxygen function [δ 3.58, 4.00 (1H each, both m, 8-H₂)], 4.18 (1H, dd-like, *J*=ca. 4, 8 Hz, 7-H)], *ortho*- and *meta*-coupled ABC-type aromatic protons [δ 6.63 (1H, dd, *J*=1.8, 8.2 Hz, 6-H), 6.74 (1H, d, *J*=1.8 Hz, 2-H), 6.74 (1H, d, *J*=8.2 Hz, 5-H)], a β-D-glucopyranosyl moiety [δ 4.54 (1H, d, *J*=7.8 Hz, Glc-1-H)], and an α-L-rhamnopyranosyl moiety [δ 1.07 (3H, d, *J*=6.4 Hz, Rha-6-H₃), 4.80 (1H, br s, Rha-1-H)] together with an acetyl group [δ 2.00 (3H, s)] and a *trans*-caffeoyl group {an *trans*-olefin [δ 6.26, 7.59 (1H each, both d, *J*=16.0 Hz, 8-, 7-H)] and *ortho*- and *meta*-coupled ABC-type aromatic protons [δ 6.77 (1H, d, *J*=8.2 Hz, 5-H), 6.95 (1H, dd, *J*=1.8, 8.2 Hz, 6-H), 7.04 (1H, d, *J*=1.8 Hz, 2-H)]}. The ¹H- and ¹³C-NMR spectra of 1 were superimposable on those of campneoside I^{1,8–10)} (6), except for the signals due to the acetyl group. Connectivities of the oligoglycoside and acyl moieties in 1 were confirmed by the heteronuclear multiple bond correlation (HMBC) experiment, which showed long-range correlations between the following proton and carbon pairs: 7-OCH₃ and 7-C (δ_C 83.3); Glc-1-H and 8-C (δ_C

Chart 1

Table 1. ^1H -NMR Data (600 MHz, CD_3OD) for Kankanosides J_1 (**1**), J_2 (**2**), K_1 (**3**), and K_2 (**4**)

Position	1 δ_{H} (J Hz)	2 δ_{H} (J Hz)	3 δ_{H} (J Hz)	4 δ_{H} (J Hz)
2	6.74 (d, 1.8)	6.73 (br s)	6.80 (d, 2.0)	6.78 (d, 1.9)
5	6.74 (d, 8.2)	6.73 (d, 8.2)	6.76 (d, 8.1)	6.76 (d, 8.2)
6	6.63 (dd, 1.8, 8.2)	6.62 (dd, 1.8, 8.2)	6.68 (dd, 2.0, 8.1)	6.67 (dd, 1.9, 8.2)
7	4.18 (dd-like, ca. 4, 8)	4.22 (dd, 3.2, 8.2)	4.34 (dd, 3.4, 8.1)	4.37 (dd, 3.1, 9.1)
8	3.58 (m)	3.63 (m)	3.62 (dd, 3.4, 11.0)	3.72 (dd, 9.1, 11.0)
7-OCH ₃	4.00 (m)	3.83 (dd, 3.2, 11.0)	4.02 (dd, 8.1, 11.0)	3.84 (dd, 3.1, 11.0)
8-O-Glc	3.21 (3H, s)	3.24 (3H, s)	3.23 (3H, s)	3.25 (3H, s)
1'	4.54 (d, 7.8)	4.64 (d, 8.2)	4.40 (d, 7.9)	4.44 (d, 7.9)
2'	4.91 (dd, 7.8, 9.2)	4.91 (dd, 8.2, 9.2)	3.43 (dd, 7.9, 9.2)	3.43 (dd, 7.9, 9.1)
3'	4.01 (dd, 9.2, 9.6)	4.04 (dd, 9.2, 9.6)	3.81 (dd, 9.2, 9.7)	3.82 (dd, 9.1, 9.7)
4'	4.99 (dd, 9.2, 9.6)	5.00 (dd, 9.6, 9.6)	5.00 (dd, 9.7, 9.7)	5.02 (dd, 9.7, 9.8)
5'	3.56 (m)	3.61 (m)	3.77 (m)	3.77 (ddd, 2.2, 5.5, 9.8)
6'	3.51 (m)	3.52 (m)	3.64 (dd, 5.5, 11.6)	3.62 (dd, 5.5, 11.9)
	3.61 (m)	3.61 (m)	3.92 (dd, 2.0, 11.6)	3.94 (dd, 2.2, 11.9)
3'-O-Rha				
1''	4.80 (br s)	4.82 (br s)	5.19 (d, 1.7)	5.20 (d, 1.6)
2''	3.25 (m)	3.65 (m)	3.91 (dd, 1.7, 3.3)	3.92 (dd, 1.6, 3.2)
3''	3.61 (m)	3.52 (m)	3.57 (dd, 3.3, 9.5)	3.57 (dd, 3.2, 9.4)
4''	3.25 (m)	3.26 (m)	3.28 (m)	3.28 (m)
5''	3.55 (m)	3.52 (m)	3.55 (dq, 9.5, 6.4)	3.55 (dq, 9.1, 6.2)
6''	1.07 (3H, d, 6.4)	1.07 (3H, d, 6.4)	1.08 (3H, d, 6.4)	1.08 (3H, d, 6.2)
6'-O-Glc				
1'''			4.31 (d, 7.9)	4.26 (d, 7.7)
2'''			3.19 (dd, 7.9, 9.1)	3.19 (dd, 7.7, 9.3)
3'''			3.34 (dd, 8.8, 9.1)	3.32 (m)
4'''			3.26 (dd, 8.8, 9.6)	3.25 (m)
5'''			3.22 (m)	3.22 (m)
6'''			3.63 (dd, 5.7, 12.1)	3.62 (dd, 5.5, 12.0)
			3.83 (dd, 2.2, 12.1)	3.82 (dd, 2.1, 12.0)
2'-O-Ac	2.00 (3H, s)	2.11 (3H, s)		
4'-O- <i>trans</i> -Caf				
2	7.04 (d, 1.8)	7.04 (d, 1.8)	7.05 (d, 1.9)	7.05 (d, 1.9)
5	6.77 (d, 8.2)	6.77 (d, 8.2)	6.78 (d, 8.3)	6.78 (d, 8.4)
6	6.95 (dd, 1.8, 8.2)	6.95 (dd, 1.8, 8.2)	6.96 (dd, 1.9, 8.3)	6.96 (dd, 1.9, 8.4)
7	7.59 (d, 16.0)	7.54 (d, 16.0)	7.60 (d, 15.8)	7.60 (d, 15.8)
8	6.26 (d, 16.0)	6.27 (d, 16.0)	6.27 (d, 15.8)	6.28 (d, 15.8)

Fig. 1. ^1H - ^1H COSY and HMBC Correlations for **1**—**4**

74.1); Glc-2-H [δ 4.91 (1H, dd, $J=7.8$, 9.2 Hz)] and the acetyl carbonyl carbon (δ_{C} 171.4); Glc-4-H [δ 4.99 (1H, dd, $J=9.2$, 9.6 Hz)] and the *trans*-caffeoyl carbonyl carbon (δ_{C} 168.1); and Rha-1-H and Glc-3-C (δ_{C} 80.5) (Fig. 1). Finally, alkaline hydrolysis of **1** with 5% potassium hydroxide (KOH) liberated *trans*-caffeic acid, which was identified by HPLC analysis, together with a deacylated product. The deacylated product was successively treated with 1.0 M hydrochloric acid (HCl) to liberate L-rhamnose and D-glucose, which were identified by HPLC analysis using an optical rotation detector.^{1–5)} Thus, the structure of kankanoside J_1 was eluci-

dated to be 2-methoxy-2-(3,4-dihydroxyphenyl)ethyl *O*- α -L-rhamnopyranosyl-(1 $\rightarrow$ 3)-2-*O*-acetyl-4-*O*-*trans*-caffeoyl- β -D-glucopyranoside (**1**).

Kankanoside J_2 (**2**) was isolated as a white powder with negative optical rotation ($[\alpha]_{\text{D}}^{25} -18.1^\circ$ in MeOH). By high-resolution positive-ion FAB-MS measurement, the molecular formula of **2** was found to be the same as that of **1**. The ^1H - and ^{13}C -NMR data of **2** (CD_3OD , Tables 1, 2) were very similar to those of **1**, except for the signals due to the ethyl bridge of the aglycone moiety {a methoxy group [δ 3.24 (3H, s, 7-OCH₃)], a methylene [δ 3.63 (1H, m), 3.83 (1H, dd,

Table 2. ^{13}C -NMR Data (150 MHz, CD_3OD) for Kankanosides J_1 (**1**), J_2 (**2**), K_1 (**3**), and K_2 (**4**)

Position	1 δ_{C}	2 δ_{C}	3 δ_{C}	4 δ_{C}
1	131.7	131.2	131.2	130.7
2	115.3	115.0	115.2	115.1
3	146.5	146.5	146.5	146.5
4	146.4	146.3	146.4	146.5
5	116.3	116.2	116.3	116.4
6	120.2	119.7	120.2	119.9
7	83.3	84.3	83.5	84.4
8	74.1	74.9	74.7	75.2
7-OCH ₃	56.8	57.1	56.7	56.7
8-O-Glc				
1'	101.7	102.2	104.1	104.5
2'	75.1	75.3	76.0	76.3
3'	80.5	80.5	81.5	81.4
4'	70.7	70.7	70.6	70.5
5'	76.3	76.2	74.9	74.7
6'	62.3	62.2	69.4	69.5
3'-O-Rha				
1''	103.4	103.3	103.0	103.0
2''	72.7	72.6	72.4	72.4
3''	71.9	71.9	72.1	72.0
4''	73.7	73.6	73.8	73.8
5''	70.8	70.8	70.4	70.4
6''	18.5	18.5	18.4	18.4
6'-O-Glc				
1'''			104.6	104.7
2'''			75.1	75.1
3'''			77.8	77.8
4'''			71.5	71.5
5'''			77.9	77.9
6'''			62.6	62.6
2'-O-Ac				
1	20.9	21.0		
2	171.4	171.5		
4'-O-Caf				
1	127.7	127.6	127.6	127.6
2	115.3	115.2	115.3	115.3
3	146.9	146.5	146.9	146.8
4	149.9	149.9	149.9	149.9
5	116.6	116.5	116.5	116.5
6	123.3	123.2	123.3	123.3
7	148.2	148.2	148.2	148.3
8	114.6	114.6	114.7	114.7
9	168.1	168.1	168.5	168.5

$J=3.2$, 11.0 Hz), 8-H₂] and a methine bearing an oxygen function [δ 4.22 (1H, dd, $J=3.2$, 8.2 Hz, 7-H)]. Alkaline hydrolysis of **2** with 5% KOH liberated *trans*-caffeic acid together with a deacylated product, and the deacylated product was successively treated with 1.0 M HCl to liberate L-rhamnose and D-glucose. As shown in Fig. 1, the same long-range correlations as in the case of **1** were observed in the HMBC experiment. Consequently, the planar structure of kankanoside J_2 (**2**) was revealed to be the same as that of **1**, and was elucidated to be 7-isomer of **1**.¹¹⁾

Structures of Kankanosides K_1 (3**) and K_2 (**4**)**
Kankanosides K_1 (**3**) and K_2 (**4**), $\text{C}_{36}\text{H}_{48}\text{O}_{21}$, were also obtained as white powders with negative optical rotations (**3**: $[\alpha]_{\text{D}}^{25} -75.3^\circ$; **4**: $[\alpha]_{\text{D}}^{25} -7.4^\circ$ both in MeOH). The ^1H - and ^{13}C -NMR spectra of **3** and **4** (CD_3OD , Tables 1, 2) showed signals assignable to a methoxy group [**3**: δ 3.23 (3H, s, 7-OCH₃); **4**: δ 3.25 (3H, s, 7-OCH₃)], a methylene and a methine bearing an oxygen function [**3**: δ [3.62 (1H, dd, $J=3.4$,

11.0 Hz), 4.02 (1H, dd, $J=8.1$, 11.0 Hz), 8-H₂], 4.34 (1H, dd, $J=3.4$, 8.1 Hz, 7-H); **4**: δ [3.72 (1H, dd, $J=9.1$, 11.0 Hz), 3.84 (1H, dd, $J=3.1$, 11.0 Hz), 8-H₂], 4.37 (1H, dd, $J=3.1$, 9.1 Hz, 7-H)], *ortho*- and *meta*-coupled ABC-type aromatic protons [**3**: δ 6.68 (1H, dd, $J=2.0$, 8.1 Hz, 6-H), 6.76 (1H, d, $J=8.1$ Hz, 5-H), 6.80 (1H, d, $J=2.0$ Hz, 2-H); **4**: δ 6.67 (1H, dd, $J=1.9$, 8.2 Hz, 6-H), 6.76 (1H, d, $J=8.2$ Hz, 5-H), 6.78 (1H, d, $J=1.9$ Hz, 2-H)], two β -D-glucopyranosyl moieties [**3**: δ 4.31 (1H, d, $J=7.9$ Hz, *terminal*-Glc-1-H), 4.40 (1H, d, $J=7.9$ Hz, *inner*-Glc-1-H); **4**: δ 4.26 (1H, d, $J=7.7$ Hz, *terminal*-Glc-1-H), 4.44 (1H, d, $J=7.9$ Hz, *inner*-Glc-1-H)], and an α -L-rhamnopyranosyl moiety [**3**: δ 1.08 (3H, d, $J=6.4$ Hz, Rha-6-H₃), 5.19 (1H, d, $J=1.7$ Hz, Rha-1-H); **4**: δ 1.08 (3H, d, $J=6.2$ Hz, Rha-6-H₃), 5.20 (1H, d, $J=1.6$ Hz, Rha-1-H)] together with a *trans*-caffeoyl group {an *trans*-olefin [**3**: δ 6.27, 7.60 (1H each, both d, $J=15.8$ Hz, 8-, 7-H); **4**: δ 6.28, 7.60 (1H each, both d, $J=15.8$ Hz, 8-, 7-H)] and *ortho*- and *meta*-coupled ABC-type aromatic protons [**3**: δ 6.78 (1H, d, $J=8.3$ Hz, 5-H), 6.96 (1H, dd, $J=1.9$, 8.3 Hz, 6-H), 7.05 (1H, d, $J=1.9$ Hz, 2-H)]; **4**: δ 6.78 (1H, d, $J=8.4$ Hz, 5-H), 6.96 (1H, dd, $J=1.9$, 8.4 Hz, 6-H), 7.05 (1H, d, $J=1.9$ Hz, 2-H)]}. The proton and carbon signals in the ^1H - and ^{13}C -NMR spectra of **3** and **4** were superimposable on those of echinacoside,^{1,4,12)} except for the signals due to the 7-methoxy group. The connectivities of the *trans*-caffeoyl group and the glycosyl moieties in **3** and **4** were elucidated on the basis of HMBC experiments as shown in Fig. 1. Finally, alkaline hydrolysis of **3** and **4** with 5% KOH gave the deacylated products together with *trans*-caffeic acid. Those deacylated products were successively treated with 1.0 M HCl to liberate L-rhamnose and D-glucose, respectively. Consequently, the structure of kankanosides K_1 and K_2 were determined to be 2-methoxy-2-(3,4-dihydroxyphenyl)ethyl *O*- α -L-rhamnopyranosyl-(1 $\rightarrow$ 3)-[β -D-glucopyranosyl-(1 $\rightarrow$ 6)]-4-*O*-*trans*-caffeoyl- β -D-glucopyranoside (**3** and **4**).^{11,13)}

Previously, methanolic extract from stems of *C. tubulosa* and several phenylethanoid constituents such as echinacoside, acteoside, and isoacteoside were found to show hepatoprotective effects on D-galactosamine (D-GalN)/lipopolysaccharide-induced liver injury in mice and inhibitory effect on D-GalN-induced cytotoxicity in primary cultured mouse hepatocytes.¹⁾ We further examined inhibitory effects of kankanosides K_1 (**3**) and K_2 (**4**), and isocampneoside I (**5**) on D-GalN-induced cytotoxicity in primary cultured hepatocytes. Although their activities were weaker than those of echinacoside ($\text{IC}_{50}=10.2\ \mu\text{M}$), acteoside ($4.6\ \mu\text{M}$), and isoacteoside ($5.3\ \mu\text{M}$), the principle phenylethanoid constituents from stems of *C. tubulosa*,¹⁾ **3**—**5** showed moderate activity.¹⁴⁾

Experimental

The following instruments were used to obtain spectral and physical data: specific rotations, Horiba SEPA-300 digital polarimeter ($l=5\ \text{cm}$); UV spectra, Shimadzu UV-1600 spectrometer; IR spectra, Shimadzu FTIR-8100 spectrometer; ^1H - and ^{13}C -NMR spectra, JEOL JNM-ECA600 (600, 150 MHz) and JEOL JNM-ECS400 (400, 100 MHz) spectrometers with tetramethylsilane as an internal standard; FAB-MS and high-resolution FAB-MS, JEOL JMS-SX 102A mass spectrometer; HPLC detector, Shimadzu RID-10A refractive index, Shimadzu SPD-10A UV-VIS, and Shodex OR-2 optical rotation detectors. HPLC column, Cosmosil 5C₁₈-MS-II and π NAP (Nacalai Tesque Inc., 250 $\times$ 4.6 mm i.d.) and (250 $\times$ 20 mm i.d.) columns were used for analytical and preparative purposes, respectively.

The following experimental conditions were used for chromatography: normal-phase silica gel column chromatography (CC), silica gel 60N (Kanto

Chemical Co., Ltd., 63—210 mesh, spherical, neutral); reversed-phase silica gel CC, Diaion HP-20 (Nippon Rensui) and Chromatorex ODS DM1020T (Fuji Silysia Chemical, Ltd., 100—200 mesh); normal-phase TLC, pre-coated TLC plates with silica gel 60F₂₅₄ (Merck, 0.25 mm); reversed-phase TLC, pre-coated TLC plates with silica gel RP-18 F_{254S} (Merck, 0.25 mm); reversed-phase HPTLC, pre-coated TLC plates with silica gel RP-18 WF_{254S} (Merck, 0.25 mm), detection was achieved by spraying with 1% Ce(SO₄)₂–10% aqueous H₂SO₄, followed by heating.

Plant Material This item was described in a previous report.¹⁾

Extraction and Isolation Fresh stems of *C. tubulosa* (2.98 kg) were finely cut and extracted three times with methanol under reflux for 3 h. Evaporation of the solvent under reduced pressure provided a methanolic extract (249.1 g, 8.36%). The methanolic extract was subjected to Diaion HP-20 CC (5.0 kg, H₂O→MeOH) to give H₂O- and MeOH-eluted fractions (167.84 g, 5.63% and 81.21 g, 2.73%, respectively). The MeOH-eluted fraction (61.00 g) was subjected to normal-phase silica gel CC [1.8 kg, CHCl₃→MeOH–H₂O (15:3:0.4→10:3:0.5→6:4:1, v/v/v)→MeOH] to give seven fractions [Fr. 1 (1.12 g), 2 (9.56 g), 3 (0.89 g), 4 (10.69 g), 5 (8.84 g), 6 (12.52 g), and 7 (4.60 g)], as was described previously.¹⁾ The fraction 4 (10.69 g) was separated by reversed-phase silica gel CC [500 g, MeOH–H₂O (30:70, v/v)→MeOH→acetone] to give four fractions [Fr. 4-1 (878.2 mg), 4-2 (7.06 g), 4-3 (1.57 g), and 4-4 (792.8 mg)]. The fraction 4-3 (1.57 g) was purified by HPLC [Cosmosil 5C₁₈-MS-II, CH₃CN–1% aqueous AcOH (20:80, v/v)] to give 11 fractions [Fr. 4-3-1 (30.4 mg), 4-3-2 (55.2 mg), 4-3-3 [=campneoside I (6, 22.1 mg, 0.0010%)], 4-3-4 [=acteoside (224.6 mg, 0.010%)], 4-3-5 (27.4 mg), 4-3-6 (43.6 mg), 4-3-7 [=isoacteoside (825.0 mg, 0.037%)], 4-3-8 [=syringalide A 3'-O- α -L-rhamnopyranoside (37.6 mg, 0.0017%)], 4-3-9 (39.8 mg), 4-3-10 [=2'-acetylacteoside (85.4 mg, 0.0038%)], and 4-3-11 (64.6 mg)], as was described previously.¹⁾ The fraction 4-3-5 (27.4 mg) was further purified by HPLC [Cosmosil π NAP, CH₃CN–1% aqueous AcOH (18:82, v/v)] to give isocampneoside I (5, 8.5 mg, 0.0004%). The fraction 4-3-6 (43.6 mg) was further purified by HPLC [Cosmosil π NAP, CH₃CN–1% aqueous AcOH (18:82, v/v)] to give 5 (3.7 mg, 0.0002%) together with kankanosides H₁¹⁾ (17.0 mg, 0.0008%) and H₂¹⁾ (3.3 mg, 0.0001%). The fraction 4-3-9 (39.8 mg) was further purified by HPLC [Cosmosil π NAP, CH₃CN–1% aqueous AcOH (18:82, v/v)] to give kankanosides J₁ (1, 3.7 mg, 0.0002%) and J₂ (2, 3.5 mg, 0.0002%) together with kankanoside I¹⁾ (15.4 mg, 0.0007%) and isoacteoside¹⁾ (3.1 mg, 0.0001%). The fraction 5 (8.84 g) was separated by reversed-phase silica gel CC [400 g, MeOH–H₂O (20:80→30:70, v/v)→MeOH→acetone] to give seven fractions [Fr. 5-1 (870.2 mg), 5-2 (478.9 mg), 5-3 (3.72 g), 5-4 (979.9 mg), 5-5 (1.19 g), 5-6 (1.27 g), and 5-7 (130.1 mg)]. The fraction 5-3-4 (72.3 mg) was further purified by HPLC [Cosmosil π NAP, CH₃CN–1% aqueous AcOH (10:90, v/v)] to give kankanosides K₁ (3, 5.1 mg, 0.0002%) and K₂ (4, 10.6 mg, 0.0005%) together with campneoside II¹⁾ (7, 10.6 mg, 0.0005%).

Kankanoside J₁ (1): A white powder, $[\alpha]_D^{25}$ –6.5° (*c*=0.25, MeOH). High-resolution positive-ion FAB-MS: Calcd for C₃₂H₄₀O₁₇Na (M+Na)⁺ 719.2163; Found 719.2170. UV [$\lambda_{\max}$ (log ϵ), MeOH]: 291 (sh, 3.68), 333 (3.83) nm. IR (KBr, cm^{–1}): 3413, 1734, 1719, 1701, 1638, 1508, 1159, 1067, 1046. ¹H-NMR (600 MHz, CD₃OD) δ : given in Table 1. ¹³C-NMR (150 MHz, CD₃OD) δ : given in Table 2. Positive-ion FAB-MS *m/z*: 719 (M+Na)⁺. Negative-ion FAB-MS *m/z*: 695 (M–H)[–].

Kankanoside J₂ (2): A white powder, $[\alpha]_D^{25}$ –18.1° (*c*=0.23, MeOH). High-resolution positive-ion FAB-MS: Calcd for C₃₂H₄₀O₁₇Na (M+Na)⁺ 719.2163; Found 719.2167. UV [$\lambda_{\max}$ (log ϵ), MeOH]: 291 (sh, 3.99), 333 (4.15) nm. IR (KBr, cm^{–1}): 3418, 1734, 1717, 1638, 1509, 1159, 1070, 1046. ¹H-NMR (600 MHz, CD₃OD) δ : given in Table 1. ¹³C-NMR (150 MHz, CD₃OD) δ : given in Table 2. Positive-ion FAB-MS *m/z*: 719 (M+Na)⁺. Negative-ion FAB-MS *m/z*: 695 (M–H)[–].

Kankanoside K₁ (3): A white powder, $[\alpha]_D^{25}$ –75.3° (*c*=0.16, MeOH). High-resolution positive-ion FAB-MS: Calcd for C₃₆H₄₈O₂₁Na (M+Na)⁺ 839.2586; Found 839.2589. UV [$\lambda_{\max}$ (log ϵ), MeOH]: 289 (sh, 4.05), 335 (4.24) nm. IR (KBr, cm^{–1}): 3415, 1734, 1717, 1686, 1636, 1614, 1509, 1159, 1074. ¹H-NMR (600 MHz, CD₃OD) δ : given in Table 1. ¹³C-NMR (150 MHz, CD₃OD) δ : given in Table 2. Positive-ion FAB-MS *m/z*: 839 (M+Na)⁺.

Kankanoside K₂ (4): A white powder, $[\alpha]_D^{25}$ –7.4° (*c*=0.35, MeOH). High-resolution positive-ion FAB-MS: Calcd for C₃₆H₄₈O₂₁Na (M+Na)⁺ 839.2586; Found 839.2594. UV [$\lambda_{\max}$ (log ϵ), MeOH]: 291 (sh, 4.06), 334 (4.23) nm. IR (KBr, cm^{–1}): 3415, 1734, 1717, 1686, 1636, 1614, 1509, 1159, 1079. ¹H-NMR (600 MHz, CD₃OD) δ : given in Table 1. ¹³C-NMR (150 MHz, CD₃OD) δ : given in Table 2. Positive-ion FAB-MS *m/z*: 839 (M+Na)⁺.

Alkaline and Acid Hydrolysis of Kankanosides J₁ (1), J₂ (2), K₁ (3), and K₂ (4) Solutions of 1–4 (each 1.0 mg) in 5% aqueous potassium hydroxide (KOH, 0.5 ml) were stirred at 40 °C for 1 h. Each solution was neutralized with Dowex HCR W2 (H⁺ form), and the resin was removed by filtration. Evaporation of the solvent from the filtrates under reduced pressure yielded the corresponding deacylated products, which were subjected to HPLC analysis [column: Cosmosil π NAP, 250×4.6 mm i.d.; mobile phase: CH₃CN–1% aqueous AcOH (15:85, v/v); detection: UV (254 nm); flow rate: 1.0 ml/min] to give *trans*-caffeic acid (*t*_R 9.9 min from 1–4). Then each was dissolved in 1.0 M HCl (1.0 ml) and heated at 80 °C for 3 h. After being cooled, the reaction mixture was neutralized with Amberlite IRA-400 (OH[–] form), and the resins were removed by filtration. After removal of the solvent under reduced pressure, the residue was separated by Sep-Pak C18 cartridge column (H₂O→MeOH). The H₂O-eluted fraction was subjected to HPLC analysis under following conditions: HPLC column, Kaseisorb LC NH₂-60-5, 4.6 mm i.d.×250 mm (Tokyo Kasei Co., Ltd., Tokyo, Japan); detection, optical rotation [Shodex OR-2 (Showa Denko Co., Ltd., Tokyo, Japan); mobile phase, CH₃CN–H₂O (85:15, v/v); flow rate 0.8 ml/min]. Identification of L-rhamnose (i) and D-glucose (ii) from 1–4 present in the H₂O-eluted fractions were carried out by comparison of their retention times and optical rotation with those of authentic samples [i, *t*_R 9.9 min (negative)] and [ii, *t*_R 17.9 min (positive)].

Acknowledgements T. M., K. N., and O. M. were supported by 'High-tech Research Center' Project for Private Universities: matching fund subsidy from Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan, 2007–2011 and also supported by a Grant-in Aid for Scientific Research from MEXT. M. Y. and H. M. were supported by the 21st COE program, Academic Frontier Project, and a Grant-in Aid for Scientific Research from MEXT.

References and Notes

- 1) Part XXXV: Morikawa T., Pan Y., Ninomiya K., Imura K., Matsuda H., Yoshikawa M., Yuan D., Muraoka O., *Bioorg. Med. Chem.*, **18**, 1882–1890 (2010).
- 2) Morikawa T., Xie H., Wang T., Matsuda H., Yoshikawa M., *Chem. Biodiv.*, **6**, 411–420 (2009).
- 3) Muraoka O., Morikawa T., Zhang Y., Ninomiya K., Nakamura S., Matsuda H., Yoshikawa M., *Tetrahedron*, **65**, 4142–4148 (2009).
- 4) Yoshikawa M., Matsuda H., Morikawa T., Xie H., Nakamura S., Muraoka O., *Bioorg. Med. Chem.*, **14**, 7468–7475 (2006).
- 5) Xie H., Morikawa T., Matsuda H., Nakamura S., Muraoka O., Yoshikawa M., *Chem. Pharm. Bull.*, **54**, 669–675 (2006).
- 6) Si C.-L., Liu Z., Kim J.-K., Bae Y.-S., *Holzforchung*, **62**, 197–200 (2008).
- 7) The ¹H- and ¹³C-NMR spectra of 1–4 were assigned with the aid of distortionless enhancement by polarization transfer (DEPT), double quantum filter correlation spectroscopy (DQF COSY), heteronuclear multiple quantum coherence (HMQC), and heteronuclear multiple bond correlation (HMBC) experiments.
- 8) Imakura Y., Kobayashi S., Mima A., *Phytochemistry*, **24**, 139–146 (1985).
- 9) Wu J., Huang J., Xiao Q., Zhang S., Xiao Z., Li Q., Long L., Huang L., *Magn. Res. Chem.*, **42**, 659–662 (2004).
- 10) Kitagawa S., Tsukamoto H., Hisada S., Nishibe S., *Chem. Pharm. Bull.*, **32**, 1209–1213 (1984).
- 11) Stereochemistries of 7-position in 1–4 have not been determined.
- 12) Kobayashi H., Oguchi H., Takizawa N., Miyase T., Ueno A., Usamanghani K., Ahmad M., *Chem. Pharm. Bull.*, **35**, 3309–3314 (1987).
- 13) Campneoside I (6), a methylated version of campneoside II (7), was reported to be isolated from water extracts of leaves of *Campsis chinensis*⁸⁾ and fruit of *Forsythia viridissima*.¹⁰⁾ In the present study, 6 could not be produced by treatment of campneoside II (7) with methanol under reflux for more than 24 h. These findings suggest that 1–4 are not artificially produced during the extraction procedure.
- 14) Inhibitory effects of presently isolated kankanosides K₁ (3) and K₂ (4), and isocampneoside I (5, IC₅₀=81.6 μ M) on D-GalN-induced cytotoxicity in primary cultured mouse hepatocytes were examined [inhibition (%) 3: 0.0±1.1, 5.0±1.2, 10.2±2.3*, 16.8±2.9**, and 31.0±3.7**; 4: 0.0±0.6, 5.2±0.1, 10.3±0.7**, 18.0±1.1**, and 24.9±3.4**; 5: 0.0±1.0, 6.0±0.4*, 16.6±0.4**, 31.7±0.9**, and 53.7±2.5** at 0, 3, 10, 30, and 100 μ M, respectively]. Values are expressed as means±S.E.M. (*n*=4). For statistical analysis, one-way analysis of variance (ANOVA) followed by Dunnett's test was used. Probability (*p*) values less than 0.05 were considered significant (**p*<0.05, ***p*<0.01). The bioassay method was described previously.¹⁾