Stereoselective Vinylogous Mukaiyama Aldol Reaction of α -Haloenals

Yoichi IWASAKI, Ryosuke MATSUI, Takahiro SUZUKI, Atsuo NAKAZAKI, and Susumu KOBAYASHI*

Faculty of Pharmaceutical Sciences, Tokyo University of Science; 2641 Yamazaki, Noda, Chiba 278–8510, Japan.

Received January 11, 2011; accepted January 27, 2011; published online January 31, 2011

We have developed a high-yielding and stereoselective vinylogous Mukaiyama aldol reaction (VMAR) of α -haloenals. Contrary to the simple α,β -unsaturated aldehyde, α -haloenals were found to be reactive affording the corresponding VMAR adducts in excellent yields. Some transformations of VMAR adducts by Pd-mediated cross-coupling were also examined in order to demonstrate the synthetic utility of VMAR of α -haloenals.

Key words vinylogous Mukaiyama aldol reaction; α -haloenal; stereoselectivity; cross-coupling

We previously reported a highly stereoselective vinylogous Mukaiyama aldol reaction (VMAR) using vinylketene silyl *N,O*-acetal **1** and **2**, which provides a unique and remarkable entry to a remote asymmetric induction (Chart 1).^{1–3} From a synthetic point of view, this method can directly afford the *anti*- δ -hydroxy- α,γ -dimethyl- α,β -unsaturated carbonyl unit which is seen in many naturally occurring products. Indeed, VMAR has successfully been utilized in natural product syntheses by many groups^{4–20} including ourselves.^{21–25} However, α,β -unsaturated aldehyde is not generally a good substrate for VMAR in terms of yield (*i.e.*, low to moderate). High yield could be achieved by the addition of a catalytic amount of water,³ or by carrying out the reaction for a prolonged period of time.^{5,11} In this context, we became interested in the employment of α -haloenals as substrates which are considered to be much more reactive compared to simple enals. We also reasoned that the VMAR products from α -haloenals could serve as versatile intermediates for introducing a variety of substituents using well-established Pd-mediated methodologies. Herein, we report a high-yielding and stereoselective VMAR of α -haloenals achieved under standard conditions (TiCl_4 in CH_2Cl_2 , 2.0 eq of aldehyde). Some transformations of VMAR products from α -haloenals by Pd-catalyzed cross-coupling are also described.

In this study, we examined VMAR of α -iodoenal **3a**,²⁶ α -bromoenal **3b**,²⁷ and α -chloroenal **3c**,^{28,29} (Table 1). According to the established protocol, VMAR of α -iodoenal **3a** (2.0 eq) with the vinylketene silyl *N,O*-acetal **2** using TiCl_4 (1.0 eq) afforded the corresponding *anti*-aldol adduct **4a** in excellent yield with high diastereoselectivity (entry 1). In addition, VMAR of α -bromoenal **3b** and α -chloroenal **3c** also provided *anti*-adducts **4b** and **4c**³⁰ in excellent yield, respectively (entries 2, 3). Notably, α -halo substituents improved

the yields of aldol adducts and shortened the reaction time compared with VMAR of tiglic aldehyde **3d** (entry 4). Independently, Hosokawa and Tatsuta *et al.* reported that the VMAR of *ent*-**2** and tiglic aldehyde **3d** afforded *ent*-**4d** in good yield (82%) over a prolonged reaction time (4 d) at low temperature (-60°C).^{5,11}

Stereochemical determination of the aldol adducts **4a** and **4b** were easily achieved by Stille cross-coupling using a catalytic amount of $\text{Pd}(\text{dba})_2$ and Me_4Sn to convert the *anti*-adduct **4d** (Chart 2). The spectroscopic data of synthetic **4d**

Table 1. VMAR of α -Haloenals

Entry	X		Time (h)	Yield (%)	d.r. ^{a)}
1	I	3a	10	95	>20 : 1
2	Br	3b	12	88	>20 : 1
3	Cl	3c	4	93	>20 : 1
4	Me	3d	14	76	>20 : 1

a) Diastereomeric ratio was determined by ¹H-NMR analysis.

Reagents and conditions: (a) Me_4Sn , $\text{Pd}(\text{dba})_2$, Ph_3As , HMPA, 60°C , 92%; (b) Me_4Sn , $\text{Pd}(\text{dba})_2$, HMPA, 60°C , 11% (brsm 72%); (c) TBSOTf, *i*- Pr_2NEt , CH_2Cl_2 , 0°C , 100%; (d) NaBH_4 , $\text{THF-H}_2\text{O}$, RT, 99%; (e) O_3 , CH_2Cl_2 -MeOH, -78°C then NaBH_4 , RT, 38%; (f) *p*-TsOH, acetone, RT, quant.

Chart 1. Vinylogous Mukaiyama Aldol Reaction

Chart 2. Determination of the Stereochemistry of **4a**, **4b** and **4c**

* To whom correspondence should be addressed. e-mail: kobayash@rs.noda.tus.ac.jp

Table 2. Pd-Mediated Cross-Coupling with Vinyl Iodide **4a**

Entry	Conditions	R	Yield (%)
1	Me ₄ Sn, Pd(dba) ₂ , AsPh ₃ , HMPA, 60 °C	Me 4d	92
2	Et ₂ Zn, Pd(PPh ₃) ₄ , THF, 0 °C to RT	Et 7a	72
3	CH ₂ =CHSnBu ₃ , Pd(dba) ₂ , AsPh ₃ , HMPA, RT	7b	96

from **4a** and **4b** agreed with those of the known *ent*-**4d**.¹¹⁾ (Synthetic **4d**: $[\alpha]_D^{25} -17.3$, $c=1.09$ in CH₂Cl₂, reported *ent*-**4d**: $[\alpha]_D^{25} +16.5$, $c=1.16$ in CH₂Cl₂).

Stereochemistry of the aldol adduct **4c** was determined by transforming to the known compound **6**.³¹⁾ After protection of the secondary alcohol with TBSOTf, the chiral auxiliary group was reductively removed with NaBH₄ to provide the alcohol **5**. Ozonolysis of **5**, reduction of the resulting aldehyde, followed by acetal formation gave the known compound **6**. The spectroscopic data of synthetic **6** was identical to those reported for **6**.³¹⁾

Next, a series of representative Pd-catalyzed cross-coupling reactions were applied to vinyl iodide **4a** in order to demonstrate the synthetic utility of the present approach (Table 2). Introduction of methyl group to **4a** (entry 1) is already described in Chart 2. Negishi cross-coupling with Et₂Zn and Pd(PPh₃)₄ afforded ethyl derivative **7a** in 72% yield (entry 2). Introduction of a vinyl group to **4a** was performed by Stille cross-coupling to afford **7b** in excellent yield (entry 3).

In conclusion, we were able to overcome the low reactivity of enals in VMAR by employing α -haloenals as substrates. Moreover, the combination of VMAR of α -haloenal and well-established Pd-catalyzed cross-coupling has been demonstrated to considerably expand the scope of VMAR.

Experimental

General Techniques All non-aqueous reactions were carried out under an argon atmosphere in oven-dried of flame-dried glassware. Tetrahydrofuran (THF) and Et₂O were purchased from Kanto Chemical Co., Inc., Tokyo, Japan in anhydrous grade. CH₂Cl₂ was distilled from CaH₂ immediately before use. TiCl₄ was distilled from granular copper and stored in ampoules. Diisopropylethylamine was distilled from CaH₂ and stored over KOH. Other reagents were used as received. All reactions were monitored by TLC, which was carried out on 0.25 mm Silica Gel 60 F₂₅₄ plates (E. Merck). Flash column chromatography separations were performed on PSQ 100B (Fuji Silysia Co., Ltd., Japan). The NMR spectra (¹H, ¹³C) were determined on a Bruker 600 MHz spectrometer (Avance DRX-600) or on a JEOL 400 MHz spectrometer (JNM-LD400), using CDCl₃ solutions, unless otherwise noted. Chemical shifts for ¹H-NMR were expressed in parts per million (ppm) downfield from tetramethylsilane (δ) in duteriochloroform as internal standard and coupling constants (J) are in hertz (Hz). Multiplicities are indicated as: br (broadened), s (singlet), d (doublet), t (triplet), q (quartet), or m (multiplet). Melting points were determined with SHIMADZU MM-2 or Yanaco MP-3S melting point apparatus and were uncorrected. Optical rotations were recorded using CHCl₃ or CH₂Cl₂ as solvents on a JASCO P-1030 digital polarimeter at room temperature, using sodium D line. Infrared spectra (IR) were recorded on a JASCO FT/IR-410 spectrometer using NaCl (neat) or KBr pellets (solid), and are reported in wavenumbers (cm⁻¹). Mass spectra (MS) were obtained on an Applied Biosystems mass spectrometer (API QSTAR pulsar i) under conditions as High resolution, using poly(ethylene glycol) as internal standard.

General Procedure for Vinylogous Mukaiyama Aldol Reaction of Vinylketene Silyl *N,O*-Acetal **2 with α -Haloenals** To a solution of aldehyde (1.41 mmol, 2.0 eq) in CH₂Cl₂ (3.5 ml), a 1.0 M TiCl₄ solution in CH₂Cl₂ (0.70 ml, 0.705 mmol, 1.0 eq) was added dropwise at -78 °C. Then a solution of **1** (239 mg, 0.705 mmol) in CH₂Cl₂ (3.5 ml) was added dropwise into the aldehyde solution at -78 °C. After stirring for 4 to 16 h at -40 °C, the cold reaction mixture was poured into 1 : 1 mixture of saturated NaHCO₃ aq. and saturated Rochelle salt aq. (25 ml). The mixture was diluted with Et₂O (25 ml), and stirred vigorously at room temperature until the white slurry was completely dissolved. The aqueous layer was extracted two times with Et₂O. The combined organic layer was washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel (elution with hexane/AcOEt) to give the aldol adducts.

Aldol adduct **4a**: TLC, $R_f=0.28$ (hexane/AcOEt=2 : 1). mp 114 °C. ¹H-NMR (600 MHz, CDCl₃) δ : 0.86 (3H, d, $J=6.6$ Hz), 0.93 (3H, d, $J=6.9$ Hz), 0.94 (3H, d, $J=7.0$ Hz), 1.84 (3H, d, $J=6.4$ Hz), 2.02 (3H, d, $J=1.3$ Hz), 2.36 (1H, dq, $J=4.7, 7.0, 6.9$ Hz), 2.94 (1H, ddq, $J=10.5, 9.0, 6.6$ Hz), 3.30 (1H, dd, $J=9.0, 2.4$ Hz), 3.64 (1H, d, $J=2.4$ Hz), 4.20 (1H, dd, $J=9.1, 5.6$ Hz), 4.35 (1H, dd, $J=9.1, 9.1$ Hz), 4.58 (1H, ddd, $J=9.1, 5.6, 4.7$ Hz), 5.82 (1H, dq, $J=10.5, 1.3$ Hz), 6.02 (1H, q, $J=6.4$ Hz). ¹³C-NMR (150 MHz, CDCl₃) δ : 14.3, 15.2, 16.0, 17.8, 21.4, 28.5, 40.0, 58.1, 63.5, 81.7, 113.9, 133.0, 133.5, 139.3, 154.4, 171.3. $[\alpha]_D^{25} -26.6$ ($c=1.00$, CHCl₃). IR (CHCl₃, cm⁻¹) 3486, 2965, 1769, 1687, 1366, 1301, 1209. high resolution (HR)-MS (electrospray ionization (ESI)) m/z : 444.0644 (Calcd for C₁₆H₂₄NO₄Na [M+Na]⁺ 444.0642).

Aldol adduct **4b**: TLC, $R_f=0.52$ (hexane/AcOEt=2 : 1). ¹H-NMR (600 MHz, CDCl₃) δ : 0.90 (3H, d, $J=6.6$ Hz), 0.93 (3H, d, $J=6.9$ Hz), 0.94 (3H, d, $J=7.0$ Hz), 1.81 (3H, d, $J=6.5$ Hz), 2.01 (3H, d, $J=1.4$ Hz), 2.36 (1H, dq, $J=4.6, 7.0, 6.9$ Hz), 3.02 (1H, ddq, $J=10.4, 8.7, 6.6$ Hz), 3.59 (1H, d, $J=2.8$ Hz), 3.79 (1H, dd, $J=8.7, 2.8$ Hz), 4.20 (1H, dd, $J=9.0, 5.5$ Hz), 4.35 (1H, dd, $J=9.0, 9.0$ Hz), 4.58 (1H, ddd, $J=9.0, 5.5, 4.6$ Hz), 5.77 (1H, dq, $J=10.4, 1.4$ Hz), 6.07 (1H, q, $J=6.5$ Hz). ¹³C-NMR (150 MHz, CDCl₃) δ : 14.2, 15.2, 16.1, 16.4, 17.8, 28.5, 38.5, 58.1, 63.5, 80.7, 127.6, 129.6, 132.9, 139.7, 154.4, 171.3. $[\alpha]_D^{25} -16.7$ ($c=1.15$, CHCl₃). IR (neat, cm⁻¹) 3494, 2965, 2875, 1772, 1685, 1455, 1367, 1301, 1209. HR-MS (ESI) m/z : 396.0769 (Calcd for C₁₆H₂₄NO₄NaBr [M+Na]⁺ 396.0780).

Aldol adduct **4c**: TLC, $R_f=0.29$ (hexane/AcOEt=2 : 1). ¹H-NMR (600 MHz, CDCl₃) δ : 0.92 (3H, d, $J=6.6$ Hz), 0.93 (3H, d, $J=7.0$ Hz), 0.94 (3H, d, $J=7.1$ Hz), 1.80 (3H, d, $J=6.6$ Hz), 2.00 (3H, d, $J=1.4$ Hz), 2.35 (1H, dq, $J=4.6, 7.1, 7.0$ Hz), 3.02 (1H, ddq, $J=10.4, 8.8, 6.6$ Hz), 3.54 (1H, d, $J=3.0$ Hz), 3.90 (1H, dd, $J=8.8, 3.0$ Hz), 4.20 (1H, dd, $J=9.0, 5.5$ Hz), 4.35 (1H, dd, $J=9.0, 9.0$ Hz), 4.58 (1H, ddd, $J=9.0, 5.5, 4.6$ Hz), 5.75 (1H, dq, $J=10.4, 1.4$ Hz), 5.85 (1H, q, $J=6.6$ Hz). ¹³C-NMR (150 MHz, CDCl₃) δ : 13.7, 14.1, 15.2, 16.1, 17.8, 28.5, 37.7, 58.1, 63.5, 80.0, 124.5, 132.8, 134.8, 139.9, 154.4, 171.3. $[\alpha]_D^{25} -14.7$ ($c=1.00$, CHCl₃). IR (CHCl₃, cm⁻¹) 3494, 2965, 2875, 1770, 1687, 1390, 1367, 1315, 1209. HR-MS (ESI) m/z : 352.1281 (Calcd for C₁₆H₂₄NO₄NaCl [M+Na]⁺ 352.1286).

Aldol adduct **4d**: TLC, $R_f=0.34$ (hexane/AcOEt=2 : 1). ¹H-NMR (600 MHz, CDCl₃) δ : 0.82 (3H, d, $J=6.7$ Hz), 0.93 (3H, d, $J=6.8$ Hz), 0.94 (3H, d, $J=7.0$ Hz), 1.64 (3H, dd, $J=6.6, 1.0$ Hz), 1.67 (3H, dd, $J=1.0$ Hz), 1.98 (3H, d, $J=1.5$ Hz), 2.35 (1H, dq, $J=4.6, 7.0, 6.8$ Hz), 2.75 (1H, ddq, $J=10.3, 9.3, 6.7$ Hz), 3.33 (1H, br s), 3.67 (1H, br d, $J=9.3$ Hz), 4.19 (1H, dd, $J=9.0, 5.7$ Hz), 4.35 (1H, dd, $J=9.0, 9.0$ Hz), 4.58 (1H, ddd, $J=9.0, 5.7, 4.6$ Hz), 5.48 (1H, dq, $J=6.6, 1.0$ Hz), 5.78 (1H, dq, $J=10.3, 1.5$ Hz). ¹³C-NMR (150 MHz, CDCl₃) δ : 10.5, 13.1, 14.0, 15.2, 16.1, 17.8, 28.4, 37.8, 58.1, 63.4, 82.2, 123.7, 131.7, 134.8, 142.0, 154.5, 171.5. $[\alpha]_D^{25} -17.3$ ($c=1.09$, CH₂Cl₂). IR (neat, cm⁻¹) 3513, 2965, 2929, 2873, 1768, 1687, 1367, 1301, 1280, 1211. HR-MS (ESI) m/z : 332.1822 (Calcd for C₁₇H₂₇NO₄Na [M+Na]⁺ 332.1832).

Alcohol **5**: To a solution of **4c** (38.5 mg, 0.117 mmol) in CH₂Cl₂ (1.2 ml), *i*-Pr₃NEt (0.061 ml, 0.350 mmol) and TBSOTf (0.064 ml, 0.280 mmol) was added at 0 °C. After stirring for 40 min, the reaction was quenched with MeOH. After stirring further 10 min, the mixture was concentrated *in vacuo*. The residue was purified by column chromatography on silica gel (elution with hexane/AcOEt=8 : 1) to give the corresponding TBS ether (51.8 mg, 100% yield). To a solution of the corresponding TBS ether (49.7 mg, 0.112 mmol) in THF (2.8 ml), NaBH₄ (29.6 mg, 0.783 mmol) in water (0.6 ml) was added at 0 °C. After stirring for 3 h at r.t., the reaction was diluted with water and Et₂O. The aqueous layer was extracted two times with Et₂O. The combined organic layer was washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel (elution with hexane/AcOEt=8 : 1) to give alcohol **5** (35.2 mg, 99% yield). Alcohol **5**: colorless oil. TLC,

$R_f=0.59$ (hexane/AcOEt=2:1). $^1\text{H-NMR}$ (600 MHz, CDCl_3) δ : -0.01 (3H, s), 0.01 (3H, s), 0.86 (9H, s), 0.88 (3H, d, $J=6.9$ Hz), 1.16 (1H, t, $J=6.2$ Hz), 1.67 (3H, d, $J=1.3$ Hz), 1.73 (3H, d, $J=6.6$ Hz), 2.80 (1H, ddq, $J=9.7$, 6.5, 6.9 Hz), 3.88 (1H, d, $J=6.5$ Hz), 3.99 (2H, d, $J=6.2$ Hz), 5.23 (1H, dq, $J=9.7$, 1.3 Hz), 5.71 (1H, q, $J=6.6$ Hz). $^{13}\text{C-NMR}$ (150 MHz, CDCl_3) δ : -5.1, -4.8, 13.4, 14.1, 17.4, 18.1, 25.7, 36.4, 69.1, 80.5, 121.5, 128.3, 135.4, 136.8. $[\alpha]_D^{21} -15.2$ ($c=1.02$, CHCl_3); IR (neat, cm^{-1}) 3328, 2958, 2929, 2857, 1660, 1461, 1255, 1095, 1006. HR-MS (ESI) m/z : 341.1658 (Calcd for $\text{C}_{16}\text{H}_{31}\text{O}_2\text{NaSiCl} [\text{M}+\text{Na}]^+$ 341.1674).

Alcohol **6**: Alcohol **5** (12.1 mg, 0.0379 mmol) was diluted in 1:1 $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (2.0 ml) and cooled to -78°C . Ozone was bubbled through the solution until a blue tint was observed (about 3 min), and then argon was passed through the solution for 15 min, after which NaBH_4 (11.5 mg, 0.303 mmol) was added. After stirring for 8 h at r.t., the reaction was quenched with saturated NH_4Cl aq. The aqueous layer was extracted two times with Et_2O . The combined organic layer was washed with brine, dried over anhydrous Na_2SO_4 , filtered, and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel (elution with hexane/AcOEt=2:1) to give the corresponding diol (3.4 mg, 38% yield). To a solution of the corresponding diol (2.2 mg, 9.39 mmol) in acetone (0.5 ml), *p*-TsOH (0.9 mg, 4.69 mmol) was added at r.t. After stirring for 16 h, the reaction was neutralized with saturated NaHCO_3 aq. The aqueous layer was extracted two times with Et_2O . The combined organic layer was dried over anhydrous MgSO_4 , filtered, and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel (elution with petroleum ether/ether=4:1) to give alcohol **6** (1.5 mg, quantitative yield). Alcohol **6**: colorless oil. TLC, $R_f=0.39$ (hexane/AcOEt=1:1). $^1\text{H-NMR}$ (600 MHz, CDCl_3) δ : 0.83 (3H, d, $J=7.0$ Hz), 1.37 (3H, s), 1.42 (3H, s), 1.89—1.82 (1H, m), 2.75 (1H, dd, $J=8.6$, 2.8 Hz), 3.66 (1H, dd, $J=8.1$, 7.5 Hz), 3.69—3.59 (2H, m), 3.95 (1H, ddd, $J=8.8$, 7.5, 6.1 Hz), 4.10 (1H, dd, $J=8.1$, 6.1 Hz). $^{13}\text{C-NMR}$ (150 MHz, CDCl_3) δ : 13.1, 25.7, 26.7, 39.2, 67.5, 68.8, 80.9, 109.4. $[\alpha]_D^{22} +30$ ($c=0.027$, CHCl_3). IR (CHCl_3 , cm^{-1}) 2921, 2852, 2042, 1070, 1045, 804. HR-MS (ESI) m/z : 183.0993 (Calcd for $\text{C}_8\text{H}_{16}\text{O}_3\text{Na} [\text{M}+\text{Na}]^+$ 183.0997).

Methyl derivative **4d**: To a solution of iodide **4a** (30.0 mg, 0.071 mmol) in degassed HMPA (0.8 ml), Me_4Sn (0.04 ml, 0.285 mmol), $\text{Pd}(\text{dba})_2$ (4.1 mg, 0.007 mmol) and AsPh_3 (8.7 mg, 0.028 mmol) were added respectively at r.t. After stirring for 40 min at 60°C , the reaction mixture was poured into H_2O . The mixture was diluted with Et_2O , and the aqueous layer was extracted two times with Et_2O . The combined organic layer was washed with brine, dried over anhydrous Na_2SO_4 , filtered, and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel (elution with hexane/AcOEt=6:1) to give **4d** (20.2 mg, 92% yield).

Ethyl derivative **7a**: To a solution of iodide **4a** (10.7 mg, 0.025 mmol) in degassed HMPA (0.5 ml), $\text{Pd}(\text{dba})_2$ (1.5 mg, 0.0025 mmol) and AsPh_3 (3.1 mg, 0.010 mmol) were added respectively at r.t. Then a 1.08 M Et_2Zn solution in hexane (0.094 ml, 0.102 mmol) was added dropwise at 0°C . After stirring for 2 h at r.t., the reaction mixture was quenched with saturated NH_4Cl aq. The aqueous layer was extracted two times with Et_2O . The combined organic layer was washed with brine, dried over anhydrous Na_2SO_4 , filtered, and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel (elution with hexane/AcOEt=6:1) to give **7a** (5.9 mg, 72% yield). Ethyl derivative **7a**: TLC, $R_f=0.38$ (hexane/AcOEt=2:1). $^1\text{H-NMR}$ (600 MHz, CDCl_3) δ : 0.86 (3H, d, $J=6.6$ Hz), 0.93 (3H, d, $J=6.9$ Hz), 0.94 (3H, d, $J=7.0$ Hz), 1.08 (3H, t, $J=7.6$ Hz), 1.67 (3H, d, $J=6.8$ Hz), 1.98 (3H, d, $J=1.4$ Hz), 2.22—2.11 (2H, m), 2.35 (1H, ddq, $J=4.6$, 7.0, 6.9 Hz), 2.76 (1H, ddq, $J=10.3$, 9.2, 6.6 Hz), 3.30 (1H, d, $J=1.9$ Hz), 3.70 (1H, br d, $J=9.2$ Hz), 4.19 (1H, dd, $J=9.0$, 5.8 Hz), 4.34 (1H, dd, $J=9.0$, 9.0 Hz), 4.58 (1H, ddd, $J=9.0$, 5.8, 4.6 Hz), 5.47 (1H, q, $J=6.8$ Hz), 5.80 (1H, dq, $J=10.3$, 1.4 Hz). $^{13}\text{C-NMR}$ (150 MHz, CDCl_3) δ : 13.0, 13.98, 14.01, 15.2, 16.5, 17.8, 19.8, 28.5, 38.5, 58.1, 63.5, 81.8, 123.5, 131.7, 141.0, 142.2, 154.4, 171.5. $[\alpha]_D^{24} -62$ ($c=0.013$, CHCl_3). IR (neat, cm^{-1}) 3515, 2964, 2931, 2873, 1772, 1685, 1390, 1365, 1301, 1280, 1209. HR-MS (ESI) m/z : 346.1996 (Calcd for $\text{C}_{18}\text{H}_{29}\text{NO}_4\text{Na} [\text{M}+\text{Na}]^+$ 346.1988).

Vinyl derivative **7b**: To a solution of iodide **4a** (30.2 mg, 0.072 mmol) in degassed HMPA (0.8 ml), vinyl(tributyl)tin (45.5 mg, 0.143 mmol), $\text{Pd}(\text{dba})_2$ (4.1 mg, 0.007 mmol) and AsPh_3 (8.8 mg, 0.029 mmol) were added respectively at r.t. After stirring for 10 min at r.t., the reaction mixture was poured into H_2O . The mixture was diluted with Et_2O , and the aqueous layer was extracted two times with Et_2O . The combined organic layer was washed with brine, dried over anhydrous Na_2SO_4 , filtered, and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel (elution with hexane, then hexane/AcOEt=8:1) to give **7b** (22.2 mg, 96% yield). Vinyl derivative **7b**: TLC, $R_f=0.48$ (hexane/AcOEt=2:1). $^1\text{H-NMR}$ (600 MHz,

CDCl_3) δ : 0.87 (3H, d, $J=6.7$ Hz), 0.93 (3H, d, $J=6.9$ Hz), 0.94 (3H, d, $J=7.0$ Hz), 1.79 (3H, d, $J=7.0$ Hz), 1.96 (3H, d, $J=1.5$ Hz), 2.35 (1H, ddq, $J=4.6$, 7.0, 6.9 Hz), 2.94 (1H, ddq, $J=10.4$, 8.7, 6.7 Hz), 3.37 (1H, d, $J=2.4$ Hz), 3.97 (1H, dd, $J=8.7$, 2.4 Hz), 4.20 (1H, dd, $J=9.0$, 5.7 Hz), 4.35 (1H, dd, $J=9.0$, 9.0 Hz), 4.59 (1H, ddd, $J=9.0$, 5.7, 4.6 Hz), 5.24 (1H, br d, $J=11.5$ Hz), 5.51 (1H, dd, $J=17.8$, 1.7 Hz), 5.73 (1H, q, $J=7.0$ Hz), 5.83 (1H, dq, $J=10.4$, 1.3 Hz), 6.58 (1H, dd, $J=17.8$, 11.5 Hz). $^{13}\text{C-NMR}$ (150 MHz, CDCl_3) δ : 13.5, 14.0, 15.2, 16.5, 17.8, 28.5, 39.1, 58.1, 63.5, 79.0, 115.9, 126.9, 131.2, 131.7, 137.5, 142.1, 154.5, 171.5. $[\alpha]_D^{24} -51$ ($c=0.19$, CHCl_3). IR (neat, cm^{-1}) 3505, 2965, 2929, 2875, 1770, 1685, 1390, 1367, 1301, 1211. HR-MS (ESI) m/z : 344.1844 (Calcd for $\text{C}_{18}\text{H}_{27}\text{NO}_4\text{Na} [\text{M}+\text{Na}]^+$ 344.1832).

Acknowledgement This research was supported in part by a Grant-in-Aid for Scientific Research (B) (KAKENHI No. 18390010) from the Japan Society for the Promotion of Science.

References and Notes

- Shirokawa S.-i., Kamiyama M., Nakamura T., Okada M., Nakazaki A., Hosokawa S., Kobayashi S., *J. Am. Chem. Soc.*, **126**, 13604—13605 (2004).
- Shinoyama M., Shirokawa S.-i., Nakazaki A., Kobayashi S., *Org. Lett.*, **11**, 1277—1280 (2009).
- Yamaoka M., Nakazaki A., Kobayashi S., *Tetrahedron Lett.*, **51**, 287—289 (2010).
- Hosokawa S., Ogura T., Togashi H., Tatsuta K., *Tetrahedron Lett.*, **46**, 333—337 (2005).
- Hosokawa S., Yokota K., Imamura K., Suzuki Y., Kawarasaki M., Tatsuta K., *Tetrahedron Lett.*, **47**, 5415—5418 (2006).
- Hosokawa S., Kuroda S., Imamura K., Tasuta K., *Tetrahedron Lett.*, **47**, 6183—6186 (2006).
- Jiang X., Liu B., Lebreton S., De Brabander J. K., *J. Am. Chem. Soc.*, **129**, 6386—6387 (2007).
- Nicolaou K. C., Guduru R., Sun Y.-P., Banerji B., Chen D. Y.-K., *Angew. Chem. Int. Ed.*, **46**, 5896—5900 (2007).
- Nicolaou K. C., Sun Y.-P., Guduru R., Banerji B., Chen D. Y.-K., *J. Am. Chem. Soc.*, **130**, 3633—3644 (2008).
- Schmauder A., Müller S., Maier M. E., *Tetrahedron*, **64**, 6263—6269 (2008).
- Hosokawa S., Yokota K., Imamura K., Suzuki Y., Kawarasaki M., Tatsuta K., *Asian J. Chem.*, **3**, 1415—1421 (2008).
- Lipshutz B. H., Amorelli B., *J. Am. Chem. Soc.*, **131**, 1396—1397 (2009).
- Wang L., Gong J., Deng L., Xiang Z., Chen Z., Wang Y., Chen J., Yang Z., *Org. Lett.*, **11**, 1809—1812 (2009).
- Hosokawa S., Mukaeda Y., Kawahara R., Tatsuta K., *Tetrahedron Lett.*, **50**, 6701—6704 (2009).
- Hosokawa S., Matsushita K., Tokimatsu S., Toriumi T., Suzuki Y., Tatsuta K., *Tetrahedron Lett.*, **51**, 5532—5536 (2010).
- Paterson I., Kan S. B. J., Gibson L. J., *Org. Lett.*, **12**, 3724—3727 (2010).
- Wang L., Xi Y., Yang S., Zhu R., Liang Y., Chen J., Yang Z., *Org. Lett.*, **13**, 74—77 (2011).
- Tatsuta K., Hosokawa S., *Chem. Rev.*, **105**, 4707—4729 (2005).
- Tatsuta K., Hosokawa S., *Chem. Rev.*, **6**, 217—233 (2006).
- Hosokawa S., Tatsuta K., *Mini-Rev. Org. Chem.*, **5**, 1—18 (2008).
- Nakamura T., Shirokawa S.-i., Hosokawa S., Nakazaki A., Kobayashi S., *Org. Lett.*, **8**, 677—679 (2006).
- Shirokawa S.-i., Shinoyama M., Ooi I., Hosokawa S., Nakazaki A., Kobayashi S., *Org. Lett.*, **9**, 849—852 (2007).
- Yamaoka M., Fukatsu Y., Nakazaki A., Kobayashi S., *Tetrahedron Lett.*, **50**, 3849—3852 (2009).
- Yamaoka M., Nakazaki A., Kobayashi S., *Tetrahedron Lett.*, **50**, 6764—6768 (2009).
- Matsui R., Seto K., Sato Y., Suzuki T., Nakazaki A., Kobayashi S., *Angew. Chem. Int. Ed.*, **50**, 680—683 (2011).
- Krafft M. E., Cran J. W., *Synlett*, **8**, 1263—1266 (2005).
- Lutjens H., Nowotny S., Knochel P., *Tetrahedron Asymmetry*, **6**, 2675—2678 (1995).
- Kamigata N., Satoh T., Yoshida M., *Bull. Chem. Soc. Jpn.*, **61**, 449—454 (1988).
- Ley S. V., Whittle A. J., *Tetrahedron Lett.*, **22**, 3301—3304 (1981).
- Especially, the electron-withdrawing ability of chloro group probably shortened the reaction time in the VMAR of α -chloroal 3c.
- Nicolaou K. C., Papahatjis D. P., Claremon D. A., Magolda R. L., Dolle R. E., *J. Org. Chem.*, **50**, 1440—1456 (1985).