After cooling, the solid obtained was recrystallized from AcOEt-petr. benzine to colorless platelets, m.p. 83.5~85°. This substance gave a negative permanganate test in acetone. *Anal.* Calcd. for $C_{11}H_{14}O_4$: C, 64.07; H, 4.89. Found: C, 63.61; H, 4.86.

3-(*o*-Carboxyphenyl)butyric Acid (V)—A solution of 2 g. of (IIIa) in 60 cc. of 10% NaOH was heated under stirring at 90° in a water bath. Raney Ni alloy was then added to the solution in small portions over a period of 90 mins. The solution was heated for additional 1 hr. After cooling, the precipitated Ni was filtered and washed with water. The combined aqueous layer was acidified with conc. HCl and the solid obtained formed white crystals from water, m.p. 147.5~149.5°. An admixture with the sample of β -(*o*-carboxyphenyl)butyric acid, m.p. 148~150°, prepared by the ring-opening of 2-cyano-3-methyl-1-indanone, gave no depression of the melting point. The yield was 1.9 g. (95%). *Anal.* Calcd. for $C_{11}H_{12}O_4$: C, 63.45; H, 5.81. Found: C, 63.27; H, 5.85.

Summary

The Reformatsky reaction of ethyl *o*-acetylbenzoate (II) with ethyl bromoacetate gave ethyl 3-methylphthalide-3-acetate (IIIa) as the product, which was led to 3-(*o*-carboxyphenyl)butyric acid (I) via 3-(*o*-carboxyphenyl)crotonic acid (IVa).

The structures of (IIIa) and (IVa) were established by means of their chemical properties and the ultraviolet and infrared absorption spectra, and it was shown that the method was valuable for the verification of each structure of the tautomerism represented by the following general formulae:

(Received October 9, 1956)

U. D. C. 547.94 : 582.751.9

3. Masao Tomita and Tatsuhiko Nakano: Studies on the Alkaloids of Rutaceous Plants. I. Alkaloids of *Phellodendron amurense* RUPR. (1).

(Pharmaceutical Institute, Medical Faculty, University of Kyoto*)

In view of the recent report concerning the isolation of N-methylisocorydine of the aporphine type from *Fagara coco* (GILL) ENGL.¹⁾ and *Xanthoxylum* spp.²⁾ of the Rutaceae family, there seems to be a possibility that some of Rutaceous plants may yield other quaternary alkaloids besides those of the berberine type since so far only berberine type of alkaloids are known throughout the family.

Our attention was first directed to *Phellodendron amurense* RUPR. (Japanese name "Kihada") when it was noted that Murayama and Shinozaki,³⁾ and Takada⁴⁾ in an investigation of this plant reported only the presence of berberine-type alkaloids.

The methanol extract of the plant material, after removal of berberine-type alkaloids as the hydrochlorides, was treated by the modification of the previously reported procedure.³⁾ After purification by reineckate precipitation there was isolated a basic substance as the picrate crystallizing in orange-yellow microscopic pillars, m.p. >300°. It gave analytical values corresponding to $CH_5N_3 \cdot C_6H_3O_7N_3$, and contains no methoxyl nor N-methyl group. The styphnate formed orange-yellow microscopic pillars, $CH_5N_3 \cdot C_6H_3O_8N_3$.

* Yoshida-konoe-cho, Sakyo-ku, Kyoto (富田真雄, 中野立彦).

1) J. Comin, V. Deulofeu: J. Org. Chem., **19**, 1774(1954).

2) J. R. Cannon, G. K. Hughes, E. Ritchie, W. C. Taylor: Australian J. Chem., **6**, 86(1953).

3) Y. Murayama, K. Shinozaki: J. Pharm. Soc. Japan, **530**, 299(1926).

4) Y. Murayama, J. Takada: *Ibid.*, **550**, 1035(1927).

m.p. >230°. An aqueous solution of the hydrochloride caused no precipitate by Meyer's reagent, but it produced pink-colored precipitate with ammonium reineckate. With ninhydrin it gave no color, but an orange-red color with Dragendorff's reagent.⁵⁾ The test⁶⁾ for a guanidino group was positive. These experimental results exhibited a close similarity to guanidine, the identity of which was further confirmed by direct comparison of the infrared spectra of their respective picrates.

The second alkaloid was isolated as the styphnate, yellow needles, m.p. 230~231° (decomp.), from the mother-liquor of guanidine picrate, which was shown by infrared spectrum determination to be identical with magnoflorine.⁷⁾

The mother-liquor left after separation of magnoflorine styphnate was treated with aqueous potassium iodide, whereby the iodide was separated as a brownish oil, which upon crystallization from a mixture of ethanol, methanol, and acetone, formed colorless prisms, m.p. 259°. Analyses of the iodide were in accord with the formula $C_{20}H_{24}O_4NI$, $[\alpha]_D^{25} -147^\circ$ (methanol). Its infrared spectrum showed the presence of a phenolic hydroxyl function. It gave a negative test for a methylenedioxy group with Gaebel's and Labat's reagents. The ultraviolet absorption spectrum of the iodide is shown in Fig. 1.

Fig. 1.

Ultraviolet Absorption Spectrum of
Phellodendrine Iodide (in MeOH soln.)

The third alkaloid appears to be new and it is proposed to designate it phellodendrine. Because of the small amount available, however, a precise description of the basic skeleton of phellodendrine must await further experiments.

The residual substance from the mother-liquor of phellodendrine iodide was obtained as a crystalline styphnate forming yellow needles and decomposing around 230° with explosion. With ninhydrin it gave no color, but with Dragendorff's reagent it developed an orange-red color. With Meyer's reagent it produced no precipitate and analyses confirmed the absence of methoxyl and N-methyl groups. The test for sulfur and the Beilstein's test were negative. This substance appears to differ somewhat from guanidine in its properties, but its small amount did not permit further investigations.

The foregoing results are summarized below.

<i>Phellodendron amurense</i> Rupr.	{	Berberine
		Palmatine
{	{	Magnoflorine
		Phellodendrine (new alkaloid)
		Guanidine
		Unknown base

5) T. Nakano : This Bulletin, **2**, 328(1954); R. Munier, M. Macheboeuf : Bull. Soc. chim. biol., **31**, 1144(1949).

6) S. Sakaguchi : J. Biochem. Japan, **5**, 25(1925).

7) T. Nakano : This Bulletin, **2**, 329(1954).

We are indebted to Mr. S. Asano of Alps Pharmaceutical Co., Furukawa-cho, Gifu, for providing us with the plant material used in this work; to Mr. Narisada of the Research Laboratory, Shionogi & Co. Ltd., for measurement of the infrared spectra, and to Mr. Takashima of Osaka University for the ultraviolet absorption spectral data.

Experimental⁸⁾

Isolation of the Alkaloids from *Phellodendron amurense* RUPR.; Guanidine—Two kg. of the dried bark of the plant material was coarsely ground and extracted 3 times with boiling MeOH. The methanolic extract was evaporated *in vacuo* and the syrupy residue was diluted with a small amount of water and acidified by addition of conc. HCl. The crude berberine hydrochloride soon deposited, which was filtered and weighed 169 g. The filtrate was then treated with aq. ammonium reineckate, and the resulting precipitate was decomposed in acetone solution with aq. Ag₂SO₄ and then with aq. BaCl₂. The filtered solution now containing quaternary bases as the chlorides was concentrated *in vacuo* below 50° to a small volume and shaken up with ether to remove impurities. The aq. layer was evaporated *in vacuo* below 50° to dryness, the residue was dissolved in a minimum amount of water, and treated with aq. sodium picrate. The resulting precipitate was crystallized from acetone-MeOH to form orange-yellow, microscopic needles, m.p. >300°. Yield, 4.2 g. This picrate contains no methoxyl nor methylimino group. *Anal.* Calcd. for CH₅N₃·C₆H₃O₇N₃: C, 29.17; H, 2.80; N, 29.17. Found: C, 29.36; H, 2.95; N, 29.54.

A portion of the picrate was dissolved in a mixture of acetone and water, decomposed with 1% HCl, and the liberated picric acid was removed by means of ether. Evaporation of the aq. layer left a crystalline residue, which caused no precipitate with Meyer's reagent but which gave pink-colored precipitate with aq. ammonium reineckate. It produced no color with ninhydrin, but a pink color with Dragendorff's reagent. The test for a guanidyl group was positive.

The styphnate formed (from MeOH-EtOH) orange-yellow, microscopic pillars, m.p. >230°. *Anal.* Calcd. for CH₅N₃·C₆H₃O₈N₃: C, 27.64; H, 2.65; N, 27.63. Found: C, 27.77; H, 2.99; N, 27.81. This base was confirmed to be identical with guanidine by direct comparison of the infrared spectra of their respective picrates (Nujol mull).

Magnoflorine—The mother liquor from guanidine picrate was treated with 1% HCl and the liberated picric acid was eliminated by extraction with ether. The aq. layer was evaporated *in vacuo* below 50° to dryness, and the residue was dissolved in a minimum amount of water and treated with aq. sodium styphnate⁹⁾. The precipitate which formed was crystallized from acetone-MeOH to give 0.5 g. of yellow needles, m.p. 230~231°(decomp.). *Anal.* Calcd. for C₂₀H₂₄O₄N·C₆H₃O₈N₃: C, 53.24; H, 4.47. Found: C, 53.51; H, 4.71. This styphnate was found to be identical with magnoflorine styphnate by direct comparison of their infrared spectra (Nujol mull).

Phellodendrine and Other Bases—The mother liquor left after separation of magnoflorine styphnate was decomposed with HCl and the liberated styphnic acid was removed by ether. The aq. layer, after evaporation *in vacuo* below 50° to dryness, left a syrupy residue, which was dissolved in a minimum amount of water and treated with conc. aq. KI. The resulting oily precipitate was induced to crystallize from MeOH-EtOH-acetone, whereupon 50 mg. of colorless prisms, m.p. 259°, was obtained. $[\alpha]_D^{25} -147^\circ$ ($c=0.0816$, $l=0.5$ dm. in MeOH). *Anal.* Calcd. for C₂₀H₂₄O₄NI: C, 51.18; H, 5.15; N, 2.98. Found: C, 51.58, 51.60; H, 5.38, 5.51; N, 3.07, 3.19. This iodide gave a negative test for a methylenedioxy group with Gaebel's and Labat's reagents. The infrared spectrum showed the presence of a phenolic hydroxyl function (Nujol mull). The ultraviolet absorption spectrum (MeOH solution) is shown in Fig. 1: $\lambda_{\max}^{\text{MeOH}}$ 287 m μ (log ϵ 4.03).

The mother liquor separated from phellodendrine iodide by decantation was treated with aq. sodium styphnate, yielding yellow precipitate. It was crystallized from acetone-EtOH-MeOH to give 0.1 g. of yellow needles decomposing around 230° with explosion. This substance produced no precipitate with Meyer's reagent but an orange-red color with Dragendorff's reagent. It contains no methoxyl nor methylimino group. It also gave no color with ninhydrin. The test for sulfur and the Beilstein's test were negative. The properties of this substance differs somewhat from guanidine styphnate, but because of its small amount no further investigation was made.

Summary

From the bark of *Phellodendron amurense* RUPR. (Rutaceae), four other quaternary alkaloids, guanidine, magnoflorine, phellodendrine (a new base), and an unknown base were isolated, besides those of the berberine type such as berberine and palmatine.

(Received October 12, 1956)

8) All melting points are uncorrected. We are indebted to Dr. K. Hozumi and the members of the Microanalytical Laboratory of this Institute for the microanalytical data.

9) T. Nakano: This Bulletin, 2, 327(1954).