Communications to the Editor

UDC 547.852.7.07

Synthesis of Cinnoline N-Oxides

In the field of benzodiazine N-oxides, syntheses of quinoxaline N-oxide,¹⁾ quinazoline N-oxide,²⁾ and, phthalazine N-oxide³⁾ have been already reported, but the synthesis of cinnoline N-oxide remained to be uninvestigated. The authors have now succeeded in the syntheses of cinnoline 1-oxide (II), 2-oxide (III), 4-methylcinnoline 1-oxide (V), and 2-oxide (VI). Heating of cinnoline (I) with hydrogen peroxide in glacial acetic acid gave cinnoline N-oxide. The product was chromatographed on alumina, and gave two isomeric cinnoline mono-N-oxides, (II), m.p. 110.5~111.5°, UV $\lambda_{\text{max}}^{\text{EtOH}}$ m μ (log ϵ): 219 (4.41), 229.5 (4.40), 303 (3.71), 315 (3.71), 351.5 (3.90), 367 (3.88), (Anal. Calcd. for $\text{C}_8\text{H}_6\text{ON}_2$: C, 65.75; H, 4.14; N, 19.17. Found: C, 66.13; H, 4.23; N, 19.01) and (III), m.p. 125~126°, UV $\lambda_{\text{max}}^{\text{EtOH}}$ m μ (log ϵ): 218 (4.48), 262 (4.44), 308 (3.76), 351 (3.70), 360 (3.70), (Anal. Calcd. for $\text{C}_8\text{H}_6\text{ON}_2$: C, 65.75; H, 4.14; N, 19.17. Found: C, 65.65; H, 4.14; N, 18.84). The apparent productive ratio of (II) and (III) isolated was about 1:2, but the precise ratio was observed to 1:1.4 by nuclear magnetic resonance spectral analysis.*¹

N-Oxidation of 4-methylcinnoline (IV) was carried out under similar conditions to cinnoline N-oxides, and the product was separated by alumina chromatography to (V), m.p. 94~95°, UV $\lambda_{\text{max}}^{\text{EtOH}}$ m μ (log ϵ): 219 (4.42), 229.5 (4.37), 253 (3.93), 308 (3.70), 318 (3.72), 357 (3.94), 369 (3.90), (Anal. Calcd. for $\text{C}_9\text{H}_8\text{ON}_2$: C, 67.48; H, 5.03; N, 17.49. Found: C, 67.66; H, 5.20; N, 17.21) and (VI), m.p. 151~152°, UV $\lambda_{\text{max}}^{\text{EtOH}}$ m μ (log ϵ): 222 (4.33), 261 (4.37), 307 (3.80), 348.5 (3.70), 357 (3.68), (Anal. Calcd. for $\text{C}_9\text{H}_8\text{ON}_2$: C, 67.48; 5.03; N, 17.49. Found: C, 67.58; H, 5.12; N, 17.25). The ratio of isolated (V) to (VI) was about 1:2, which was almost identical with the ratio observed by nuclear magnetic resonance spectra.

Chart 1.

In order to determine the structure of these N-oxides (II), (III), (V), and (VI), cinnoline 1-oxide (II) was synthesized from 5,6,7,8-tetrahydrocinnoline (VIII) by the following method.

N-Oxidation of 5,6,7,8-tetrahydrocinnoline (VIII) gave two isomers of 5,6,7,8-tetrahydrocinnoline N-oxides; (IX), m.p. 100~100.5°, (Anal. Calcd. for $\text{C}_8\text{H}_{10}\text{ON}_2$: C, 63.98; H, 6.71;

1) H. McIlwain: J. Chem. Soc., **1943**, 322; J.K. Landquist: J. Chem. Soc., **1953**, 2816.

2) K. Adachi: Yakugaku Zasshi, **77**, 507 (1957).

3) E. Hayashi, T. Higashino, C. Iijima, Y. Kōno, T. Doihara: Yakugaku Zasshi, **82**, 584 (1962).

*¹ Studies of nuclear magnetic resonance spectra will be reported in the near future.

N, 18.65. Found: C, 63.94; H, 6.72; N, 18.48) and (X), m.p. 127~128°, (*Anal.* Calcd. for $C_8H_{10}ON_2$: C, 63.98; H, 6.71; N, 18.65. Found: C, 63.40; H, 6.61; N, 18.19), which were separated by alumina chromatography. On the other hand, N-oxidation of 3-chloro-5,6,7,8-tetrahydrocinnoline (VII) gave 3-chloro-5,6,7,8-tetrahydrocinnoline N-oxide (XI), m.p. 133~134°, (*Anal.* Calcd. for $C_8H_9ON_2Cl$: C, 52.33; H, 4.88; N, 15.18. Found: C, 52.28; H, 5.15; N, 15.32). Catalytic reduction of (XI) on palladium-carbon gave 5,6,7,8-tetrahydrocinnoline N-oxide, m.p. 100~100.5°, which was confirmed to be identical with (IX) by a comparison of their infrared absorption spectra. As shown in the previous paper of this series, N-oxidation of 3-chloro-5-methylpyridazine^{4,5)} and 3-chloro-6-methylpyridazine^{5,6)} gave the corresponding 1-oxides respectively. Therefore N-oxide (XI) obtained from (VII) should be also 1-oxide. Further evidence for the structure of (IX) and (X) were obtained by nuclear magnetic resonance spectroscopy. It has been found by our investigation⁷⁾ on the nuclear magnetic resonance spectra of pyridazine N-oxide and its derivatives that the peaks of the ring protons appeared in the following order $\tau_{H_3} < \tau_{H_6} < \tau_{H_5} < \tau_{H_4}$ on the spectra of methyl substituted pyridazine N-oxides. This order should be retained in other alkyl substituted pyridazine N-oxides. These signal peaks of aromatic ring protons of (IX) and (X) centered at 1.82 τ , 3.17 τ and 2.19 τ , 2.84 τ respectively, and the peaks at 1.82 τ , 3.17 τ can be assigned to the protons H_3 and H_4 in (IX), and those at 2.19 τ , 2.81 τ to the protons H_6 and H_5 in (X), respectively.

Treating with sodium methoxide, (XI) gave 3-methoxy-5,6,7,8-tetrahydrocinnoline 1-oxide (XII).

Bromination of (IX) and (XI) with N-bromosuccinimide in CCl_4 gave monobromo compounds (XIII), m.p. 145~146°, (*Anal.* Calcd. for $C_8H_9ON_2Br$: C, 41.94; H, 3.93; N, 12.24. Found: C, 41.89; H, 4.04; N, 12.44), and (XIV), m.p. 93~94°, (*Anal.* Calcd. for $C_8H_8ON_2ClBr$: C, 36.36; H, 3.03; N, 10.60. Found: C, 36.55; H, 3.14; N, 10.09) respectively, but (XII) gave a dibromo compound (XVII), m.p. 151~152°, (*Anal.* Calcd. for $C_8H_{10}O_2N_2Br_2$: C, 31.95; H, 2.96; N, 8.28. Found: C, 31.84; H, 3.15; N, 8.06). Further bromination of (XIII) and (XIV) with N-bromosuccinimide in CCl_4 gave dibromo compounds (XV), m.p. 149~150°, (*Anal.* Calcd. for $C_8H_8ON_2Br_2$: C, 31.36; H, 2.60; N, 9.09. Found: C, 31.54; H, 2.87; N, 9.18) and (XVI), m.p. 137~138°, (*Anal.* Calcd. for $C_8H_7ON_2Br_2Cl$: C, 28.02; H, 2.04; N, 8.17. Found: C, 28.34; H, 2.31; N, 8.41). The structure of these compounds (XIII) will be reported in the near future.

Treating with sodium methoxide, (XV) afforded cinnoline 1-oxide, (II), m.p. 110.5~111.5°, which was proved to be identical with (II) obtained by N-oxidation of cinnoline (I) by a comparison of their infrared absorption spectra. Accordingly, the structure of (III) is deduced to be cinnoline 2-oxide. By the same treatment of (XVI) and (XVII), 3-chlorocinnoline 1-oxide (XVIII), m.p. 168~169°, (*Anal.* Calcd. for $C_8H_5ON_2Cl$: C, 53.18; H, 2.77; N, 15.51. Found: C, 52.85; H, 2.53; N, 15.11) and 3-methoxycinnoline 1-oxide

4) M. Ogata, H. Kano: This Bulletin in press.

5) H. Watanabe, M. Ogata, H. Kano: *Ibid.* in press.

6) T. Nakagome: Yakugaku Zasshi, 81, 1048 (1961).

7) K. Tori, M. Ogata, H. Kano: This Bulletin in press.

(XIX), m.p. 94~95° (*Anal.* Calcd. for $C_9H_8O_2N_2$: C, 61.36; H, 4.58; N, 15.90. Found: C, 61.01; H, 4.57; N, 15.52) were obtained. The structure of 4-methylcinnoline N-oxide (V and VI) were elucidated by nuclear magnetic resonance spectral analysis.

In nuclear magnetic resonance spectrum* of cinnoline 1-oxide (II), signal peaks at 1.33 τ , 1.67 τ and 2.50 τ were assigned to that of H_8 , H_3 and H_4 , respectively. On the other hand, in the spectrum of 2-oxide (III), signal peaks at 1.79 τ and 1.94 τ were assigned to that of H_3 and H_4 respectively. The nuclear magnetic resonance spectrum of 4-methylcinnoline N-oxide (V) shows characteristic peaks at 1.35 τ and 1.87 τ , while that of (VI) shows at 1.90 τ . From these results, (V) was considered to be 4-methylcinnoline 1-oxide (V) and (VI) was 2-oxide (VI).

The authors express their appreciation to Prof. Emeritus E. Ochiai of the University of Tokyo and Dr. K. Takeda, Director of this laboratory, for their helpful advices and encouragements.

Shionogi Research Laboratory
Shionogi & Co., Ltd.
Fukushima-ku, Osaka.

Masaru Ogata (尾形 秀)
Hideo Kano (加納日出夫)
Kazuo Tori (通 和夫)

June 27, 1962