authors' were consistent with the proposed structure IA, information available was insufficient for the structure to be considered fully established, because an alternative structure IB could not be excluded. The alternate structure IB was abandoned for the following reasons. In nuclear magnetic resonance spectrum, ketolactone IV shows octet peaks in the region of 5.92~6.87 analysed as AB parts of ABX system (τ_A ; 6.14, τ_B ; 6.66, J_{AB} ; 17.4 c.p.s., J_{AX} ; 8.7 c.p.s., J_{BX} ; 4.0 c.p.s.). Nuclear magnetic resonance spectra of IV, as well as those of its related compounds (I, II, and III), indicate that these octet peaks mentioned above originated from the two methylene protons on C-11 adjacent to the carbonyl carbon C-12, and furthermore, this methylene group is neighbored by methine function (C-9-H). Since the structure IB has been eliminated spectroscopically as shown above, sciadin, dimethylsciadinonate and sciadinone have been proved to have the structure IA, II, and III, respectively.

The authors express their gratitude to Mr. Y. Kawazoe and Mr. H. Hotta, Faculty of Pharmaceutical Sciences, Tokyo University, for the interpretation of NMR spectra. They also express their deep thanks to Tokyo University Forests for the gift of the materials.

Research Institute of Dental
Materials, Tokyo Medical &
Dental University, Yushima,
Bunkyo-ku, Tokyo.

Chikara Kaneko (金子主税)
Takashi Tsuchiya (土屋 隆)
Masayuki Ishikawai (石川正幸)

November 13, 1962

UDC 543.41:547.724.1+547.456

The Mechanism of the New Color Reaction of Anthrone with Furfural and Pentose

The studies on the color reaction of anthrone with carbohydrate introduced by Dreywood¹⁾ have been reported by many investigators. Sattler and Zerban,²⁾ Shriver, Webb, and Swanson,³⁾ and Yemm and Willis⁴⁾ assumed from the results of spectrophotometrical studies that the color was due to furfural compound derived from carbohydrate by strong acid. On the other hand, Momose, *et al.*⁵⁾ isolated one of the main dyes as trianthronylidenepentane from the reaction mixture of anthrone with pentose or hexose, and discussed a new reaction mechanism without the formation of furfural compound.

Recently the authors have reported in the preceding paper⁶⁾ that furfural gave a specific blue color with anthrone when the reaction mixture was cooled enough to keep away from the evolution of heat. When pentose was previously heated in concentrated acid in the absence of anthrone, it also gave the same blue color with anthrone, though pentose itself gave no color without heating. As this specific color was very unstable by heat, the reaction mixture should be kept in cold. The intensity of the blue color

- 1) R. Dreywood: Ind. Eng. Chem., Anal. Ed., 18, 499 (1946).
- 2) L. Sattler, F. W. Zerban: Science, 108, 207 (1948).
- 3) E. H. Shriver, M. B. Webb, J. W. Swanson: TAPPI., 33, 578 (1950).
- 4) E. W. Yemm, A. J. Willis: Biochem. J., 57, 508 (1954).
- 5) T. Momose, Y. Ueda, K. Sawada, A. Sugi: This Bulletin, 5, 31 (1957).
- 6) R. Sawamura, T. Koyama: Yakugaku Zasshi, 81, 1689 (1961).

thus formed was proportional to furfural concentration, and could be used for the determination of pentose.

In the present communication, the specific color substance was isolated and its structure was investigated.

The color substance of this reaction was separated on alumina or CaHPO_4 with benzene or petr. ether as eluate, and one of the main dyes was obtained as yellow syrup I. I gave a blue color in concentrated sulfuric acid, and it seemed that I had the formula $\text{C}_{19}\text{H}_{12}\text{O}_2$ from the data of elementary analyses. UV $\lambda_{\text{max}}^{\text{petr. ether}}$ $m\mu$: 320, 400. UV $\lambda_{\text{max}}^{70\% \text{ H}_2\text{SO}_4}$ $m\mu$: 600. IR cm^{-1} : $\nu_{\text{C=O}}^{\text{liq. film}}$ 1661. Pentose also gave I in the same reaction with anthrone after it was previously heated in concentrated acid in the absence of anthrone, but another carbohydrate, hexose, methylpentose, etc., gave not I.

On the other hand, it was reported that 10-benzylidene-9-anthrone (III) was prepared by the condensation of anthrone and benzaldehyde in pyridine in the presence of piperidine.⁷⁾ From the reaction mixture of anthrone and furfural in the same procedure, yellow syrup II was yielded.

I and II have the same formula $\text{C}_{19}\text{H}_{12}\text{O}_2$ and also give a blue color in concentrated sulfuric acid respectively. The absorption spectra of I and II in the infrared and ultra-violet region are identical, and are similar to those of III. Furthermore, the infrared absorption at 740, 880, and 1020 cm^{-1} shows that I and II are furan compound.^{8,9)}

Thus, the main color substance I formed from the new color reaction introduced by the present authors⁸⁾ is considered as a new substance having the structure of 10-furfurylidene-9-anthrone. This substance is rapidly decomposed by heat, and therefore, it is presumed that there is no furfurideneanthrone in the reaction mixture of the color reaction introduced by Dreywood.¹⁾

*Pharmaceutical Institute,
College of Science and Engineering,
Nihon University.
Kanda-surugadai, Chiyoda-ku, Tokyo.*

Ryoji Sawamura (沢村良二)
Takashi Koyama (小山 隆)

November 16, 1962

7) M. R. Padova: *Ann. Chim.*, (8), 19, 378 (1910).

8) M. Yamaguchi: *Bunseki Kagaku*, 7, 210 (1958).

9) T. Momose, Y. Ueda, M. Nakamura: *This Bulletin*, 8, 827 (1960).