National Institute of Radiological Sciences,
9-1, Anagawa 4-chome,
Chiba, Japan

Mikio Yamazaki (山崎幹夫)

Faculty of Pharmaceutical Sciences,
University of Tokyo,
Hongo, Tokyo, Japan

Shoji Shibata (柴田承二)

Received November 13, 1965

[Chem. Pharm. Bull.
14(1) 97-99 (1966)]

UDC 581.19 : 582.542.1 : 547.597

Further Characterization of Triterpenoids of Migrated Hopane and Arborane Groups from Gramineae Plants

Arundoin, one of the two major triterpenoids¹⁾ from rhizomes of *Imperata cylindrica* P. BEAUV. var. *media* HUBBARD (チガヤ) (Gramineae), has been proved to be identical²⁾ with the triterpene from *Arundo conspicua* (Gramineae)³⁾ and the structure has been revised to 3 β -methoxyfern-9(11)-ene (Ia).²⁾ The other triterpene, named cylindrin, has been characterised¹⁾ as methyl ether (II) of isoarborinol.⁴⁾ Considerations on biogenesis⁵⁾ of migrated hopane, *i.e.* fernane, and arborane derivatives from present knowledges on triterpene chemistry,⁶⁾ and the similarity of mass spectral fragmentation,⁵⁾ which would be much influenced by the stereochemical relations of the double bond and the rings D and E,^{4,7)} of fern-9(11)-ene and arbor-9(11)-ene derivatives, lead to our proposal²⁾ of the most probable structure (III) for arbor-9(11)-ene derivatives. This has now been proved to be correct by X-ray analysis of 2 α -bromoarborinone (IV).⁸⁾

Further examinations on the minor constituents of the same plant afforded three triterpenoids of rearranged hopane and arborane groups: simiarenol(adian-5-en-3 β -ol) (Va),⁹⁾ m.p. 209~211°, acetate, m.p. 216~217°, $[\alpha]_D^{25} +70^\circ$ (CHCl₃); a new triterpene (Ib), m.p. 192~193°, $[\alpha]_D^{25} -19^\circ$ (CHCl₃), acetate, m.p. 222~223°, $[\alpha]_D^{25} -9^\circ$ (CHCl₃); and isoarborinol (arbor-9(11)-en-3 β -ol) (IIIb),^{4,10)} m.p. 295~300°, acetate, m.p. 296~298°, in respective yields of 0.001, 0.0005, and 0.0005%. Identification of Va and IIIb has been made by the direct comparison with the authentic specimens.*¹

Although simiarenol (Va) has been correlated with adian-5-ene (Vb),¹¹⁾ placement of the hydroxyl group at 3 β -position has based on ORD and NMR data; thus the acid

*¹ The authors thanks are due to Miss W.H. Hui, University of Hong Kong, for her kind donation of the authentic specimen of simiarenyl acetate.

- 1) T. Ohmoto, K. Nishimoto, M. Ito, S. Natori: This Bulletin, **13**, 224 (1965).
- 2) K. Nishimoto, M. Ito, S. Natori, T. Ohmoto: Tetrahedron Letters, **1965**, 2245.
- 3) G. Eglinton, R. J. Hamilton, M. Martin-Smith, S. J. Smith, G. Subramanian: *Ibid.*, **1964**, 2323.
- 4) H. Vorbrüggen, S. C. Pakrashi, C. Djerassi: *Ann.*, **668**, 57 (1963).
- 5) K. Nishimoto, M. Ito, S. Natori, T. Ohmoto: Kth Symposium on the Chemistry of Natural Products, Abstracts of Papers, p. 185 (Osaka, October 1965).
- 6) *i.e.* L. Ruzicka: *Proc. Chem. Soc.*, **1959**, 341; J. H. Richards, J. B. Hendrickson: "The Biogenesis of Steroids, Terpenes, and Acetogenins," W. A. Benjamin, Inc., New York (1964).
- 7) H. Budikievicz, J. M. Wilson, C. Djerassi: *J. Am. Chem. Soc.*, **85**, 3688 (1963).
- 8) O. Kennard, L. R. di Sanseverino, H. Vorbrüggen, C. Djerassi: Tetrahedron Letters, **1965**, 3433.
- 9) H. R. Arthur, W. H. Hui, R. T. Aplin: *Ibid.*, **1965**, 937.
- 10) W. H. Hui, C. N. Lam: *Phytochemistry*, **4**, 333 (1965).
- 11) H. Ageta, K. Iwata, S. Natori: Tetrahedron Letters, **1964**, 3413.

catalyzed migration reaction of the acetate of Va. has been carried out. From the reaction mixture, 3 β -acetoxyfern-8-ene (VI),²⁾ along with 3 β -acetoxyadian-5(10)-ene (VII), m.p. 276~278°, $[\alpha]_D$ -35.8° (CHCl₃), and a hydrocarbon, was isolated. Since VI has been further derived³⁾ to hopenone-II,¹²⁾ this offers confirmative evidence for the location of hydroxyl group in simiarenol (Va).

The new triterpene, m.p. 192~193° (Ib), designated fernenol (Ib), C₃₀H₅₀O, is a pentacyclic triterpene, having one each of tri-substituted double bond (IR : Ib, 815, 790 cm⁻¹, the acetate, 815, 790 cm⁻¹) and hydroxyl group (IR : Ib, 3530 cm⁻¹, the acetate, 1725, 1250 cm⁻¹). The mass spectrum of Ib (426 (M⁺), 259 (M⁺-167) (base peak), 241 (M⁺-167-H₂O)) suggested the presence of the double bond at 9(11)-position, probably on fernane or arborane nucleus,^{5,7)} since the fragmentation pattern is quite similar to those compounds and different from Δ^7 compounds^{7,13,14)} and other groups of triterpenoids.⁷⁾ Methylation of Ib with methyl iodide and potassium in benzene afforded arundoin (Ia) and oxidation of Ib with chromium trioxide-pyridine to 3-keto compound, followed by the Wolff-Kishner reduction, afforded fern-9(11)-ene (Ic);¹³⁾ thus Ia, Ib, and Ic have been correlated and the structure of the new triterpene has been established as fern-9(11)-en-3 β -ol (Ib).

From benzene extract of whole herbs of *Zoysia japonica* STEUD. (シバ) (Gramineae), after the treatment by the same method as that for *Imperata cylindrica*,¹⁾ four triterpenoids have been isolated each in 0.001% yield and identified with arundoin (Ia), cylindrin (IIIa), fernenol (Ib), and isoarborinol (IIIb).

The triterpene, isolated from the leaves of *Tripetaleia paniculata* (ホソツジ) (Ericaceae) by Kondo, *et al.*,^{*2,15)} m.p. 210~211°, has been proved to be identical with simiarenol (Va), which was isolated originally from a Ericaceae plant.⁹⁾

Biogenesis of migrated hopane derivatives (Ia, Ib, Va) and arborane derivatives (IIIa, IIIb) will be explained as shown in Chart 1^{5,8,13)} and the relationship of hopane-zeorinane group (VIII) and fernane group (I) to arborane group (III) corresponds to those of dammarane group and euphane-tircallane group to lanostane group in tetracyclic triterpenoids, the difference coming from chair and boat conformations of the ring B of all-*trans*-squalene at the initial stage of cation induced cyclization.⁶⁾ The occur-

*2 The authors thank Prof. T. Takemoto, Tohoku University, for his kind gift of the specimen.

12) H. Fazakerley, T.G. Halsall, E.R.H. Jones : J. Chem. Soc., 1959, 1877.

13) H. Ageta, K. Iwata, S. Natori : Tetrahedron Letters, 1963, 1447.

14) S. Nakamura, T. Yamada, H. Wada, Y. Inoue, T. Goto, Y. Hirata : *Ibid.*, 1965, 2017.

15) Y. Kondo, T. Nagaya, Y. Takeuchi, T. Takemoto : Yakugaku Zasshi, 83, 674 (1963).

rence of the pairs of compounds such as Ia and IIIa, and Ib and IIIb, having the same structure in ring A and enantiomorphous in rings C, D, and E, in the same Gramineae plants might be worth while to note from biogenetical and chemotaxonomical points of view.

Chart 1. Biogenesis of Migrated Hopane and Arborane Derivatives

This work has been carried out in close contact with Professor H. Ageta, Showa College of Pharmacy, to whom the authors' thanks are due. They are grateful to Takeda Chemical Industries for the determination of the mass spectra.

National Institute of
Hygienic Sciences,
Tamagawayoga, Setagaya, Tokyo

Faculty of Pharmacy,
Toho University,
Narashino, Chiba

Kazumitsu Nishimoto (西本和光)
Miyoko Ito (伊藤巳代子)
Shinsaku Natori (名取信策)
Taichi Ohmoto (大本太一)

Received November 16, 1965