Stereochemistry and Biogenesis of Serratinine^{1,2)}YASUO INUBUSHI, HISASHI ISHII, BOMPEI YASUI,
and TAKASHI HARAYAMA*Faculty of Pharmaceutical Sciences, Osaka University³⁾*

(Received March 31, 1967)

The stereochemistry of serratinine containing its absolute configuration (E) has been established. The tentative biogenesis of serratinine from lycodoline type alkaloid has been also proposed.

In the preceding two papers⁴⁾ of this series, the structure establishment of serratinine(I) has been described. In this paper, we wish to present the full stereochemistry of serratinine including its absolute configuration and also refer to the tentative biogenesis of this alkaloid.

A/B Ring Junction and the Configuration of C₁₃-OH—von Braun degradation of diacetylserratinine,^{*,6)} (II) gave a neutral cyanobromide (III) in good yield which showed an

I⁵⁾

N-cyano band at 2188 cm⁻¹ in the IR spectrum. An attempt to hydrolyze two acetoxyl groups of the compound (III) with methanolic sodium hydroxide solution led to an unexpected result to give nearly quantitatively a ketal (IV), C₁₇H₂₄O₃N₂. In the IR spectrum of the ketal (IV), a hydroxyl band at 3509 cm⁻¹ and an N-cyano band at 2188 cm⁻¹ were observed but no carbonyl band did appear. Acetylation of the compound (IV) with acetic anhydride in pyridine gave acetyl ketal (V) whose IR spectrum showed an acetyl carbonyl band at 1721 cm⁻¹ but no hydroxyl absorption. The NMR spectrum of this

compound (V) showed a 3H doublet ($J=6$ cps) due to a secondary methyl group at 9.06 τ and a 3H singlet attributable to an acetyl methyl at 7.97 τ together with a multiplet due to a proton geminal to an acetoxyl group at 4.97 τ . In addition to these signals, the spectrum showed also the complex signals corresponding to three protons between 5.84 τ and 6.18 τ which could be assigned to protons attached to the carbon atoms bearing ether linkages. These observations can be well rationalized by assuming that treatment of cyanobromide (III) with alkali caused the ketal formation as shown in Chart 1. The only remaining problem to be elucidated is which of the two hydroxyl groups in serratinine participated in this ketal formation. This ambiguity was then clarified by inspection of NMR spectrum of dehydration product from the ketal (IV). Treatment of the ketal with phosphorus oxychloride in pyridine gave anhydro ketal (VI) whose NMR spectrum showed a 3H singlet due to a vinyl methyl group at 8.29 τ . This fact indicates that the free hydroxyl group of the ketal should

- 1) Studies on the Constituents of Domestic Lycopodium Genus Plants. Part VI. Part V: *Chem. Pharm. Bull.* (Tokyo), **16**, 92 (1968).
- 2) The preliminary report of this work appeared in *Tetrahedron Letters*, **1966**, 1551.
- 3) Location: 6-5, Toneyama, Toyonaka, Osaka-fu.
- 4) a) Part IV: *Chem. Pharm. Bull.* (Tokyo), **16**, 82 (1968). b) Part V: *ibid.* **16**, 92 (1968).
- 5) The numbering system of lycopodium alkaloids which is usually introduced, is based on the biogenesis of this type alkaloids suggested by Conroy.¹⁵⁾ To that of serratinine, we adapted the analogous manner based on the tentative biogenesis proposed by us to serratinine.
- 6) Physical constants and preparation of the compound marked with an asterisk in this paper appeared in the preceding paper.^{4a)}

geminal to an acetoxyl group at 5.82τ . The high field shifts of these two signals in this compound can be probably explained by the shielding effect of a newly introduced benzene ring. On the basis of examination of models, the situation in which both methyl and proton concerned are subject simultaneously to the shielding effect, is only possible in a case that both $\text{C}-\text{CH}_3$ and C_8 proton exist in *trans* diequatorial arrangement. Therefore, it may safely be said that the C_8 hydroxyl group in serratinine (I) should be *cis* to the $\text{C}-\text{CH}_3$ group, namely this hydroxyl group being axial.

This assignment to the C₈ hydroxyl group was also supported by the following observations. Dehydration of serratinine (I) with phosphorus oxychloride in pyridine at room temperature gave a monoanhydride,^{*,8)}(X) which was verified as 8-anhydroserratinine by the facts that

8) Physical constants and preparation of the compound marked with double asterisk in this paper appeared in the preceding paper.^{4b)}

acetylation of the compound (X) furnished 13-acetyl-8-anhydroserratinine* (XI) described in the previous paper, and hydrolysis of 13-acetyl-8-anhydroserratinine regenerated the compound (X). Consequently, the facile dehydration of the C₈ hydroxyl group by such a mild condition suggests that this group would have axial orientation.

The second evidence was obtained from a comparison of the properties between serratinine (I) and 8-episerratinine (XII). Reduction of 13-acetyl-8-dehydroserratinine* (VIII) with sodium borohydride in ethanol, followed by hydrolysis with alkali gave a mixture which showed virtually two spots on thin-layer chromatography. When the reduction product was chromatographed on alumina, the first eluate was found to be identical with serratinine and the successive elution with ethyl acetate then furnished another crystals (XII) whose IR spectrum showed hydroxyl bands at 3520 and 3340 cm⁻¹ and a ketonic band at 1737 cm⁻¹. Oxidation of the compound (XII) with Jones' reagent gave bisdehydroserratinine* (XIII) which was obtained from serratinine with the same reagent as mentioned in the previous paper.^{4a)} This result coupled with analytical data suggests that the latter compound (XII) would be 8-episerratinine.

The possibility of epimerization at C₇ and/or C₁₅ of the compound (VIII) during reduction process, would be ruled out since the crude reduction product, without being purified, showed virtually two spots corresponding to serratinine (I) and 8-episerratinine (XII), respectively, on thin-layer chromatography, and besides the total yield of both isomers was nearly quantitatively. Furthermore, inspections of models of the compounds which would be expected to be formed by the supposed epimerization, revealed that the epimerized compounds should have abnormally unfavorable structures with marked strain.

It has been well known that the epimer with the equatorial hydroxyl group is more strongly adsorbed on a chromatographic alumina column than the axial epimer, thus suggesting that the C₈ hydroxyl group in serratinine would be axial, that in 8-episerratinine being equatorial. Now, a pair of epimers with regard to C₈ hydroxyl group came to our hands.

Chart 3

On the basis of the structure and stereochemical informations described so far, three conformational structures with regard to ring A are possible, *i.e.* one chair form (A) and two twist forms (B) and (C). Among these structures, in both (A) and (B) the secondary methyl group is *cis* to C₇-H but in (C) *trans* to C₇-H.

Finally, the geometry around C₈ was revealed by means of application of NMR to the conformational problems. Acetylation of 8-episerratinine with acetic anhydride in pyridine gave a monoacetate (XIV) whose IR spectrum showed a hydroxyl band at 3270 cm⁻¹ and carbonyl bands at 1747 and 1732 cm⁻¹. The compound (XIV) was then oxidized with Jones' reagent to give a corresponding ketone (XV) which was not identical with 13-acetyl-8-dehydroserratinine* (VIII) obtained from 13-acetylserratinine* (XVI) under the same oxidation condition, thus indicating that acetylation of 8-episerratinine had occurred at the C₈ hydroxyl group. The NMR spectrum of the monoacetate (XIV) showed a 1H clean quartet (*J*₁=5.5 cps and *J*₂=11 cps) due to a proton geminal to an acetoxyl group centered

at 5.17 τ , while the signals of the corresponding proton in 8-acetylserratinine* (XVII), an epimer of 8-acetyl-8-episerratinine, appeared as a 1H rather sharp multiplet (half band-width 5 cps) at 4.94 τ . This finding suggests that the C₈ hydrogen of 8-acetyl-8-episerratinine (XIV) is axial and that of 8-acetylserratinine* (XVII) is equatorial. The NMR spectrum of 8-acetyl-8-episerratinine was then utilized in conformational analysis around C₈ acetoxyl group of this compound.

When the hydrogens at C₇, C₈ and C₁₅ are designated as H_a, H_b and H_c, respectively, the magnitude of coupling constants between the coupled protons concerned were calculated.⁹⁾ Dreiding model of conformation (A) reveals that the dihedral angle between H_a and H_b is approximately 45°, that between H_b and H_c being nearly 170°. Thus, these values correspond to 4 cps for J_{a-b} and to 11 cps for J_{b-c} , respectively. In conformation (B), the similar values would be expected. In the case of conformation (C), however, H_b is situated in parallel with both H_a and H_c, so that each dihedral angle would be about 180°, suggesting that both coupling constants for J_{a-b} and J_{b-c} would be approximately 11 cps. Since the observed coupling constants were 5.5 cps and 11 cps which would be allocated to J_{a-b} and J_{b-c} , respectively, 8-acetyl-8-episerratinine (XIV) could be depicted by a conformation (A) or (B), but not (C). Conclusively, we may deduce that both C₈ hydroxyl group and a secondary methyl group are *cis* to C₇-H.

Configuration of C₄-N Bond—It has been shown that the large difference in basicity due to hydrogen bonding can be utilized in establishing stereochemistry of some alkaloids. The pK_a values of some acetyl derivatives of serratinine are shown in Chart 4.

That acetylation of C₁₃ hydroxyl group caused a fairly large decreasing in basicity could be accounted for in terms of the existence of specific interaction between C₁₃ hydroxyl group and nitrogen.

Chart 4

Inspections of models reveal that only in the *cis* relationship of C₄-N bond with respect to C₁₂-C₁₃ bond are both hydroxyl and nitrogen functions close enough together to give rise to an intramolecular hydrogen bonding. This deduction, however, is not definitely because the differences in basicity are not large enough to establish conclusively the stereochemistry of C₄-N bond.

9) H. Conroy, "Advances in Organic Chemistry Methods and Results," Vol. II, 1960, Interscience Publishers, Inc., New York, p. 310.

The correctness of this postulation was then shown firmly by the following chemical evidence.

We have already described in a previous paper that lithium aluminum hydride reduction of serratinine (I) gave α -dihydro-serratinine** (XVIII). On the other hand, reduction of serratinine with sodium in *n*-propanol produced another dihydro compound, β -dihydro-serratinine (XIX) which was confirmed to be an epimer of α -dihydro-serratinine with regard to C₅ hydroxyl group by the fact that both dihydro compounds were oxidized with Jones' reagent to give bisdehydro-serratinine* (XIII). On heating at 160° under reduced pressure, β -dihydro-serratinine (XIX) methoxide was subjected to Hofmann degradation to afford a basic methine (XX), C₁₇H₂₉O₃N, in excellent yield. The IR spectrum of this methine showed a hydroxyl band at 3401 cm⁻¹ and a ketonic band at 1709 cm⁻¹¹⁰⁾ and the NMR spectrum showed a 3H singlet due to an N-methyl group at 7.73 τ and no signal due to an olefinic proton. Meanwhile, under the same condition, α -dihydro-serratinine (XVIII) methoxide afforded an unseparable mixture which showed many spots on thin-layer chromatography. These facts indicate that the C₅ hydrogen of β -dihydro-serratinine should be *trans* to C₄-N bond as shown in formula (D).

(D)

At this point, the configuration of C₅ hydrogen of β -dihydro-serratinine must be settled. Reduction of 8-acetyl-13-dehydro-serratinine* (XXI) with sodium borohydride gave a mixture which showed virtually two spots on thin-layer chromatography. Chromatography on

10) The band of this carbonyl group gave rather lower frequencies than those expected for five membered ketones but we now have some evidences that this anomalous position could be ascribed to hydrogen bonding. These findings will be reported in other paper.

alumina resulted in separation of two compounds, *i.e.* serratinine and α -dihydroserratinine** (XVIII) which were identified with authentic specimens, respectively. In contrast with this result, the compounds leaving the C₁₃ hydroxyl group intact, like serratinine or 13-acetyl-8-dehydroserratinine* (VIII), resisted to reduction of C₅ ketone

function with the same reagent even under the forced condition. The insusceptibility to the reagent in the latter cases may be ascribed to the steric hindrance to approach of the metal hydride ion to either side of the C₅ carbonyl function, and this steric interference should be partly due to the C₁₃ hydroxyl group. Therefore, in 8-acetyl-13-dehydroserratinine (XXI) which possesses a carbonyl group instead of a hydroxyl group at C₁₃ position, the metal hydride ion can only approach from the less hindered side, *i.e.* the side on which the C₁₃ hydroxyl group was originally held. In this case, the hydride ion could approach competitively to either the C₅ ketone group or the C₁₃ ketone group. If the hydride ion approaches first to the C₅ ketone group, the initial reduction product could be then subject to further reduction to give α -dihydroserratinine and if the hydride ion approaches to the C₁₃ ketone first, the hydroxyl group newly introduced, rejects further approach of the reagent to the C₅ ketone group, so that the reaction does not proceed any more and serratinine is the only reduction product. Consequently, these considerations allow us to deduce that the configuration of C₅ hydrogen in α -dihydroserratinine (XVIII) should be the same configuration as that of C₁₃ hydroxyl group of serratinine. In other words, the C₅ hydrogen in β -dihydroserratinine (XIX) is *trans* to the C₁₃ hydroxyl group. Since it has been shown by Hofmann degradation studies on two epimeric dihydroserratinines that the C₅ hydrogen in β -dihydroserratinine (XIX) would be *trans* to the C₄-N bond, the configuration of the C₄-N bond in serratinine must be *cis* to the C₁₃ hydroxyl group. Consequently, serratinine should be represented by the stereostructure (E) or its mirror image.

Absolute Configuration—The final problem to be discussed on the stereochemistry of serratinine is concerned with the absolute configuration of this alkaloid. Wolff-Kishner reduction¹¹⁾ of 8-acetyl-13-dehydroserratinine (XXI) afforded an alcohol, 5-deoxo-13-deoxyserratinine (XXII), which was then oxidized with chromium trioxide in 90% acetic acid to give a ketone, 8-dehydro-5-deoxo-13-deoxyserratinine (XXIII). The optical rotatory dispersion curve of this ketone gave a positive Cotton effect with extreme at 273 and 308 m μ ; $a = 47.58$. The positive sign of this curve is in accord with the predicted sign from the octant diagram (F) for the formula (XXIII). Consequently, serratinine should be represented by the formula (E) and not its mirror image.

This conclusion was also supported by the result obtained from application of the benzoate rule.

Treatment of 8-episerratinine (XII) with benzoyl chloride in pyridine gave 8-benzoyl-8-episerratinine (XXIV). Application of the benzoate rule,^{12,13)} to a pair of 8-episerratinine

11) D.H.R. Barton, D.A.J. Ives, and B.R. Thomas, *J. Chem. Soc.*, **1955**, 2056.

12) J.H. Brewster, *Tetrahedron*, **13**, 106 (1961).

13) In a preliminary communication,²⁾ we have applied the benzoate rule to a pair of 13-acetylserratinine and its benzoate. However, the derivations of their $[M]_D$ values by changes of solvents employed were fairly large. For instance, 13-acetyl-8-benzoylserratinine showed $[M]_D$: $+62.3^\circ$ in ethanol solution, but $[M]_D$ -23.9° in chloroform solution. Since the scattering $[M]_D$ values can be probably ascribed to the deviation of ring A by the solvation, application of the benzoate rule to such compounds would not be suitable. 8-Benzoyl-8-episerratinine seems to be more rigid compound than 13-acetyl-8-benzoylserratinine because the former has a more favorable equatorial benzoyloxy group at C₈. Since deviation of conformation of the compound could be perceived by measuring NMR spectrum and its conformation has been presumed by examining the signal due to a proton geminal to a benzoyloxy group in its NMR spectrum taken in chloroform solution, it is desirable that the optical rotation should be measured in the same solvent as that employed in NMR measurement.

Chart 6

(XII) and its benzoate was attempted: $[M]_D(\text{benzoate}) + 212.06^\circ - [M]_D(\text{ol}) + 77.66^\circ = \Delta[M]_D + 134.40^\circ$ (in chloroform). Several years ago, Brewster clearly has explained the benzoate rule by his dissymmetry rule.¹⁴⁾ When we adapt his interpretation, our experimentally observed result can be rationalized as follows. Newman's projection of the surroundings of the bond between O and C_8 in 8-benzoyl-8-episerratinine (XXIV) shows three possible conformers with regard to O-CO bond as shown in Chart 6. However, two of them, (b) and (c), can be excluded for the reason that both of them have 1,3-diaxial-like interaction and seem to be unfavorable forms. The O-CO bond in the most preferable conformer (a) is located between $\text{C}_8\text{--C}_7$ and $\text{C}_8\text{--H}$ bonds.

Since the observed $\Delta[M]_D$ showed positive sign which is expected to the Newman's projection (G) for 8-benzoyl-8-episerratinine (XXIV), it could be concluded that the absolute configuration of the asymmetric center at C_8 in 8-episerratinine (XII) is R-form and that in serratinine (I), accordingly, being S-form. These conclusions are consistent with the former result obtained from the positive Cotton effect for 8-dehydro-13-deoxyserratinine.

Biogenesis—It has been suggested by Conroy¹⁵⁾ that the skeleton inherent in lycopodium alkaloids might owe to the condensation of two eight carbon polyacetate straight chains.

14) J.H. Brewster, *J. Am. Chem. Soc.*, **81**, 5475, 5483, 5493 (1959).

15) H. Conroy, *Tetrahedron Letters*, **1960**, 34.

Lycopodine type alkaloid

Chart 7

However, serratinine is a new representative to which the biogenesis suggested so far could not directly be applied. We, now, wish to propose the extended biogenetic schemes to that of serratinine on the basis of Conroy's biogenesis. If lycopodine or a close related derivative is the precursor of serratinine, this alkaloid is plausibly formed through the postulated steps from the supposed precursor as shown in Chart 7, although the sequence is obscure. This postulation seems to be reasonable, since lycopodine was isolated from the same plant together with serratinine and the presented absolute configuration of serratinine concures with that of an alkaloid which might be expected to arise from the lycopodine type alkaloid through the postulated transformations. The additional C₈ hydroxyl group in serratinine seems to be quite reasonable, since the oxygen functions, lycopodium alkaloids are most commonly found at C₅, C₈ and C₁₃, especially lycofawcine¹⁶ possessing three oxygen functions at these positions. Although these biogenetic schemes are plausible enough and the experimental support is required, this proposed biogenetic pathway would be still useful for predicting the structures of this series of alkaloids which seem to be isolated from the plant.

Experimental¹⁷⁾

Diacetylserratinine Cyanobromide (III)—A mixture of 1 g of diacetylserratinine* (II) and 10 g of cyanogen bromide in 40 ml of dry CHCl₃ was allowed to stand at room temperature for 40 hr. The residue was taken up in CHCl₃ and washed with aq. 5% HCl solution. Some quantity of starting material (II) was recovered from the acidic washings. The chloroform solution was dried over anhydr. K₂CO₃ and removal of the solvent gave 0.91 g of crystals which were recrystallized from acetone-ether to afford leaflets (III) mp 200–202°, [α]_D²⁵ –23.1° (c =1.02, EtOH). *Anal.* Calcd. for C₂₁H₂₉O₅N₂Br: C, 53.73; H, 6.23. Found: C, 53.73; H, 6.24. IR cm⁻¹: $\nu_{\text{C}\equiv\text{N}}$ 2188; $\nu_{\text{C}=\text{O}}$ 1739, 1727; $\nu_{\text{C}-\text{O}}$ 1230. NMR τ : 4.93 (1H, m., >CH–OAc); 5.36 (1H, m., >CH–OAc); 7.97 (3H, s., –OCOCH₃); 8.08 (3H, s., –OCOCH₃); 9.12 (3H, d., J =6 cps, >CH–CH₃).

Ketal (IV)—A solution of 150 mg of diacetylserratinine cyanobromide (III) in 15 ml of MeOH containing 400 mg of KOH was refluxed for 30 minutes, cooled and neutralized with aq. 5% AcOH solution. The solution was concentrated *in vacuo*, diluted with water and extracted with CHCl₃. The chloroform extract was dried over anhydr. K₂CO₃ and removal of the solvent gave 63 mg of crystals which were recrystallized from acetone to give colorless needles, mp 239–240°, [α]_D²⁵ –27.5° (c =1.01, EtOH). *Anal.* Calcd. for C₁₇H₂₄O₃N₂: C, 67.08; H, 7.95. Found: C, 66.94; H, 8.02. IR cm⁻¹: $\nu_{\text{O}-\text{H}}$ 3509; $\nu_{\text{C}\equiv\text{N}}$ 2188.

Acetylketal (V)—A mixture of 100 mg of ketal (IV) and 3 ml of acetic anhydride in 3 ml of pyridine was heated at 100° for 3 hr, cooled and evaporated to dryness *in vacuo*. The residue was taken up in water and extracted with CHCl₃. The extract was dried over anhydr. K₂CO₃ and the solvent was evaporated. The residue in benzene was chromatographed on alumina. Elution with benzene afforded 70 mg of solid mass which was recrystallized from ether to give colorless leaflets (V), mp 184.5–185.5°, [α]_D²⁵ –19.6° (c =0.79, EtOH). *Anal.* Calcd. for C₁₉H₂₆O₄N₂: C, 65.87; H, 7.57. Found: C, 65.62; H 7.67. IR cm⁻¹: $\nu_{\text{C}\equiv\text{N}}$ 2193; $\nu_{\text{C}=\text{O}}$ 1721; $\nu_{\text{C}-\text{O}}$ 1245. NMR τ : 4.97 (1H, m., >CH–OAc); 5.84–6.18 (3H, m., >CH–O–C(=O)–CH₂–); 7.97 (3H, s., –OCOCH₃); 9.06 (3H, d., J =6 cps, >CH–CH₃).

16) W.A. Ayer, W.R. Bowman, P. Kébarle, and R.H. Burnell, *Can. J. Chem.*, **43**, 328 (1965).

17) Refer to the footnote 19) in the preceding paper (Part IV).

Anhydroketal (VI)—To a solution of 50 mg of ketal (IV) in 1 ml of dry pyridine was added 2 drops of POCl_3 . The mixture was allowed to stand at room temperature overnight, poured into water, made acidic with conc. HCl and extracted with CHCl_3 . The extract was dried over anhydr. K_2CO_3 and the solvent was evaporated. The residue in benzene was chromatographed on alumina. Elution with benzene gave 15 mg of solid mass which was recrystallized from benzene-ether to furnish colorless prisms (VI), mp 175–178°. *Anal.* Calcd. for $\text{C}_{17}\text{H}_{22}\text{O}_2\text{N}_2$: C, 71.30; H, 7.74. Found: C, 71.38; H, 7.92. IR cm^{-1} : $\nu_{\text{C}\equiv\text{N}}$ 2183. NMR τ : 4.37 (1H, m., olefinic proton); 5.77–6.18 (3H, m., $\text{>CH-O-C-O-CH}_2\text{-}$); 6.5–6.75 (2H, m., $\text{-CH}_2\text{-N-CN}$); 8.29 (3H, s., vinyl methyl).

Dehydroketal (VII)—A solution of 100 mg of ketal (IV) in 1 ml of pyridine was added to a stirred solution of 1 ml of pyridine containing 100 mg of chromium trioxide. The mixture was stirred at room temperature for 20 hr, poured into ice-water, made alkaline with NH_4OH , and extracted with CHCl_3 . The extract was dried over anhydr. K_2CO_3 and evaporated to dryness *in vacuo*. The residue in CHCl_3 was chromatographed on alumina. Elution with CHCl_3 gave 58 mg of crystals which were recrystallized from acetone to afford colorless prisms (VII), mp 284–285°. *Anal.* Calcd. for $\text{C}_{17}\text{H}_{22}\text{O}_3\text{N}_2$: C, 67.52; H, 7.33; N, 9.27. Found: C, 67.79; H, 7.62; N, 9.13. IR cm^{-1} : $\nu_{\text{C}\equiv\text{N}}$ 2183; $\nu_{\text{C=O}}$ 1692. NMR τ : 5.7–6.2 (3H, m., $\text{>CH-O-C-O-CH}_2\text{-}$); 8.96 (3H, d., $J=6$ cps, >CH-CH_3).

Formation of Dehydroketal (VII) from 13-Acetyl-8-dehydroserratinine* (VIII)—A mixture of 150 mg of 13-acetyl-8-dehydroserratinine* (dehydromonoacetylserratinine II (VIII)) in dry CHCl_3 containing 1.5 g of cyanogen bromide was allowed to stand at room temperature for 40 hr. The chloroform and excess cyanogen bromide were removed under reduced pressure. The residue was taken up in CHCl_3 , washed with aq. 5% HCl solution and dried over anhydr. K_2CO_3 . Removal of the solvent afforded 90 mg of crystals which were recrystallized from acetone-ether to give colorless plates, mp 169–172°. *Anal.* Calcd. for $\text{C}_{19}\text{H}_{25}\text{O}_4\text{N}_2\text{Br}$: C, 53.65; H, 5.93. Found: C, 53.61; H, 5.93. IR cm^{-1} : $\nu_{\text{C}\equiv\text{N}}$ 2193; $\nu_{\text{C=O}}$ 1745, 1733, 1698; $\nu_{\text{C-O}}$ 1233.

A solution of 100 mg of 13-acetyl-8-dehydroserratinine cyanobormide in 12 ml of MeOH containing 0.3 g of KOH was heated for 30 minutes under reflux, cooled and neutralized with aq. 5% AcOH solution. The solution was then concentrated *in vacuo*, diluted with water and extracted with CHCl_3 . The extract was dried over anhydr. K_2CO_3 . Removal of the solvent and recrystallization of the residue from acetone gave 21 mg of colorless prisms, (VII), mp 283–285°, which were identified with the sample derived from the ketal (IV) by comparison of IR spectra and mixed melting point determination.

Diacetylbenzylidene Serratinine (IX)—A solution of 1 g of benzylidene serratinine* in 15 ml of pyridine containing 15 ml of acetic anhydride was heated at 100° for 4.5 hr, cooled and evaporated to dryness *in vacuo*. The residue was dissolved in water, made alkaline with NH_4OH and extracted with CHCl_3 . The extract was dried over anhydr. K_2CO_3 and the solvent was evaporated. The residue in benzene was chromatographed on alumina. Elution with benzene followed by ether gave 786 mg of crystals which were recrystallized from *n*-hexane to afford pale yellow plates, (IX), mp 184–185°. *Anal.* Calcd. for $\text{C}_{22}\text{H}_{33}\text{O}_5\text{N}$: C, 71.81; H, 7.37. Found: C, 71.90; H, 7.39. IR cm^{-1} : $\nu_{\text{C=O}}$ 1727; $\nu_{\text{C=C}}$ 1625, ν_{aromatic} 1600, 1572; $\nu_{\text{C-O}}$ 1239, 1224. UV λ_{max} m μ (log ϵ): 228 (3.88), 291.5 (4.26); $\nu_{\text{max}}^{\text{EtOH-HCl}}$ m μ (log ϵ): 236 (3.71), 309 (4.25). NMR τ : 2.3–2.8 (5H, m., aromatic protons); 4.56 (1H, m., olefinic protons); 5.31 (1H, m., >CH-OAc); 5.82 (1H, m., >CH-OAc); 7.95 (3H, s., -OCOCH_3); 8.07 (3H, s., -OCOCH_3); 9.43 (3H, diffused d., $J=6$ cps, >CH-CH_3).

8-Anhydroserratinine (X)—To a solution of 150 mg of serratinine (I) in 3 ml of pyridine was added 0.5 ml of POCl_3 and the mixture was then allowed to stand at room temperature overnight and evaporated to dryness *in vacuo*. The residue was dissolved in aq. 5% HCl solution, washed with ether, made alkaline with NH_4OH and extracted with CHCl_3 . The extract was dried over anhydr. K_2CO_3 and the solvent was evaporated. The residue in benzene was chromatographed on alumina. Elution with benzene followed by ether gave 55 mg of solid mass which was recrystallized from benzene to afford colorless needles (X), mp 196–198°. This substance was identical with anhydroserratinine II* (X), independently obtained by dehydration of monoacetylserratinine II* (*i.e.* 13-acetyl-8-hydroxy derivative of serratinine) (XI), followed by hydrolysis, as evidenced by mixed melting point determination and infrared spectra comparison. Acetylation of 50 mg of 8-anhydroserratinine (X) by the same manner as described in acetylation of the ketal (IV), gave 25 mg of 13-acetyl-8-anhydroserratinine* (XI).

Reduction of 13-Acetyl-8-dehydroserratinine* (VIII) with NaBH_4 (8-episerratinine (XII))—To a solution of 108 mg of 13-acetyl-8-dehydroserratinine* (VIII) in 200 ml of EtOH was gradually added 400 mg of NaBH_4 in small portions under reflux during 4 hr. A solution of 100 mg of NaOH in 10 ml of water was then added to the reaction mixture and the mixture was heated for further 15 minutes. After decomposition of the excess reagent with AcOH , the reaction mixture was diluted with water, made alkaline with NH_4OH and extracted with CHCl_3 . The chloroform extract was dried over anhydr. K_2CO_3 and the solvent was evaporated. The residue which upon thin-layer chromatography showed virtually two spots, was dissolved in CHCl_3 and the chloroform solution was chromatographed on alumina. Elution with CHCl_3 gave 25 mg of colorless pillars, mp 244–245°, whose IR spectrum was identical with that of serratinine (I). Further elution with ethyl acetate afforded 63.6 mg of crystalline mass which was recrystallized from ethyl acetate

to give colorless needles, 8-episerratinine (XII), mp 234–237°, $[\alpha]_D^{25} + 27.8^\circ$ ($c=1.0$, CHCl_3); $[\alpha]_D^{25} + 22.43^\circ$ ($c=1.11$, EtOH). *Anal.* Calcd. for $\text{C}_{16}\text{H}_{25}\text{O}_3\text{N}$: C, 68.78; H, 9.02. Found: C, 68.77; H, 8.95. IR cm^{-1} : $\nu_{\text{O-H}}$ 3520, 3340; $\nu_{\text{C=O}}$ 1737.

Oxidation of 8-Episerratinine (XII) to Bisdehydroserattinine* (XIII)—To a stirred solution of 45 mg of 8-episerratinine (XII) in 5 ml of acetone was added 1 ml of Jones' reagent under cooling. The mixture was stirred at 40° for 1 hr. After decomposition of the excess reagent with MeOH, the solution was concentrated to a half volume, diluted with water and extracted with ether. The extract was dried over anhydr. K_2CO_3 and the solvent was evaporated. Distillation of the residue at 160° (2 mm Hg) and recrystallization of the distillate from ether gave 23 mg of colorless prisms, mp 158–160°, whose IR spectrum was identical with that of bisdehydroserattinine* (XIII).

Monoacetyl-8-episerratinine (XIV)—A solution of 103 mg of 8-episerratinine (XII) in 1.5 ml of pyridine containing 1 ml of acetic anhydride was heated at 100° for 4 hr, cooled and evaporated to dryness *in vacuo*. The residue was taken up in water, made alkaline with NH_4OH and extracted with CHCl_3 . The chloroform extract was dried over anhydr. K_2CO_3 and the solvent was evaporated. Recrystallization of the residue from acetone-ether gave 85 mg of colorless pillars (XIV), mp 220.5–221°. *Anal.* Calcd. for $\text{C}_{18}\text{H}_{27}\text{O}_4\text{N}$: C, 67.26; H, 8.47. Found: C, 66.83; H, 8.70. IR cm^{-1} : $\nu_{\text{O-H}}$ 3270; $\nu_{\text{C=O}}$ 1747, 1732; $\nu_{\text{C-O}}$ 1236. NMR τ : 5.17 (1H, clean q., $J_1=5.5$ cps, $J_2=11$ cps, $>\text{CH-OAc}$); 6.48 (1H, m., $>\text{CH-OH}$); 7.96 (3H, s., $-\text{OCOCH}_3$); 9.07 (3H, d., $J=6$ cps, $>\text{CH-CH}_3$).

8-Acetyl-13-dehydro-8-episerratinine (XV)—To a cooled solution of 25 mg of 8-acetyl-8-episerratinine (XIV) in 3 ml of acetone was added 0.3 ml of Jones' reagent with stirring and stirring was continued for further 30 minutes at 40°. After decomposition of excess reagent with MeOH, the solution was evaporated to dryness *in vacuo*. The residue was dissolved in water, made alkaline with NH_4OH and extracted with ether. The ethereal extract was dried over anhydr. K_2CO_3 and the solvent was evaporated. The residue was distilled at 165° (2 mm Hg) to give an oily substance. Recrystallization of the distillate for *n*-hexane-ether gave 8 mg of colorless prisms (XV), mp 169–170°. *Anal.* Calcd. for $\text{C}_{18}\text{H}_{25}\text{O}_4\text{N}$: C, 67.69; H, 7.89. Found: C, 67.52; H, 8.06. IR cm^{-1} : $\nu_{\text{C=O}}$ 1740, 1730, 1705; $\nu_{\text{C-O}}$ 1235. NMR τ : 4.79 (1H, q., $J_1=5.5$ cps, $J_2=10.0$ cps, $>\text{CH-OAc}$); 7.92 (3H, s., $-\text{OCOCH}_3$); 8.93 (3H, d., $J=6$ cps, $>\text{CH-CH}_3$).

β -Dihydroserattinine (XIX)—To a solution of 100 mg of serratinine (I) in 20 ml of *n*-propanol was added 100 mg of metal sodium in small portions under reflux. After sodium had been completely dissolved, the reaction mixture was evaporated to dryness *in vacuo*, diluted with water and extracted with CHCl_3 . The extract was dried over anhydr. K_2CO_3 and the solvent was evaporated. Recrystallization of the residue from methanol-benzene gave 93 mg of colorless prisms, (XIX), mp 210–211°. *Anal.* Calcd. for $\text{C}_{16}\text{H}_{27}\text{O}_3\text{N}$: C, 68.29; H, 9.67. Found: C, 68.11; H, 9.80. IR cm^{-1} : $\nu_{\text{O-H}}$ 3410, 3289. β -Dihydroserattinine methiodide was prepared as usual in acetone and recrystallized from methanol to give colorless leaflets, mp 256–257°. *Anal.* Calcd. for $\text{C}_{16}\text{H}_{27}\text{O}_3\text{N} \cdot \text{CH}_3\text{I}$: C, 48.23; H, 7.14. Found: C, 48.48; H, 7.13. IR cm^{-1} : $\nu_{\text{O-H}}$ 3360. Oxidation of β -dihydroserattinine (XIX) with Jones' reagent gave bisdehydroserattinine* (XIII).

β -Dihydroserattinine Methine (XX)—A solution of 100 mg of β -dihydroserattinine methiodide in 5 ml of MeOH was stirred with freshly precipitated silver hydroxide at room temperature for 20 minutes, filtered and evaporated to dryness *in vacuo*. The residue was pyrolyzed at 160° to give an oily distillate which was soluble in ether. Recrystallization of the distillate from ether gave 35 mg of colorless prisms, (XX), mp 174–175°. Occasionally, this material crystallized as colorless prisms which melted at 158–159°. These two different crystals were shown to be dimorphisms by cross seeding methods and comparison of their IR spectra in solution. *Anal.* Calcd. for $\text{C}_{17}\text{H}_{29}\text{O}_3\text{N}$: C, 69.11; H, 9.90. Found: C, 69.37; H, 9.89. IR cm^{-1} : $\nu_{\text{O-H}}$ 3401; $\nu_{\text{C=O}}$ 1709 (Nujol); $\nu_{\text{O-H}}$ 3584, 3472; $\nu_{\text{C=O}}$ 1719 (CHCl_3). NMR τ : 6.14 (2H, m., $>\text{CH-OH}$); 7.73 (3H, s., $>\text{N-CH}_3$); 9.02 (3H, d., $J=7$ cps, $>\text{CH-CH}_3$).

Methiodide—Prepared as usual in acetone and obtained as colorless prisms, mp 285°, by recrystallization from MeOH-acetone. *Anal.* Calcd. for $\text{C}_{17}\text{H}_{29}\text{O}_3\text{N} \cdot \text{CH}_3\text{I}$: C, 49.43; H, 7.38. Found: C, 49.68; H, 7.33.

Reduction of 8-Acetyl-13-dehydroserattinine (XXI) with NaBH_4 —A solution of 130 mg of 8-acetyl-13-dehydroserattinine (XXI) and 700 mg of NaBH_4 in 20 ml of MeOH was heated for 3 hr under reflux. After addition of 10 ml of aq. 10% NaOH solution, the mixture was heated for further 1 hr and decomposition of the excess hydride was effected by addition of AcOH. The solution was then diluted with water, made alkaline with NH_4OH and extracted with CHCl_3 . The chloroform extract was dried over anhydr. K_2CO_3 . Removal of the solvent furnished crystals which upon thin-layer chromatography showed two spots. The mixture in CHCl_3 was chromatographed on alumina and elution with CHCl_3 gave 75 mg of serratinine (I), mp 244–245°, which was identified with an authentic sample of serratinine by mixed melting point determination and infrared spectra comparison. Successive elution of chromatographic column with MeOH- CHCl_3 (1:1) gave 15 mg of colorless plates, mp 272–276°, which were identified with α -dihydroserattinine** (XVIII) by comparison of infrared spectra and mixed melting point determination.

5-Deoxo-13-deoxyserratinine (XXII)—A solution of 0.2 g of sodium in 10 ml of triethyleneglycol was heated at 180° and completely anhydrous hydrazine was distilled into this solution until it refluxed freely at 180°. To the cooled solution obtained above was added 100 mg of 8-acetyl-13-dehydroserattinine* (XXI) and the mixture was refluxed at 180° for 16 hr. Excess hydrazine was distilled off from the reaction

mixture until the temperature of the solution had raised to 220° and refluxing was continued for further 24 hr. The cooled solution was diluted with water and extracted with CHCl_3 . After drying over anhydr. K_2CO_3 , the solvent was evaporated to left solid mass which was chromatographed over chloroform-alumina. Elution with CHCl_3 gave 65 mg of crystals which were recrystallized from *n*-hexane to afford colorless needles (XXII), mp 244–245°, $[\alpha]_D^{25} -31.6^\circ$ ($c=0.57$, EtOH). *Anal.* Calcd. for $\text{C}_{16}\text{H}_{25}\text{ON}$: C, 77.06; H, 10.91. Found: C, 77.29; H, 10.94. IR cm^{-1} : $\nu_{\text{O-H}}$ 3155.

8-Dehydro-5-deoxo-13-deoxyserratinine (XXIII)—A solution of 50 mg of 5-deoxo-13-deoxyserratinine (XXII) in 2 ml of aq. 90% AcOH was added dropwise to a stirred solution of 25 mg of chromium trioxide in 0.7 ml of aq. 90% AcOH at room temperature and the mixture was stirred at room temperature for 15 hr. After decomposition of excess reagent with MeOH, the solution was evaporated to dryness *in vacuo*. The residue was taken up in water, made alkaline with NH_4OH and extracted with CHCl_3 . The extract was dried over anhydr. K_2CO_3 and the solvent was evaporated. The residue in benzene was chromatographed on alumina. Elution with benzene gave colorless oil which was distilled at 90° (0.014 mm Hg) to afford 37 mg of colorless plates (XXIII), mp 81–85°; $[\alpha]_D^{25} +71.9^\circ$ ($c=1.64$, EtOH); $[\phi]_{273} -1023^\circ$ (trough), $[\phi]_{308} +3735^\circ$ (peak) (RD in MeOH). *Anal.* Calcd. for $\text{C}_{16}\text{H}_{25}\text{ON}$: C, 77.68; H, 10.19. Found: C, 77.52; H, 10.13. IR cm^{-1} : $\nu_{\text{C=O}}$ 1704.

8-Benzoyl-8-episerratinine (XXIV)—To a solution of 150 mg of 8-episerratinine (IX) in 6 ml of pyridine was added 2 ml of benzoyl chloride. The mixture was allowed to stand at room temperature overnight, poured into ice-water, made alkaline with NH_4OH and extracted with ether. The ethereal extracts were shaken with aq. 5% HCl solution and the acidic solution was then made alkaline with NH_4OH , and extracted with ether again. The ethereal extract was dried over anhydr. K_2CO_3 and the solvent was evaporated. The residue in benzene was chromatographed on alumina. Elution with benzene gave 100 mg of crystals which were recrystallized from ether to give colorless pillars, (XXIV), mp 196–197°, $[\alpha]_D^{25} +55.3^\circ$ ($c=0.96$, CHCl_3); $[\alpha]_D^{32} +65.9^\circ$ ($c=0.954$, EtOH). *Anal.* Calcd. for $\text{C}_{23}\text{H}_{29}\text{O}_4\text{N}$: C, 72.03; H, 7.62. Found: C, 71.87; H, 7.89. IR cm^{-1} : $\nu_{\text{O-H}}$ 3520, 3150; $\nu_{\text{C=O}}$ 1740, 1694; ν_{aromatic} 1600, 1578. NMR τ : 1.8–2 (2H, m., aromatic protons); 2.3–2.6 (3H, m., aromatic protons); 4.87 (1H, q., $J_1=12$ cps, $J_2=6$ cps, $>\text{CH}-\text{OCO}\phi$); 6.40 (1H, m., $>\text{CH}-\text{OH}$); 8.98 (3H, d., $J=6$ cps, $>\text{CH}-\text{CH}_3$).

Acknowledgement We are grateful to Miss Y. Mano, Kyoto University, for elementary analyses and to Dr. T. Shingu, Kyoto University, for NMR spectral measurements.