cm⁻¹: 1773, τ 9.14 (C₁₀-Me), 8.74 (C₄-Me), by the treatment (reflux, *p*-TsOH in benzene). Unlike the other 6 β lactones (XIII) and (XIV), the lactone (XX) was so stable that it could be recrystallized and could not be epimerized under the aforementioned thermal condition.

Consequently, synthesis of the 6 α -hydroxy-hexahydro series (*trans* B/C-ring fusion) was performed by the other way. Catalytic hydrogenation (Pd-C, MeOH) of 6 α -hydroxy-tetrahydro-ester (X) afforded oily product, which was chromatographed (Al₂O₃) to convert to a lactone (XXI) (*trans* B/C-ring fusion), C₁₇H₂₆O₂, mp 86—87°, $\nu_{\max}^{\text{KBr}}$ cm⁻¹: 1758, τ 9.06 (C₁₀-Me), 8.73 (C₄-Me). The lactone (XXI) is distinguishable from the 6 β lactone (XX) and the other isomeric lactones (XIV and VII) (*cis* B/C-ring fusion). Alkaline hydrolysis (KOH-EtOH-H₂O) of the lactone (XXI) gave the corresponding 6 α -hydroxy acid (XXII), C₁₇H₂₈O₃, mp 227.5—228.5°, which was readily returned (reflux, 10% HCl aq. MeOH) to the original lactone (XXI) and was methylated (CH₂N₂) to the corresponding ester (XXIII), C₁₈H₃₀O₃, mp 142.5—144.5°. Further evidence on C₆-configuration of both the lactones (XX and XXI) was adduced by their reduction (LiAlH₄) to give the respective diol (XXIV), C₁₇H₃₀O₂, mp 141.5—143° and (XXV), C₁₇H₃₀O₂, mp 157.5—158.5°, which were also synthesized from the ester (XXVIII) and (XXIII) having reliable C₆-configuration, respectively.

In conclusion, it is clarified that the lactonization mode is very variable depending on the structure of B/C-ring fusion. A quantitative study on the epimerization is necessary and now is in progress.

Rikagaku Kenkyusho (The Institute of Physical
and Chemical Research),
Yamato-machi, Saitama

AKIRA TAHARA
YASUO OHTSUKA

Received February 18, 1969

[Chem. Pharm. Bull.]
17(7)1531—1533(1969)

UDC 547.458.02 : 576.8.095.3

Structural Feature of *Pneumococcus* Type XIX Specific Polysaccharide¹⁾

Some preliminary analytical data of *Pneumococcus* Type XIX specific polysaccharide have been given by Brown,²⁾ Levine, *et al.*³⁾ and Baddiley, *et al.*,⁴⁾ little known about its components or that of some serological cross-reactivities of a type XIX antiserum.⁵⁾

The present communication is concerned with structural feature of a fragment which is considered to be a major unit in the polysaccharide.

Crude type specific material was fractionated by Cetavlon treatment,⁶⁾ DEAE-cellulose column chromatography using borate,⁷⁾ followed by gel-filtration (Sephadex G-100). During these treatments, particularly DEAE-cellulose treatment with sodium borate caused a remarkable fragmentation and serological activity of the material to the antiserum decreased.

- 1) A part of this work was presented at the Winter Meeting of the American Chemical Society, Phoenix, Ariz., January, 1966.
- 2) R. Brown, *J. Immunol.*, **37**, 445 (1939); *ibid.*, **47**, 7 (1943).
- 3) S. Levine, H.J.R. Stevenson and P.W. Kabler, *Arch. Biochem. Biophys.*, **45**, 65 (1953).
- 4) Z.A. Shabarova, J.G. Buchanan and J. Baddiley, *Biochim. Biophys. Acta*, **57**, 146 (1962).
- 5) M. Heidelberger, "Fortschritte der Organischer Naturstoffe," Vol. 18, Springer-Verlag in Vienna, 1960, p. 503.
- 6) J.E. Scott, *Chem. Ind. (London)*, **1955**, 1568.
- 7) H. Neukon, H. Deuel, W.J. Heri and W. Kündig, *Helv. Chim. Acta*, **43**, 64 (1960).

After gel-filtration, a serologically active fraction was re-fractionated by Cetavlon followed by ethanol precipitation.

A main constituent thus obtained, $[\alpha]_D +60^\circ$ ($c=0.3$, H_2O), electrophoretically pure, gave L-rhamnose, D-glucose and phosphate in an approximate ratio 4:2:2 by acid hydrolysis. Quantitative estimation of the component sugars of the material by means of the colorimetric method using cysteine-sulfuric acid⁸⁾ also showed a ratio of rhamnose: glucose of 2:1. A titration curve was similar with that of a dibasic acid derivative. On the other hand, treatment with Dowex 50 (H) resin of the material in the cold produced a fragmentation and changed to serologically inactive (unable to give a precipitation band with the antiserum by the procedure of Ouchterlony⁹⁾).

Periodate oxidation of the material led to the consumption of approximately 5.2 moles of periodate per phosphorylated hexasaccharide unit, and the productions of 2.1 moles of formic acid, 1.68 mole of acetaldehyde per unit and no formaldehyde. Reduction by sodium borohydride of the oxidation product followed by hydrolysis with acidic ion-exchange resin in the cold gave glycerol, 1,2-propandiol and unidentified phosphate, and on further hydrolysis (with 2N hydrochloric acid, at 100°), L-rhamnose, D-glucose, erythritol and a small amount of D-glucose 6-phosphate in addition. The molar ratio of liberated rhamnose: glucose was 1.5:1 approximately.

From these results it appears that: 1) reducing end group of the material is rhamnose because of the productions of acetaldehyde and formic acid, 2) some of the rhamnose residues are presented as 1→2 linkage which should be produced of 1,2-propandiol, 3) other part of rhamnose residues did not contain hydroxyl groups on adjacent carbon atoms, 4) some of the glucose residues have hydroxyl groups on adjacent carbon atoms C-2 and C-3 which is erythritol-producible linkage, 5) some of the glucose residues have no adjacent hydroxyl groups, 6) a part of the glucose residues are substituted at C-6 by phosphate.

Phosphate ester in the material was presented as a mixture of mono- and diester type linkages. An alkaline titration curve of the material showed the presence of typical dibasic acid. Treatment of the material with alkaline phosphatase liberated slowly, about half of the phosphate content, though using of acid phosphatase liberated less phosphate. The enzyme-treated material after removal of inorganic phosphate then titrated as a monobasic acid showing the presence of diester type phosphate.

Treatment of the starting material with 1N sodium hydroxide (7 days, under nitrogen, at room temperature), sodium borohydride (16 hr, at room temp.) and followed by Dowex 50 (H) resin gave a dialyzable oligosaccharide as a main product, which contained L-rhamnose, D-glucose, L-rhamnitol and phosphate in the same ratio. The oligosaccharide was paper chromatographically and electrophoretically pure and showed $[\alpha]_D +70^\circ$ ($c=0.2$, H_2O). Its titration curve showed the presence of dibasic acid. Periodate oxidation study of the oligosaccharide showed that consumption of 2.28 moles of periodate, productions of 0.95 mole of formic acid and 1.23 mole of acetaldehyde per phosphorylated trisaccharide unit.

The derived oligosaccharide I, therefore, appears to be one of the repeating unit which may be joined through phosphoric acid residues to other sugar units II, III and IV thus yielding a polymer.

The most probable structures of these sugar units in the polysaccharide will be as follows:

8) Z. Dische and L.B. Shettles, *J. Biol. Chem.*, **175**, 595 (1948).

9) Ö. Ouchterlony, *Acta Pathol. Microbiol. Scand.*, **26**, 507 (1949).

The recognition of amino sugar and total structure of the type specific polysaccharide will be reported in the near future.

Acknowledgement Most of the work has been done at Queen's University. The authors thank to Dr. Rachael Brown and Prof. M. Heidelberger for gift of the type specific material and to the National Research Council of Canada for financial support of the work.

*Tokyo College of Pharmacy,
600 Kashiwagi 4-chome,
Shinjuku-ku, Tokyo*

*Dept. of Chemistry, Queen's University,
Kingston, Ontario, Canada*

TOSHIO MIYAZAKI

J.K.N. JONES

Received March 28, 1969