[Chem. Pharm. Bull.
18(12)2590-2591 (1970)]

UDC 547.597.02 : 547.991.02 : 581.192 : 582.752.1

Bitter Principle of *Picrasma ailanthoides* PLANCHON. Nigakihemiacetal C

A number of bitter principles have been isolated from *Picrasma ailanthoides* PLANCHON (= *P. quassioides* BENNETT), and their structures have been determined.¹⁻⁵ In addition to the previously reported nigakilactones A, B, C, D (=quassin),¹ E, F,³ G (=picrasin A⁴) and H⁵ along with nigakihemiacetals A and B (=neoquassin),⁵ we have recently isolated a new bitter substance, nigakihemiacetal C. The evidence establishing structure I (R=OH, H; R'=H) for nigakihemiacetal C is reported in the present paper.

Nigakihemiacetal C⁶ crystallized as colourless needles from ethanol, with mp 265—265.5° and $[\alpha]_D^{25} +49^\circ$ (MeOH). The mass spectrum indicates the formula C₂₁H₃₂O₆ (M⁺ at *m/e* 380). The infrared (IR) ($\nu_{\text{max}}^{\text{Nujol}}$ 3520, 3280, 3200, 1658, 1634 cm⁻¹) and ultraviolet (UV) ($\lambda_{\text{max}}^{\text{MeOH}}$ 272 nm, ϵ 4600) spectra reveal characteristic absorptions for hydroxyl group and an α,β -unsaturated ketone. No absorption was observed in the region which would be expected for a lactone grouping. The proton magnetic resonance (PMR) spectrum of nigakihemiacetal C showing the presence of two secondary and two tertiary methyls, a methoxyl group and an olefinic proton, is closely related to the spectrum of known nigakilactone A,¹ I (R=O, R'=H) (Table I).

TABLE I. PMR Spectral Data (δ in ppm, in CDCl₃)

Compounds	<i>sec</i> -CH ₃	<i>t</i> -CH ₃	-O-CH ₃	$\begin{array}{c} \text{C} \\ \\ \text{C}-\text{CH}-\text{O}- \\ (7) \end{array}$	C=CH
Nigakihemiacetal C	1.05 d <i>J</i> =7 1.13 d <i>J</i> =6	1.16 s 1.45 s	3.53 s		5.48 d <i>J</i> =2
Nigakilactone A	1.01 d <i>J</i> =6.5 1.10 d <i>J</i> =6	1.24 s 1.42 s	3.54 s	4.10 m	5.35 d <i>J</i> =2.5

Nigakihemiacetal C, when treated with Ac₂O-pyridine at room temperature, afforded a diacetate (C₂₅H₃₆O₈, M⁺ at *m/e* 464, $\nu_{\text{max}}^{\text{Nujol}}$ 3450, 1740, 1672, 1630, 1240 cm⁻¹; $\lambda_{\text{max}}^{\text{MeOH}}$ 272 nm, ϵ 5300), which still shows hydroxyl bands in its IR spectrum. In the PMR spectrum (in CDCl₃), signals due to proton on acetoxy-bearing carbon appear at δ 4.95 (1H), δ 5.72 (0.5H) and δ 6.33 (0.5H).

These observations suggest that nigakihemiacetal C is a hemiacetal, I (R=OH, H; R'=H), related to nigakilactone

- 1) T. Murae, T. Tsuyuki, T. Nishihama, S. Masuda and T. Takahashi, *Tetrahedron Letters*, **1969**, 3013.
- 2) H. Hikino, T. Ohta and T. Takemoto, *Chem. Pharm. Bull.* (Tokyo), **18**, 219 (1970).
- 3) T. Murae, T. Ikeda, T. Tsuyuki, T. Nishihama and T. Takahashi, *Bull. Chem. Soc. Japan*, **43**, 969 (1970).
- 4) H. Hikino, T. Ohta and T. Takemoto, *Chem. Pharm. Bull.* (Tokyo), **18**, 1082 (1970).
- 5) T. Murae, T. Ikeda, T. Tsuyuki, T. Nishihama and T. Takahashi, *Bull. Chem. Soc. Japan*, **43**, 3201 (1970).
- 6) The hemiacetals of this type exist as mixture of isomers at C₁₆. Separation on preparative thin-layer chromatogram followed by elution of each component with acetone resulted in formation of the same mixture.

A. Indeed, nigakihemiacetal C gave nigakilactone A,¹⁾ I (R=O, R'=H), on oxidation with Ag₂O. Thus, nigakihemiacetal C should be represented by I (R=OH, H; R'=H). The presence of hydrogen bond between carbonyl at C₁ and hydroxyl group at C₁₁ leads to structure I (R=OAc, H; R'=Ac) for the diacetate.

Spectral data of nigakilactone I, mp 255—256°, C₂₁H₂₈O₆ (M⁺ at *m/e* 376), suggest that this bitter substance would be identical with picrasin B.²⁾ This was confirmed by direct comparison with the authentic specimen.²⁾

Acknowledgement We wish to thank Professors T. Takemoto and H. Hikino, Tohoku University, for a generous gift of the authentic sample of picrasin B together with its spectral data.

Department of Chemistry,
Faculty of Science,
The University of Tokyo
Bunkyo-ku, Tokyo

TATSUSHI MURAE
TÔRU IKEDA
TAKAHIKO TSUYUKI
TADAAKI NISHIHAMA
TAKEYOSHI TAKAHASHI

Received September 7, 1970

[Chem. Pharm. Bull.
18(12):2591—2593(1970)]

UDC 547.94.07 : 581.192 : 582.675.1

Total Synthesis of Optically Active Natural O-Methylthalicberine

O-Methylthalicberine is the first bisbenzylisoquinoline alkaloid isolated from a Ranunculaceae plant, that is, *Thalictrum Thunbergii* DC (Japanese name "Akikaramatsu") by Fujita and Tomimatsu.¹⁾ Its structure²⁾ and absolute configuration^{2,3)} has been clarified and its unique 19-membered macro-ring system was the first example⁴⁾ in the bisbenzylisoquinoline alkaloids. We describe here the total synthesis of optically active natural O-methylthalicberine, which confirms its structure and absolute configuration **1b**.

An important intermediate, (S)-(+)-O-benzyl-8-bromo-N-nor-laudanidine (**2a**), mp 98—99°, [α]_D +38.7° (CHCl₃), was yielded by the optical resolution of the synthetic racemate of **2a** via its (—)-tartaric acid salt. N-Methylation of the resolved base **2a** gave (+)-O-benzyl-8-bromolaudanidine (**2b**). The structure and optical purity of the compounds **2a** and **2b** was assured by the comparison of the melting point and the specific rotation with those of their enantiomers which have been synthesized by Inubushi, *et al.*,⁵⁾ respectively.

The Ullmann condensation of the base **2b** with N-*t*-butoxycarbonyl-3-hydroxy-4-methoxyphenethyl amine (**3**), mp 101—102°, C₁₄H₂₁O₄N, synthesized from O-benzylisovanillin via a usual pathway afforded the compound **4a** as an oily product⁶⁾ in 39% yield. Hydrogenolysis

1) E. Fujita and T. Tomimatsu, *Yakugaku Zasshi*, **79**, 1256 (1959).

2) E. Fujita and T. Tomimatsu, *Yakugaku Zasshi*, **79**, 1260 (1959); T. Tomimatsu, *ibid.*, **79**, 1386 (1959); E. Fujita, T. Tomimatsu, and Y. Kano, *ibid.*, **80**, 1137 (1960); T. Tomimatsu and Y. Kano, *ibid.*, **83**, 153 (1963); E. Fujita, T. Tomimatsu, and Y. Kano, *ibid.*, **83**, 159 (1963); E. Fujita, K. Fuji, and T. Suzuki, *Bull. Inst. Chem. Res., Kyoto Univ.*, **43**, 449 (1965).

3) M. Tomita and J. Kunitomo, *Yakugaku Zasshi*, **82**, 741 (1962).

4) For the similar system found later, see a) N.M. Mollov, H.B. Dutschewska, and H.G. Kirjakov, *Chem. Ind. (London)*, **1965**, 1595; b) M.R. Falco, J.X. de Vries, A.G. de Brovotto, Z. Macció, S. Rebuffo, and I.R.C. Bick, *Tetrahedron Letters*, **1968**, 1953.

5) Y. Inubushi, Y. Masaki, S. Matsumoto, and F. Takami, *Tetrahedron Letters*, **1968**, 3399; *idem*, *J. Chem. Soc. (C)*, **1969**, 1547.

6) The infrared (IR) and nuclear magnetic resonance (NMR) spectra of the amorphous or oily substance marked with this symbol showed reasonable patterns for the corresponding structure to each compound and homogeneity was also verified by thin-layer chromatography and NMR spectrum measurements.