Usnic Acid. VII.¹⁾ The Pyrolysis of Methyl Dihydrous Usnic Acid

KÔTARO TAKAHASHI and MASAKO TAKANI

Faculty of Pharmaceutical Sciences, University of Kanazawa²⁾

(Received February 20, 1970)

- 1) The structures of two pyrolysis products of methyl dihydrous usnic acid (I) were established to be II and III, respectively.
- 2) The mechanism of the formation of II and III from I were discussed.
- 3) The mass spectra of V and isoanhydromethyl dihydrous usnic acid monoacetate, and X and anhydromethyl dihydrous usnic acid monoacetate were studied.

The structural formula of usnic acid³⁾ was established, and recently the structure of isousnic acid was elucidated by Shibata and Taguchi.⁴⁾ Although some explanations³⁾ have been presented for the remarkable difference between usnic- and dihydrous usnic acid in the degradation reactions, any unequivocal experimental proof has not been provided as yet. In a previous paper,⁵⁾ the author reported that 3-methyl-2,6-dihydroxyacetophenone, a resorcinol type of compound, but not 6-hydroxy-3,5-dimethyl-7-acetylcoumaranone-2³⁾ was obtained from methyl dihydrous usnic acid (I) by the potassium permanganate oxidation, followed by the vacuum distillation, to suggest that it was derived from the ring B of I. The present paper deals with the pyrolysis products of I. On heating at 220–240° for 30 min under ordinary pressure, I liberated acetic acid and acetylacetone. The residue afforded on subsequent distillation at 220–240° under diminished pressure (1 mmHg) faint yellow crystals $C_{19}H_{18}O_6$ (II), mp 243° and yellow crystals $C_{17}H_{16}O_5$ (III), mp 232–233°, which might have been resulted from I by the loss of H_2O , and H_2O and $COCH_3$, respectively. I gave also II and III by the distillation with $CaCl_2$, but no coumaranone type of compound such as 6-hydroxy-3,5-dimethyl-7-acetylcoumaranone-2 was obtained. II gave violet brown coloration with $FeCl_3$ and afforded bisoxime monoanhydride $C_{19}H_{18}O_5N_2$ (IV), mp 287°, by the action of hydroxylamine hydrochloride, suggesting the presence of a carbonyl group located at the *ortho* position of an enolic OH group. By the acetylation with acetic anhydride and conc. H_2SO_4 or with acetic anhydride and sodium acetate, II gave monoacetate $C_{21}H_{20}O_7$ (V), mp 128°, which gave by the ozonolysis, $C_{12}H_{12}O_6$ (VI), mp 170–172°. VI gave methyl ester $C_{13}H_{14}O_6$ (VII), mp 123–125°, with diazomethane, which was deacetylated to $C_{11}H_{12}O_5$ (VIII), mp 102–103°. VI gave 3-methyl-2,6-dihydroxyacetophenone,⁵⁾ $C_9H_{10}O_3$ (IX), mp 134–136°, by the hydrolysis with 10% ethanolic KOH, followed by decarboxylation. Monoacetate (V) was deacetylated to II with conc. H_2SO_4 . III gave a red coloration with $FeCl_3$ and afforded diacetate $C_{21}H_{20}O_7$ (X), mp 165–166°, with acetic anhydride and conc. H_2SO_4 or with acetic anhydride and sodium acetate. X gave VI by the ozonolysis and monoacetate $C_{19}H_{18}O_6$ (XI), mp 218–219°, by the ethanolysis. X was deacetylated to III with conc. H_2SO_4 . These chemical evidences mentioned above suggest that the aromatisation is effected in the course of the pyrolysis, not in the course of the acetylation by the Dienone-Phenol rearrangement,⁶⁾ and that the pyrolysis products II and III possess a 3-methyl-2,6-dihydroxyacetophenone structure in the molecules (Chart 1).

1) Part VI: K. Takahashi and S. Miyashita, *Chem. Pharm. Bull.* (Tokyo), **16**, 988 (1968).

2) Location: *Takaramachi, Kanazawa*.

3) Y. Asahina and S. Shibata "The Chemistry of Lichen Substances", Japan Society for the Promotion of Science, Tokyo, Japan, 1954, p. 190.

4) S. Shibata and H. Taguchi, *Tetrahedron Letters*, **1967**, 4867.

5) K. Takahashi and S. Miyashita, *Chem. Pharm. Bull.* (Tokyo), **10**, 603 (1962).

6) K. Takahashi and S. Miyashita, *Chem. Pharm. Bull.* (Tokyo), **11**, 209 (1963).

Chart 1

The structure of $C_{12}H_{12}O_6$ (VI) was elucidated as follows: VI gave violet coloration with $FeCl_3$ and exhibited infrared (IR) bands (cm^{-1}) at 1770 (acetate), 1655, 1650 (chelated $COOH$ and aryl- $COCH_3$). VII showed IR bands (cm^{-1}) at 1755 (acetate), 1690 (chelated $COOCH_3$) and 1675 (aryl- $COCH_3$). VI, VII and VIII were negative to the Gibbs reaction indicating that the *para* positions to OH groups are substituted. Therefore, it is reasonable to assume that VI, from which IX was derived, is methylresacetophenone carboxylic acid monoacetate and VII is methyl methylresacetophenone carboxylate monoacetate. There are three possible structures a, b and c for VII.

Now, by the comparison of τ -values of aryl-CH₃ (7.76) and aryl-H (3.75) of A⁶ with those of aryl-CH₃ (7.54) and aryl-H (3.75) of B,⁶ the following conclusion could be deduced that the COOCH₃ group causes the diamagnetic shift of the methyl group at *ortho* position (7.76 to 7.54), but does not effect the chemical shift of the aryl-H at *meta* position. VII and IX show the aryl-CH₃ signals at nearly equal τ -values (7.89 and 7.82), but VII shows the signal of the aryl-H at lower field (2.24) than those of the aryl-H of IX to suggest that VII has a COOCH₃ group at *meta* position to the CH₃ group and a proton at *ortho* position to the COOCH₃ group (Table I). These nuclear magnetic resonance (NMR) and IR spectral data indicate that VII would be a.

TABLE I. The NMR Data (τ -Value in CDCl₃, 60 Mc)

	Ar-CH ₃	OCOCH ₃	COCH ₃	COOCH ₃	Ar-H	OH
VII	7.89 (s) 3H	7.70 (s) 3H	7.40 (s) 3H	6.03 (s) 3H	2.24 (s) 1H	-1.19 (s) 1H
(A)	7.76 (s) 3H	—	7.26 (s) 3H	—	3.75 (s) 2H	0.13 (s) 2H
(B)	7.54 (s) 3H	—	7.28 (s) 3H	6.08 (s) 3H	3.75 (s) 1H	-3.58 (s) 1H -3.97 (s) 1H
(K)	7.82 (s) 3H	—	7.23 (s) 3H	—	3.79, 3.65 (d) 1H 2.91, 2.77 (d) 1H	2.38 (s) 1H -1.55 (s) 1H

abbreviation: s; singlet, d; doublet

The ultraviolet (UV) spectrum of II shows absorption maxima ($m\mu$, $\log \epsilon$) at 237.2 (4.24), 274.4 (4.03), 284.0 (4.00, sh) and 328 (3.63), and the IR bands (cm^{-1}) at 1660 ($\alpha\beta\text{-}\alpha'\beta'$ -unsatd. C=O), 1648 and 1635 (two chelated C=O), 1530—1550 (broad, triketone) and 1125, 1018 (—C—O—C—). These spectral data closely resemble those of isoanhydromethyldihydrousnic acid,⁷ with UV maxima⁸ at 238 (4.54), 270 (4.27, sh), 283 (4.32) and 330 (3.95) and the IR bands⁸ at 1657 ($\alpha\beta\text{-}\alpha'\beta'$ -unsatd. C=O), 1650 and 1640 (two chelated C=O) and 1530—1550 (broad, triketone) and 1182, 1070 (—C—O—C—). The NMR of II could be assigned as shown in Table II, with reference to those of isoanhydromethyldihydrousnic acid and anhydromethyldihydro-

7) K. Takahashi, S. Miyashita and Y. Ueda, *Chem. Pharm. Bull.* (Tokyo), **11**, 473 (1963).

8) The UV and IR spectra were newly measured by Hitachi EPS-2U recording spectrophotometer and Nippon Bunko D.S. 402G IR spectrometer, respectively.

usnic acid.⁶⁾ These spectral data suggest that the chromophores of both compounds II and isoanhydromethyldihydrousnic acid are almost same and almost equally situated in the molecules, except the aryl-CH₃ groups.

TABLE II. The NMR Data (τ -Value and δ -Value, 60 Mc)

	$\begin{array}{c} \text{CH}_3 \\ \diagup \\ \text{C} \\ \diagdown \\ \text{CH}_3 \end{array}$	Ar-CH ₃	COCH_3		CH ₂	Ar-H	OH		
			Ring A	Ring B			Chelated	Chelated enol	
II	8.49; 8.27 (d) 6H	7.65 (s) 3H	7.32; 7.16 (d) 3H	7.09 (s) 3H	—	1.96; 1.83 (d) 1H	-3.20 (s) 1H	-7.92; -7.95 (d) 1H	in CDCl ₃ (τ -value)
III	8.55 (s) 6H	7.71 (s) 3H	—	7.16 (s) 3H	5.96 (s) 2H	2.03 (s) 1H	-2.74 (s) 1H	—	
Isoanhydro- methyldihydro- usnic acid ^{a)}	8.40; 8.28 (d) 6H	7.20; 7.14 (d) 3H	7.35; 7.22 (d) 3H	7.17 (s) 3H	—	3.31 (s) 1H	-2.73 (s) 1H	-8.56; -8.88 (d) 1H	
Anhydromethyl- dihydrousnic acid	8.49 (s) 6H	7.20 (s) 3H	7.23 (s) 3H 7.14 (s) 3H	—	—	3.27 (s) 1H	-2.78 (s) 1H	-7.60 (s) 1H	
II	1.83 (s) 6H	2.39 (s) 3H	2.84 (s) 3H 3.07 (s) 3H	—	—	7.99 (s) 1H	—	—	in CF ₃ COOD (δ -value)
Isoanhydro- methyldihydro- usnic acid	1.83 (s) 6H	2.97 (s) 3H	2.89 (s) 3H 3.18 (s) 3H	—	—	7.04 (s) 1H	—	—	
Anhydromethyl- dihydrousnic acid	1.66 (s) 6H	2.88 (s) 3H	2.96 (s) 3H 3.10 (s) 3H	—	—	6.88 (s) 1H	—	—	

a) The NMR spectrum was newly measured by JNM-C-60-H high resolution NMR instrument and some of assignments ⁷⁾ of signals were revised.

II exhibits the NMR signals of gem-dimethyl, COCH₃ group of the ring A, aryl-H and chelated enol OH group as doublets in CDCl₃, but those of the gem-dimethyl, the COCH₃ group of the ring A and the aryl-H as singlets in CF₃COOD and D₂O solution, respectively, thus suggesting that II is an isotype of compound and exists in two types of enol form and is equilibrium between

as isoanhydromethyldihydrousnic acid.

Acetate (V) of II exhibits NMR signals as shown in Table III. V exhibits the signal of a chelated enol OH group, but does not exhibit a signal of a chelated OH group, indicating that the aryl-OH group of II was acetylated to give V. The splitting of the signals of the gem-dimethyl, the COCH₃ group of the ring A, the aryl-H and the chelated enol OH group indicates that V is an isotype of compound and exists in two types of enol form as II.

The IR bands (cm⁻¹) of V, 1770 (acetate), 1687 (aryl-COCH₃), 1662 ($\alpha\beta$ - $\alpha'\beta'$ -unsatd. ring C=O), 1620 (chelated C=O), 1515—1554 (broad, triketone) and 1125, 1018 (-C-O-C-) resemble those of isoanhydromethyldihydrousnic acid monoacetate,⁷⁾ 1767 (acetate), 1688 (aryl-COCH₃), 1658 ($\alpha\beta$ - $\alpha'\beta'$ -unsatd. ring C=O), 1620 (chelated C=O), 1525—1545 (broad, triketone) and 1145, 1055 (-C-O-C-). The UV maxima of V (m μ , log ϵ), 231.3 (4.44), 258.2 (4.22, sh) and 302.8 (4.10) also resemble those of isoanhydromethyldihydrousnic acid mono-

TABLE III. The NMR Data (τ -Value in CDCl_3 , 60 Mc)

V	$\begin{array}{c} \text{CH}_3 \\ \diagup \\ \text{C} \\ \diagdown \\ \text{CH}_3 \end{array}$	Ar-CH ₃	OCOCH ₃	COCH ₃		Ar-H	Chelated enol OH
				Ring A	Ring B		
V	8.42; 8.31 (d) 6H	7.69 (s) 3H	7.62 (s) 3H	7.33; 7.17 (d) 3H	7.23 (s) 3H	1.96; 1.82 (d) 1H	-8.41; -8.43 (d) 1H
Isoanhydromethyl- dihydro- usnic acid monoacetate	8.41; 8.29 (d) 6H	7.18; 7.11 (d) 3H	7.69 (s) 3H	7.38; 7.25 (d) 3H	7.29 (s) 3H	3.20 (s) 1H	-8.47; -8.73 (d) 1H
Anhydromethyl- dihydrousnic acid monoacetate ^{a)}	8.50 (s) 6H	7.16 (s) 3H	7.69 (s) 3H	7.24, (s) 3H 7.33, (s) 3H		3.18 (s) 1H	-7.90 (s) 1H

a) The NMR spectrum was newly measured by JNM-C-60-H high resolution NMR instrument and some of assignments⁹⁾ of signals were revised.

acetate, 234.7 (4.45), 260.7 (4.21, sh) and 304.2 (4.01). These chemical and spectral data indicate that II, V, VI, VII and VIII could be formulated as shown in Chart 1.

$\text{C}_{17}\text{H}_{16}\text{O}_5$ (III) exhibits the IR bands (cm^{-1}) at 1725 (six-membered ring C=O), 1680 ($\alpha\beta$ -unsatd. six-membered ring C=O), 1635 (chelated aryl-COCH₃) and the UV maxima ($m\mu$, log ϵ) at 266.2 (4.38), 282.0 (4.23, sh) and 302.0 (3.81, sh). The NMR signals are shown in Table II. The NMR signals (τ -value) at 8.55 (s, 6H, a gem-dimethyl) and 5.96 (s, 2H, a methylene group which could be considered to be produced by the hydrolytic fission of COCH₃ group of the ring A of I) and 2.03 (s, 1H, aryl-H), and the IR bands indicate that III exists in a diketoform as 2,2,5,5-tetramethylhexane-1,3-dione.⁹⁾ Diacetate (X) of III, which gave VI by the ozonolysis, exhibits the IR bands (cm^{-1}) at 1776 and 1765 (two acetates), 1693 (aryl-COCH₃), 1660 ($\alpha\beta\gamma\delta$ -unsatd. ring C=O) and 1175, 1045 ($-\text{C}-\text{O}-\text{C}-$) and the NMR signals (τ -

Chart 2. The Fragmentations of V ($\text{R}=\text{CH}_3$, $\text{R}'=\text{H}$) and Isoanhydromethyl-dihydrousnic Acid Monoacetate ($\text{R}=\text{H}$, $\text{R}'=\text{CH}_3$)

The figures in round and square brackets are the relative intensities of the fragment ions of the former and the latter, respectively.

Chart 3. The Fragmentations of X (R=CH₃, R'=H) and Anhydromethyldihydrousnic Acid Monoacetate (R=H, R'=CH₃)

The figures in round and square brackets are the relative intensities of the fragment ions of the former and the latter, respectively. Typical broad peaks of metastable ions were observed in the spectrum of X, but not in the spectrum of anhydromethyldihydrousnic acid monoacetate.

TABLE IV

Formula	m/e (Calcd.)	m/e (Found)					
		Species	V	Isoanhydro- methyldihydro- usnic acid monoacetate	Species	X	Anhydro- methyldihydro- usnic acid monoacetate
C ₂₁ H ₂₀ O ₇	384.121	M ⁺	0.122	0.121	M ⁺	0.123	0.124
C ₁₉ H ₁₈ O ₆	342.110	A	0.112	0.109	a	0.111	0.112
C ₁₈ H ₁₅ O ₆	327.087	B	0.086	0.087	c	0.088	0.086
C ₁₇ H ₁₆ O ₅	300.100				b	0.100	0.099
C ₁₇ H ₁₅ O ₅	299.092	C	0.090	0.092	i	0.091	0.090
C ₁₆ H ₁₃ O ₅	285.076				d	0.079	0.076
C ₁₆ H ₁₆ O ₄	272.105				f	0.105	0.104
C ₁₆ H ₁₅ O ₄	271.097	D	0.095	0.102	j	0.095	0.098
C ₁₅ H ₁₄ O ₄	258.089	E	0.092	0.088			
C ₁₅ H ₁₃ O ₄	257.081	J	0.081	0.081	e	0.083	0.083
C ₁₅ H ₁₅ O ₃	243.102				k	0.106	0.103
C ₁₄ H ₁₅ O ₃	231.102	F	0.102	0.103			
C ₁₄ H ₁₄ O ₃	230.094	G	0.095	0.095			
C ₁₄ H ₁₃ O ₃	229.086	K	0.082	0.090	g	0.086	0.083
C ₁₃ H ₁₁ O ₃	215.071	H	0.068	0.071			
C ₁₃ H ₁₃ O ₂	201.092				h	0.092	0.091
C ₁₂ H ₁₁ O ₂	187.076	I	0.074	0.074			
C ₅ H ₃ O ₃	111.008	L	0.007	—

Chart 4

value) at 8.58 (s, 6H, a gem-dimethyl), 7.72 (s, 3H, aryl-CH₃), 7.68 (s, 3H) and 7.64 (s, 3H) (two acetates), 7.24 (s, 3H, aryl-COCH₃), 3.13 (s, 1H, -CH=C<) and 1.98 (s, 1H, aryl-H) in CDCl₃ and the UV maxima ($m\mu$, log ϵ) at 248.0 (4.50), 281 (3.82, sh) and 347 (4.08). The presence of a signal at 3.13 (1H) and the absence of a signal at 5.96 (2H) indicate that X has an enol acetyl group. The alcoholysis product (XI) of X exhibits the IR bands (cm⁻¹) at 1750 (acetate), 1720 (six-membered ring C=O), 1690 (aryl-COCH₃), 1680 ($\alpha\beta$ -unsatd. ring C=O), 1195 and 1070 (-C-O-C-) and the NMR signals (τ -value) at 8.55 (s, 6H, a gem-dimethyl), 7.72 (s, 3H, aryl-CH₃), 7.65 (s, 3H, acetate), 7.28 (s, 3H, COCH₃), 5.95 (s, 2H, -CH₂-) and 1.93 (s, 1H, aryl-H), indicating that the enol acetyl group of the ring A of X was eliminated to give XI by the alcoholysis and that XI exists in a diketofrom as III. These chemical and spectral data indicate that X is a normal type of compound as anhydromethyldihydrousnic acid⁶ and III, X and XI could be formulated as shown in chart 1.

Mass spectral analysis.¹⁰ Analyses of peaks of the spectra of V and isoanhydromethyldihydrousnic acid monoacetate (isotypes of compound) (Chart 2) and those of X and anhydromethyldihydrousnic acid monoacetate (normal types of compound) (Chart 3) were carried out in millimass order (Table IV). Some of the fragmentations and the structures of the fragment-

10) Cf. H. Budzikiewicz, C. Djerassi and D.H. Williams, "Interpretation of Mass Spectra of Organic Compound," Holden-Day Inc. San Francisco, 1964.

ions of these compounds could be assumed as shown in Chart 2 and 3. The fragmentations of species m/e 342 (A) to species m/e 258 (E) by path (i) and species m/e 231 (F) by path (ii) might be characteristic of the isotypes of compound.

The reaction mechanism of the formation of II and III from methyldihydrousnic acid (I) could tentatively be postulated as shown in Chart 4. Namely, the first stage of this reaction is the fission of $-C-O-C-$ bond in the furan ring with production of two hydroxyl groups, of which the OH group in the ring B is then dehydrated to produce a double bond. At the next stage, the OH group in the ring A and the enol OH group in the ring B combine with each other to give $-C-O-C-$ bond in two ways, giving two compounds A and B. A might give II and B, after the hydrolytic fission of $COCH_3$ group in the ring A, might give III through π -complex intermediate C by the Jacobson rearrangement.¹¹⁾ In this case, the effect of the *meta*-directing group ($COCH_3$) and of the *ortho*-directing group (OH) would be considered to cause the 1,3-migration of the angular methyl group, but further investigation on this point should be performed (Chart 4).

Chart 5

It is of interest to assume that methyldihydrousnic acid (I) gives 3-methyl-2,6-dihydroxyacetophenone (IX) either by the potassium permanganate oxidation, followed by the vacuum distillation⁵⁾ or by the vacuum distillation, followed by the ozone oxidation as shown in Chart 5.

Experimental

The IR spectra were measured in KBr pellet by Nippon Bunko DS-402G spectrometer, the UV spectra in EtOH by Hitachi EPS-2U recording spectrometer, the NMR spectra in $CDCl_3$ by JNM-C-60-H high resolution NMR instrument at 60 Mc with $(CH_3)_4Si$ as internal reference and the mass spectra by JMS-01SG mass spectrometer, the ionizing current kept at 200 μA , while the ionizing energy being maintained at 75 eV, not otherwise stated.

Dry-Distillation of Methyldihydrousnic Acid (I)—Three grams of I was heated at 220–240° for 30 min. After removing acetylacetone (a rose coloration with *o*-phenylenediamine) and acetic acid, the residue was distilled *in vacuo* under 1 mmHg pressure to give reddish yellow distillate, which solidified by treatment with methanol. The yellow substance was recrystallised from benzene to give faint yellow crystal of mp 243°. Yield: 100 mg. $FeCl_3$ reaction: violet-brown. R_f : 0.25 (benzene: AcOEt=4:1, Kieselgel G). *Anal.* Calcd. for $C_{19}H_{18}O_6$: C, 66.66; H, 5.30. Found: C, 66.84; H, 5.47. From the mother liquor, yellow needle (III) of mp 232–233° was obtained. Yield: 20 mg. $FeCl_3$ reaction: red. R_f : 0.61 (benzene: AcOEt=4:1, Kieselgel G). *Anal.* Calcd. for $C_{17}H_{16}O_5$: C, 67.99; H, 5.37. Found: C, 67.95; H, 5.40.

11) P. de Mayo, "Molecular Rearrangements," Part I, p. 299.

Dry-Distillation of I with CaCl_2 —I (1.5 g) was mixed with CaCl_2 (1.5 g) and heated at 200° for 20 min. The residue was then distilled under 1 mmHg pressure to give yellow distillate which solidified by treatment with methanol. The yellow substance was chromatographed on Kieselgel H (30 g) with benzene-AcOEt (9:1) as eluent. The first yellow eluate gave crystal of mp 230° , which was proved to be identical with III by the mixed fusion and IR spectra. The second yellow eluate gave crystal of mp $240\text{--}241^\circ$, which was proved to be identical with II by the mixed fusion and IR spectra. *Anal.* Calcd. for $\text{C}_{19}\text{H}_{18}\text{O}_6$: C, 66.66; H, 5.30. Found: C, 66.69; H, 5.31.

Acetylation of II—A mixture of 0.37 g of II, 0.37 g of sodium acetate and 6 ml of acetic anhydride was heated on an oil bath at 160° for 1.5 hr. After cooling, the mixture was poured into ice water to give precipitate, which was crystallised from EtOH to give V of mp 128° as colorless plate. Yield: 140 mg. FeCl_3 reaction: red. *Rf*: 0.35 (benzene: AcOEt=4:1, Kieselgel G). *Anal.* Calcd. for $\text{C}_{21}\text{H}_{20}\text{O}_7$: C, 65.61; H, 5.24. Found: C, 65.47; H, 5.57.

Acetylation of III—A mixture of 0.5 g of III, 0.5 g of sodium acetate and 6 ml of acetic anhydride was warmed on a water bath for 1.5 hr. After cooling, the mixture was poured into ice water and the mixture was allowed to stand overnight. Precipitate was crystallised from EtOH to give X of mp $165\text{--}166^\circ$ as faint yellow needles. Yield: 50 mg. *Rf*: 0.68 (benzene:AcOEt=4:1, Kieselgel G). *Anal.* Calcd. for $\text{C}_{21}\text{H}_{20}\text{O}_7$: C, 65.61; H, 5.24. Found: C, 65.55; H, 5.30.

Acetylation of the Dry-Distillate of I—A mixture of the yellow distillate (3.8 g), obtained from I mentioned above, 38 ml of acetic anhydride and 6 drops of conc. H_2SO_4 was warmed on a water bath for 1.5 hr. After cooling, the mixture was poured into water to give brown precipitate, which was crystallised from EtOH to give crystal of mp $165\text{--}166^\circ$ which was proved to be identical with X by the mixed fusion and IR spectra. Yield: 0.7 g. *Anal.* Calcd. for $\text{C}_{21}\text{H}_{20}\text{O}_7$: C, 65.61; H, 5.24. Found: C, 65.64; H, 5.34. From the mother liquor, crystal of mp 128° was obtained, which was proved to be identical with V by the mixed fusion and IR spectra. Yield: 0.15 g. *Anal.* Calcd. for $\text{C}_{21}\text{H}_{20}\text{O}_7$: C, 65.61; H, 5.24. Found: C, 65.38; H, 5.16.

Deacetylation of V to II—V (97 mg) was dissolved in ice-cold conc. H_2SO_4 (0.5 ml) and the solution was kept in cold for 20 min and poured into ice water to give crystal of mp $241\text{--}242^\circ$ (from benzene), which was proved to be identical with II by the mixed fusion and IR spectra. Yield: 50 mg. *Anal.* Calcd. for $\text{C}_{19}\text{H}_{18}\text{O}_6$: C, 66.66; H, 5.30. Found: C, 66.52; H, 5.30.

Deacetylation of X to III—X (100 mg) was dissolved in ice-cold conc. H_2SO_4 (1 ml) and the solution was allowed to stand in cold for 20 min and poured into ice water to give III of mp 233° . *Anal.* Calcd. for $\text{C}_{17}\text{H}_{16}\text{O}_5$: C, 67.99; H, 5.37. Found: C, 67.66; H, 5.31.

Oximation of II—A mixture of II (200 mg) in EtOH and hydroxylamine hydrochloride (600 mg) and sodium acetate (600 mg) in water was warmed on a water bath for 5 hr and worked up as usual to give colorless crystal of mp 287° (decomp.) (IV) from EtOH. *Anal.* Calcd. for $\text{C}_{19}\text{H}_{18}\text{O}_5\text{N}_2$: C, 64.40; H, 5.12; N, 7.91. Found: C, 64.32; H, 5.32; N, 7.39. IR ν_{max} (cm^{-1}): 1665, 1601, 1545, 1495, 1400, 1297, 1140, 1025, 968, 888.

Ozonolysis of V—V (700 mg) was dissolved in CHCl_3 (15 ml) and ozonized oxygen was passed through for 2 hr under ice cooling. The CHCl_3 solution was then distilled off *in vacuo* and water (15 ml) was added to the residue and the solution was warmed on a water bath at 45° for 30 min and allowed to stand at room temperature for 2 days. The oxidation product was extracted with ether five times and after drying with anhyd. Na_2SO_4 , ether was distilled off to give colorless needle (VI) of mp $170\text{--}172^\circ$ from benzene. Yield: 17 mg. FeCl_3 reaction: violet. *Rf*: 0.33 (benzene: dioxane: AcOH=90:25:4, Kieselgel G). UV λ_{max} ($m\mu$, (log ϵ)): 223 (4.31), 321 (3.75). *Anal.* Calcd. for $\text{C}_{12}\text{H}_{12}\text{O}_6$: C, 57.14; H, 4.80. Found: C, 57.32; H, 4.79.

Methylation of VI—One-tenth gram of VI in ether (20 ml) was methylated with diazomethane to give colorless plate (VII) of mp $123\text{--}125^\circ$ from methanol. FeCl_3 reaction: violet. *Rf*: 0.69 (benzene: AcOEt=4:1, Kieselgel G). UV λ_{max} ($m\mu$, (log ϵ)): 223.2 (4.45), 251.6 (3.89, shoulder), 320 (3.91). *Anal.* Calcd. for $\text{C}_{13}\text{H}_{14}\text{O}_6$: C, 58.64; H, 5.30. Found: C, 58.74; H, 5.39.

Deacetylation of VII—VII (42 mg) was dissolved in ice-cold conc. H_2SO_4 (0.8 ml) and the mixture was treated as mentioned above to give crystal (VIII) of mp $102\text{--}103^\circ$ from methanol. FeCl_3 reaction: violet. UV λ_{max} ($m\mu$, (log ϵ)): 236.4 (4.45), 263.7 (4.33), 281 (4.22, shoulder), 341 (3.70). IR ν_{max} (cm^{-1}): 1670 (chelated COOCH_3), 1628 (chelated COCH_3). *Anal.* Calcd. for $\text{C}_{11}\text{H}_{12}\text{O}_5$: C, 58.92; H, 5.40. Found: C, 58.88; H, 5.43.

Hydrolysis, followed by Decarboxylation of VI—VI (90 mg) was warmed in 10% alcoholic KOH (15 ml) on a water bath for 1 hr and the solution was evaporated *in vacuo* and 3 ml of water was added. After acidification with dil. HCl, the resulting yellow precipitate was heated at 210° . Decarboxylation took place under bubbling. After the end of bubbling, the product was sublimated at 120° under 1 mmHg pressure to give crystal (IX) of mp $134\text{--}136^\circ$, which was proved to be identical with 3-methyl 2,6-dihydroxy acetophenone⁶ by the mixed fusion and UV, IR spectra. FeCl_3 reaction: blue-green. UV λ_{max} ($m\mu$, (log ϵ)): 224.5 (4.00), 273 (4.01), 352 (3.45). *Anal.* Calcd. for $\text{C}_9\text{H}_{10}\text{O}_3$: C, 65.05; H, 6.07. Found: C, 65.11; H, 6.09.

Ozonolysis of X—X (1.2 g) was dissolved in CHCl_3 (30 ml) and ozonised oxygen was passed through for 2.5 hr under ice cooling and the CHCl_3 solution was treated as mentioned above to give colorless needle

of mp 170—172° from benzene, which was proved to be identical with VI by the mixed fusion, UV and IR spectra. *Anal.* Calcd. for $C_{12}H_{12}O_6$: C, 57.14; H, 4.80. Found: C, 57.05; H, 4.71.

Alcoholysis of X—X (0.3 g) in EtOH or MeOH (20 ml) was boiled on a water bath for 45 hr. The solution gave yellow crystal (0.2 g) after cooling, which was crystallised from EtOH to give crystal (XI) of mp 218—219°. UV $\lambda_{\max}$ ($m\mu$, ($\log \epsilon$)): 249.5 (4.40), 257.2 (4.42), 281 (3.87, shoulder), 307 (3.44, shoulder). *Anal.* Calcd. for $C_{19}H_{18}O_6$: C, 66.66; H, 5.30. Found: C, 66.18; H, 5.41. Mass Spectrum (m/e , relative intensity): M^+ 342 (3), 300 (100), 299 (2), 285 (2), 272 (18), 271 (1), 257 (10), 244 (15), 230 (80), 229 (5), 202 (39), 201 (5) and 187 (17). (200 μ A, 30 eV. electrical detection).

Acknowledgement The authors express their gratitude to Dr. Y. Asahina, Emeritus Professor of University of Tokyo, for his encouragement throughout this study. Thanks are due to prof. T. Fujii of this faculty for discussing the reaction mechanism and Mrs. S. Sasaki (née Yamada) for her technical assistance at early stage of this study and Miss. S. Kitagawa of this faculty for measuring mass spectra and Mr. Y. Itatani of this faculty for measuring NMR spectra and for elemental analyses.