Rihagaku Kenkyusho,
(The Institute of Physical
and Chemical Research)
Wako shi, Saitama

Kaken Chemical Co., Ltd., Tokyo

AKIRA TAHARA
TADASHI NAKATA
YASUO OHTSUKA

SHOICHI TAKADA

Received August 2, 1971

[Chem. Pharm. Bull.]
[19(12)2655-2657(1971)]

UDC 547.918.02

Mimoside: A Glucosidic Metabolite of Mimosine in *Mimosa pudica* and *Leucaena leucocephala*

Mimosine is found in large quantities in the seeds and shoots of the legume species *Mimosa pudica*¹⁾ and *Leucaena leucocephala*²⁾; it shows various physiological activities, i.e. growth retardation, alopecia, cataract and infertility, in experimental animal,³⁻¹⁰⁾ and also inhibits the growth of *Escherichia coli*¹¹⁾ and mung bean seedlings(*Phaseolus aureus*).¹²⁾

Hylin¹³⁾ and Notation, Tiwari, Penrose and Spenser¹⁴⁾ studies the biosynthesis of mimosine in *Mimosa pudica* and *Leucaena leucocephala* by ¹⁴C-labelled precursor techniques: the pyridone ring and the side chain of mimosine could originate from lysine and serine, respectively.

The enzymic synthesis of mimosine from 3-hydroxy-4-pyridone (3,4-dihydroxypyridine) and O-acetylserine using an extract of *Leucaena* seedlings was reported by Murakoshi, Kuramoto, Haginiwa and Fowden¹⁵⁾; neither serine nor O-phosphoserine could serve as a direct substrate in the enzymic reaction.

In continuing our study of mimosine metabolism in plants, we now report that the amino acid is rapidly converted into mimosine-O-glucoside(I), which we have named mimoside.

When *Mimosa pudica* plants were supplied ¹⁴C-mimosine (labelled either universally or specifically in the alanyl side chain) via a cotton wick inserted through the stem, mimoside (I) represented the major metabolic product. I is normally a minor component of *Mimosa pudica* plants in the period before flowering.

Fig. 1 shows the distribution of ¹⁴C in the amino acid fraction of the upper part (stem and attached leaves) of two-month-old *Mimosa pudica* at different times after feeding label-

- 1) J. Renz, *Z. Physiol. Chem.*, **244**, 153 (1936); H.P. Tiwari and I.D. Spenser, *Can. J. Biochem.*, **43**, 1687 (1965).
- 2) M.P. Hegarty and R.D. Court, *Aust. J. Agr. Res.*, **15**, 165 (1964).
- 3) M. Matsumoto, E.G. Smith and G.D. Schermen, *Arch. Biochem. Biophys.*, **33**, 201 (1951).
- 4) von S. Lidwig, G. Petricia and C. Edeanor, *Am. J. Ophth.*, **47**, 107 (1959).
- 5) J.Y. Lin and K.H. Ling, *J. Formosan Med. Assoc.*, **60**, 657 (1961).
- 6) R.G. Crounse, J.D. Maxwell and H. Blank, *Nature*, **194**, 694 (1962).
- 7) K.T. Lin, J.K. Lin and T.C. Tung, *J. Formosan Med. Assoc.*, **63**, 278 (1964).
- 8) M.P. Hegarty, P.G. Schinckel and R.D. Court, *Aust. J. Agr. Res.*, **15**, 153 (1964).
- 9) J.W. Hylin and I.C. Lichton, *Biochem. Pharmacol.*, **14**, 1167 (1965).
- 10) S.S. Yang, C.T. Chin, K.H. Ling and Y.F. Yang, *Transaction of the Ophthalmological Society, Republic of China*, vol 7, p. 12 (1968).
- 11) S. Suda, *Bot. Mag. (Tokyo)*, **73**, 142 (1960).
- 12) I.K. Smith and L. Fowden, *J. Exp. Botany*, **17**, 750 (1966).
- 13) J.W. Hylin, *Phytochem.*, **3**, 161 (1964).
- 14) A.D. Notation and I.D. Spenser, *Can. J. Biochem.*, **42**, 1803 (1964); H.P. Tiwari, W.R. Penrose and I.D. Spenser, *Phytochem.*, **6**, 1245 (1967).
- 15) I. Murakoshi, H. Kuramoto, J. Haginiwa and L. Fowden, *Phytochem.*, in press.

Fig. 1. Flow Diagram to Illustrate the Distribution of ^{14}C in the Amino Acid Fraction of *Mimosa pudica*^{a)} at Various Times after Feeding [$\text{U-}^{14}\text{C}$]-mimosine and ^{14}C -Mimosine labelled only on the Side Chain

a) standard operating condition: 150 cm column, 50° , 0.2N sodium citrate buffer, pH 3.25 followed by 4.25, flow rate 0.5 ml/min. Mimoside (I) was eluted in the area between S-methylcysteine-sulfoxide and methionine-sulfoxide under this condition.

A similar conversion of mimosine into the related glucoside(I) was demonstrated in *Leucaena leucocephala* plants.

The concentration of I in dry *Leucaena* seed is very low, but its concentration increased rapidly during the first 3—5 days growth of seedlings at 30° in the dark. In the later stages of the plant's growth, the concentration of I diminished and it again became a minor component. Therefore I was isolated from 4 or 5-day-old seedlings of *Leucaena leucocephala* by a Dowex 50W ($\times 4$) resin column using 0.2M ammonium formate buffer, pH 2.4, followed on paper, using butan-1-ol-acetic acid-water (3,1,1, by vol.), in 0.013% yield. The isolated compound was identified as a glucoside of mimosine from the profile on an automatic amino acid analyzer.

I shows an absorption peak at 277 nm (ϵ 11400) in water; mimosine absorbed maximally at 284 nm (ϵ 15000) in water. I has mp $178\text{--}179^\circ$, $[\alpha]_D^{25} -60.9$ ($c=0.46$, water); it is very soluble in water, gives a fluorescent spot with the aniline-phthalate reagent for sugars (a general characteristic of glucosides), and a violet spot with ninhydrin on paper. I did not give an immediate colour with 0.1% ferric chloride in 0.1N hydrochloric acid, but a violet colour gradually developed during 25—30 hr at room temperature. This behaviour suggested that slow hydrolysis occurred in the presence of dil. hydrochloric acid to yield mimosine as described below.

I was hydrolyzed by 5% (w/v) hydrochloric acid ($95\text{--}100^\circ$, 1 hr) to give mimosine and glucose in quantitative yield. Mimosine was detected by paper chromatographic procedures, using ninhydrin or ferric chloride as chromogenic reagents, as described in previous report.¹⁵⁾

Further confirmation of the identity of the aglycone as mimosine was obtained using an automatic amino acid analyzer: mimosine eluted from the column at a position close to an isoleucine reference peak under standard operating conditions (Fig. 1).

Glucose present in the hydrolysate was similarly identified by co-chromatography with authentic material; it was detected as a brown spot after spraying with aniline-phthalate and ammoniacal silver nitrate reagents. The following solvent systems were employed: 1, butan-1-ol-acetic acid-water (90,10,29, by vol.); 2, ethyl acetate-water-pyridine (4,4,2, by vol., upper phase); 3, ethyl acetate-water-pyridine (10,3,4, by vol., upper phase); 4, benzene-butan-1-ol-pyridine-water (1,5,3,3, by vol., upper phase). R_f values determined for glucose in these solvents were 0.13, 0.25, 0.18, and 0.17, respectively.

led mimosine: the illustration shows the profile of amino acids eluted from an automatic analyzer (Shibata model AA-500, Tokyo) coupled to a Packard flow monitor system (model 3002) and ratemeter (model 282 A).

Three days after feeding U- ^{14}C -mimosine, 85% of radioactivity still present in the plant (*i.e.* 59% of that administered) was in the amino acid fraction of the upper stem and leaves: at this time, radioactivity associated with organic acids (oxalic, citric, malic, succinic and glutaric acid), sugars (glucose, fructose and two unknowns), and the insoluble residue fraction were 1.1, 5.9, and 6.2%, respectively. The roots contained only 1.6% of the total activity.

For further confirmation of the glycone moiety as glucose, a specific β -D-glucose oxidase (Worthington) was employed. When the enzymic system was coupled to O-dianisidine as a chromogenic reagent (colour maximum at 530 nm¹⁶⁾, colour production was theoretical for glucose. Quantitative analysis showed that glucose was conjugated with mimosine in equimolar amounts.

Conjugation was assumed to involve the C-1 of glucose, because I (mimoside) lacked reducing properties, and the C-3 hydroxy group of mimosine, since mimoside (I) gave no immediate colour with 0.1% ferric chloride.

The glucoside (I) also could be degraded to mimosine and glucose (30% yield in 1 hr at 30°) by the β -D-glucosidase of sweet almonds (Miles).

Therefore the structure of mimoside (I) was considered to be 2-N-(3- β -D-glucosyloxy-4-pyridone)-1-aminopropionic acid (formula I).

Mimoside (I) was synthesized by an extract of *Leucaena* seedlings from mimosine and uridine diphosphate-glucose, and also conversely was degraded into mimosine and D-glucose by the hydrolytic enzyme in *Leucaena* seedlings. Those results will be reported in the following paper.

Acknowledgement The kind encouragement of Prof. L. Fowden, Department of Botany and Microbiology, University College London, during the course of this work is acknowledged. We are indebted also to Drs. Y. Kashida and K. Arai, National Institute of Radiological Sciences, Chiba, for preparing U-¹⁴C-mimosine. Finally, thanks are due to Prof. S. Okuda and Mrs. H. Hino, Institute of Applied Microbiology, University of Tokyo, for microanalyses.

Faculty of Pharmaceutical Sciences,
University of Chiba
1-33, Yayoi-cho, Chiba

ISAMU MURAKOSHI
SHIGERU OHMIYA
JOJU HAGINIWA

Received August 17, 1971

16) M. Kitamura, H. Migami, and Y. Arimatsu, *J. Med. Technology* (Tokyo), 6, 18 (1962).