[Chem. Pharm. Bull.]
21(7) 1612 (1973)

UDC 547.831.3.04.09 : 547.466.1.057

Studies on Peptides. XXXVI.¹⁾ Chloro-substituted 8-Hydroxyquinoline Derivatives, as Racemization Suppressing Reagents in Peptide Synthesis

HARUAKI YAJIMA, MASAYUKI KUROBE, and KANAME KOYAMA

Faculty of Pharmaceutical Sciences, Kyoto University²⁾

(Received January 20, 1973)

As an active ester in peptide synthesis, the 8-hydroxyquinolyl ester of N-protected amino acids was first introduced by Jakubke³⁾ in 1965. This bifunctional active ester, as examined in detail by the same authors,⁴⁾ suppresses the rate of racemization during the aminolysis.

Applying the system of Bodanszky and Conklin,⁵⁾ known as one of convenient and sensitive tests for racemization, we have measured the degree of racemization which occurred during the coupling reaction of Ac-L-Ile-OH with H-Gly-OEt by dicyclohexylcarbodiimide (DCC) plus 8-hydroxyquinoline or its commercially available chloro-derivatives: 5-chloro or 5,7-dichloro-8-hydroxyquinoline. As shown in Table I, racemization decreases as the number of chlorosubstituents of these reagents increases. It appears, therefore, that the pK_a values of these reagents⁶⁾ are closely associated with this tendency.

Under identical conditions, the reaction of DCC alone gave a racemate in 27.4%,⁷⁾ while DCC plus N-hydroxybenztriazole⁸⁾ gave only 2.4%. These results suggest that 5,7-dichloro-8-hydroxyquinoline is an effective racemization-depressant nearly comparable to N-hydroxybenztriazole, known as the most effective reagent for the racemization-free peptide synthesis.

TABLE I. Degree of Racemization

Reagent	8-Hydroxy-quinoline	5-Chloro-8-hydroxy-quinoline	5,7-Dichloro-8-hydroxy-quinoline
Racemization %	21.3	12.4	5.3

Experimental

Ac-Ile-Gly-OEt—To a solution of Ac-L-Ile-OH (0.87 g, 0.5 mmole) in dimethylformamide (20 ml), H-Gly-OEt (prepared from 0.7 g of the hydrochloride with 0.7 ml of triethylamine) in dimethylformamide (10 ml) was combined. To this solution, DCC (0.5 mmole) and 8-hydroxyquinoline or 5-chloro-8-hydroxyquinoline or 5,7-dichloro-8-hydroxyquinoline (0.5 mmole each) was added. The solution, after stirring at 20° for 18 hr, was filtered and then evaporated *in vacuo* under 45°. The residue was dissolved in AcOEt, which after washing with 10% citric acid, 5% Na₂CO₃ and H₂O-NaCl, was dried over Na₂SO₄ and then evaporated. The solid residue was dried over P₂O₅ *in vacuo*. The racemate, allo-D-Ile-OH, was determined quantitatively, after acid hydrolysis with 6N HCl for 18 hr, by the amino acid analyzer (Spackman-Stein-Moore method).⁹⁾ Acid hydrolysate of the starting Ac-L-Ile-OH gave no measurable racemate and the results were listed in Table I.

1) Part XXXV: H. Yajima, F. Tamura, Y. Kiso, and M. Kurobe, *Chem. Pharm. Bull.* (Tokyo), **21**, 1380 (1973).

2) Location: Sakyo-ku, Kyoto.

3) H.D. Jakubke, *Z. Naturforsch.*, **20b**, 273 (1965).

4) H.D. Jakubke and A. Voigt, *Chem. Ber.*, **99**, 2419, 2944 (1966); H.D. Jakubke, A. Voigt, and S. Burkhardt, *ibid.*, **100**, 2367 (1967).

5) M. Bodanszky and L.E. Conklin, *Chem. Commun.*, **1967**, 773.

6) M.R. Chakrabarty, E.S. Hanrahan, N.D. Heindel, and G.F. Watts, *Anal. Chem.*, **39**, 238 (1967).

7) Y. Kiso and H. Yajima, *Chem. Commun.*, **1972**, 492.

8) W. König and R. Geiger, *Chem. Ber.*, **103**, 788 (1970).

9) D.H. Spackman, W.H. Stein, and S. Moore, *Anal. Chem.*, **30**, 1190 (1958).