Studies on Ketene and Its Derivatives. LXXIII.¹⁾ Synthesis of Pyrido[1,2-*a*]-pyrimidin-4-one and Pyrido[1,2-*a*]-*sym*-triazin-4-one Derivatives

TETSUZO KATO and SHINICHI MASUDA

Pharmaceutical Institute Tohoku University²⁾

(Received January 25, 1975)

Ethyl N-(2-pyridyl)formimidate (Va) reacted with diketene to give 3-acetyl-4*H*-pyrido[1,2-*a*]pyrimidin-4-one (VIIa) and ethyl 2-(2-pyridylaminomethylene)acetoacetate (VIIIa).

Reaction of diketene with ethyl N-(2-pyridyl)acetimidate (Vb) and ethyl N-(2-pyridyl)propionimidate (Vc) afforded 4*H*-pyrido[1,2-*a*]pyrimidine derivatives (VIIb and VIIc), respectively. Similarly 2-benzylideneaminopyridine (VIa) reacted with diketene to give 3-acetyl-4-hydroxy-2-phenyl-4*H*-pyrido-[1,2-*a*]pyrimidine (XI).

On the other hand, reaction of trimethylsilyl isocyanate with Va, Vb, and Vc gave rise to 4*H*-pyrido[1,2-*a*]-*sym*-triazin-4-one derivatives (XIXa, XIXb, and XIXc), respectively. Reaction of phenyl isocyanate with VIa afforded 2,3-diphenyl-4*H*-pyrido[1,2-*a*]-*sym*-2,3-dihydrotriazin-4-one (XX).

In the previous paper of this series, we have reported that diketene reacted with imidates such as ethyl benzimidate (I) and Schiff bases such as N-benzylideneaniline (II) to give 2-ethoxy-6-methyl-2-phenyl-3,4-dihydro-2*H*-1,3-oxazin-4-one (III)³⁾ and 2-benzylideneacetoacetanilide (IV),⁴⁾ respectively. The reactions involve the addition of diketene to the C=N double bond of I and II forming a six and four membered cyclic intermediate followed by prototropy to give III and IV, respectively.

Chart 1

As a continuation of the study we investigated the reaction of diketene with N-(2-pyridyl)-imidate (V) and 2-benzylideneaminopyridine (VI) in an effort to prepare the pyridyl-1,3-oxazine and benzylideneacetoacetamidopyridine derivatives, and found that the reactions did not lead to the expected oxazine and acetoacetamido derivatives, but lead pyrido[1,2-*a*]-pyrimidone derivatives presumably by the 1,4-dipolar addition reaction.

1) Part LXXII: T. Kato, Y. Yamamoto, and M. Kondo, *Heterocycles*, **4**, 293 (1975).

2) Location: Aobayama, Sendai, 980, Japan.

3) T. Kato and Y. Yamamoto, *Chem. Pharm. Bull.* (Tokyo), **15**, 1334 (1967).

4) T. Kato and Y. Yamamoto, *Chem. Pharm. Bull.* (Tokyo), **13**, 959 (1965).

Although the mechanism of the formation of these products is not clear at present, likely pathways are shown in Chart 3. Namely, the electrophilic addition of the carbonyl carbon of diketene to the ring nitrogen of V would form the dipolar intermediate XIII which cyclizes to the dihydropyrido[1,2-*a*]pyrimidone intermediate (XIV) followed by the elimination of EtOH to give VII. (path-a). If the electrophilic addition of diketene occurs at the imide nitrogen, the four membered cyclic intermediate (XVI) will form *via* the dipolar ionic intermediate XV. Ring fission of XVI accompanied with prototropy gives rise to VIIIa (path-b).⁷⁾

Another probable mechanism for the formation of VIIIa will be proposed as follows: ethyl acetoacetate, produced from diketene and EtOH which was isolated during the formation of VIIa, reacts with Va to give VIIIa.

According to the similar procedure given for VII from V, the electrophilic attack of diketene to the ring nitrogen of VIa gives rise to the adduct XVII, the dipolar cycloaddition of which affords the pyrido-pyrimidone intermediate (XVIII), which by prototropy is transformed to XI.

Chart 3

2. Reaction of Isocyanates with N-(2-Pyridyl)imidates (V) and 2-Benzylideneaminopyridine (VIa)

When ethyl N-(2-pyridyl)formimidate (Va, R=H) was allowed to react with trimethylsilyl isocyanate at room temperature, colorless needles of mp 210°, C₇H₅ON₃ (XIXa, R=H), were obtained in good yield. Elemental analysis and spectral data uniquely suggested the structure

7) T. Kato, *Accounts of Chem. Research*, **7**, 265 (1974).

being 4*H*-pyrido[1,2-*a*]-*sym*-triazin-4-one. Similarly, ethyl *N*-(2-pyridyl)acetimidate (Vb, R=CH₃) and ethyl *N*-(2-pyridyl)propionimidate (Vc, R=C₂H₅) reacted with trimethylsilyl isocyanate to give good yields of 2-methyl-4*H*-pyrido[1,2-*a*]-*sym*-triazin-4-one (XIXb, R=CH₃) and 2-ethyl-4*H*-pyrido[1,2-*a*]-*sym*-triazin-4-one (XIXc, R=C₂H₅), respectively.

Though the similar reaction of 2-benzylideneaminopyridine (VIa, R=H) with trimethylsilyl isocyanate did not afford any product corresponding to the pyridotriazine derivative such as XIX, the reaction of VIa with phenyl isocyanate gave rise to an adduct, C₁₉H₁₅ON₃ (XX), in 83% yield, which was identified as 2,3-diphenyl-4*H*-pyrido[1,2-*a*]-*sym*-2,3-dihydrotriazine-4-one on the basis of elemental analysis and spectral data.

Similar reaction of Va—c with phenyl isocyanate gave crystalline substances in pretty good yield, however, they were very unstable and purification by recrystallization or chromatography failed.

Chart 4

Experimental

IR spectra were measured by a Nippon-Bunko Model IR-S spectrophotometer. NMR spectra were determined on a Hitachi-Perkin Elmer R=20 spectrophotometer at 60 MHz. Values are given in ppm relative to tetramethylsilane as an internal standard. Abbreviations are as follows: s=singlet, d=doublet, t=triplet, q=quartet, m=multiplet, br=broad. All melting points were uncorrected.

Reaction of Diketene with Ethyl *N*-(2-Pyridyl)formimidate (Va, R=H)—A mixture of Va (1.5 g, 0.01 mole) and diketene (3.4 g, 0.04 mole) was allowed to stand at room temperature for 3 days. Crystals separated were collected by suction, and recrystallized from EtOH to orange needles of mp 151–152° (VIIa) (lit.⁵ mp 152°). Yield, 0.5 g.

The filtrate was condensed to dryness, and the resulting residue was purified by silica gel column chromatography using a mixture of petroleum ether and ether (1:1) and then acetone as eluants. From the petroleum ether–ether eluant a crystalline substance was obtained. Recrystallization from petroleum ether–ether gave colorless prisms of mp 89–90° (VIIIa). Yield, 1.2 g (51%). *Anal.* Calcd. for C₁₂H₁₄O₃N₂ (VIIIa): C, 61.52; H, 6.02; N, 11.96. Found: C, 61.62; H, 6.03; N, 12.15. IR $\nu_{\text{max}}^{\text{CHCl}_3}$ cm⁻¹: 1690, 1630. NMR (CDCl₃) ppm: 1.35 (3H, t, *J*=6.75 Hz), 2.58 (3H, s), 4.28 (2H, q, *J*=6.75 Hz), 6.80–8.50 (4H, m), 9.18 (1H, d, *J*=11.4 Hz), 12.70 (1H, d, *J*=11.4 Hz).

The acetone eluted fraction afforded 0.2 g of VIIa. Total yield of VIIa was 0.7 g (38%).

Ethyl 2-(2-Pyridylaminomethylene)acetacetate (VIIIa)—A mixture of Va (1.5 g) and ethyl acetacetate (2 g) was allowed to stand at room temperature for 3 days. The reaction mixture was condensed *in vacuo*, and the crystalline residue was purified by recrystallization from ether–petroleum ether to give 1.9 g (88%) of VIIIa.

Cyclization of VIIIa to VIIa—A solution of VIIIa (0.5 g) in Ac₂O (5 ml) was refluxed for 20 min. After removal of Ac₂O by vacuum distillation, the reaction mixture was washed with ether to give a crystalline

residue. Purification by recrystallization from EtOH gave 0.35 g (88%) of VIIa.

Reaction of Diketene with Ethyl N-(2-Pyridyl)acetimidate (Vb, R=CH₃)—A mixture of Vb (1.6 g) and diketene (3.4 g) was allowed to stand at room temperature for 3 days. Crystals separated were collected by suction. Recrystallization from MeOH afforded yellow needles of mp 159° (VIIb). Yield, 0.5 g. *Anal.* Calcd. for C₁₁H₁₀O₂N₂ (VIIb): C, 65.33; H, 4.98; N, 13.86. Found: C, 65.45; H, 4.65; N, 13.83. IR $\nu_{\text{max}}^{\text{CHCl}_3}$ cm⁻¹: 1700 (sh), 1670, 1630. NMR (CDCl₃) ppm: 2.59 (3H, s), 2.68 (3H, s), 7.08—8.07 (3H, m), 9.10 (1H, m).

The filtrate was chromatographed on a silica gel column using petroleum ether, ether and acetone as eluants. The petroleum ether-ether (3:1) eluted fraction gave 0.6 g of ethyl acetoacetate. The ether eluted fraction gave 2-acetamidopyridine (IXb), mp 69° (lit.⁸) mp 67°. The acetone elution gave 0.7 g of VIIb. Total yield of VIIb was 1.2 g (65%).

2-Methyl-4H-pyrido[1,2-a]pyrimidin-4-one—A solution of VIIb (0.2 g) in 70% H₂SO₄ (10 ml) was heated at 140° for 3 hr. The reaction mixture was poured into ice-water, neutralized with NaHCO₃, and extracted with CHCl₃. The CHCl₃ solution was condensed, and the resulting residue was recrystallized from ether to pale yellow needles of mp 120°, undepressed on admixture with an authentic sample prepared according to the literature.⁶⁾

Reaction of Diketene with Ethyl N-(2-Pyridyl)propionimidate (Vc, R=C₂H₅)—A mixture of Vc (3.6 g) and diketene was allowed to stand at room temperature for 4 days. After removal of excess diketene by vacuum distillation, the resulting residue was extracted with ether. The ether soluble fraction was purified by silica gel column chromatography using petroleum ether and ether as eluants. From the petroleum ether-ether (3:1) fraction, 0.9 g of ethyl acetoacetate was isolated. The 1:1 mixture of petroleum ether and ether gave 0.4 g (14%) of 2-propionamidopyridine (IXc), mp 62°. The ether elution gave pale yellow prisms of mp 90° (VIIc). Yield, 1.9 g (72%). *Anal.* Calcd. for C₁₂H₁₂O₂N₂ (VIIc): C, 66.65; H, 5.59; N, 12.96. Found: C, 66.78; H, 5.72; N, 12.93. IR $\nu_{\text{max}}^{\text{CHCl}_3}$ cm⁻¹: 1700 (sh), 1668, 1630. NMR (CDCl₃) ppm: 1.31 (3H, t, J=7.2 Hz), 2.70 (3H, s), 2.88 (2H, q, J=7.2 Hz), 7.10—8.10 (3H, m), 9.15 (1H, m).

Reaction of Diketene with 2-Benzylideneaminopyridine (VIa, R=H)—Diketene (1.3 g) was added dropwise to VIa (1.8 g) under ice-cooling with stirring for 5 hr. The reaction mixture, which solidified, was filtered. The crystalline substance was washed with a small amount of cold MeOH. Recrystallization from absolute MeOH gave yellow needles of mp 211° (decomp.) (XI). Yield, 2.0 g (76%). *Anal.* Calcd. for C₁₆H₁₄O₂N₂ (XI): C, 72.16; H, 5.30; N, 10.52. Found: C, 71.96; H, 5.47; N, 10.36. IR $\nu_{\text{max}}^{\text{CHCl}_3}$ cm⁻¹: 2700—3300, 1685, 1645. NMR (DMSO-d₆) ppm: 3.38 (3H, s), 5.72 (1H, d, J=3 Hz), 6.70—8.00 (8H, m), 8.70 (1H, m), 9.92 (1H, d, J=3 Hz).

3-Acetyl-2-phenyl-4H-pyrido[1,2-a]pyrimidin-4-one (XII)—A suspension of XI (0.27 g, 0.001 mole) and chloranil (0.25 g, 0.001 mole) in benzene (20 ml) was refluxed for 1 hr. The reaction mixture was extracted with 10% HCl. The HCl solution was neutralized with NaHCO₃, and extracted with CHCl₃. The CHCl₃ solution was condensed, and the resulting residue was purified by recrystallization from MeOH to pale brown needles of mp 158° (XII). Yield, 0.2 g (74%). *Anal.* Calcd. for C₁₆H₁₂O₂N₂ (XII): C, 72.71; H, 4.58; N, 10.60. Found: C, 72.84; H, 4.80; N, 10.75. IR $\nu_{\text{max}}^{\text{CHCl}_3}$ cm⁻¹: 1700, 1668, 1635. NMR (CDCl₃) ppm: 2.47 (3H, s), 7.00—8.00 (8H, m), 9.10 (1H, m).

4H-Pyrido[1,2-a]-sym-triazin-4-one (XIXa, R=H)—A mixture of Va (R=H) (3.0 g, 0.02 mole) and trimethylsilyl isocyanate was stirred at room temperature for 2 days. Crystals separated were collected by suction. Recrystallization from acetone gave colorless needles of mp 210° (XIXa, R=H). Yield, 2.4 g (81%). *Anal.* Calcd. for C₇H₅ON₃ (XIXa): C, 57.14; H, 3.43; N, 28.56. Found: C, 56.99; H, 3.56; N, 28.97. IR $\nu_{\text{max}}^{\text{CHCl}_3}$ cm⁻¹: 1715, 1638. NMR (CF₃CO₂H) ppm: 7.90—9.00 (3H, m), 8.68 (1H, s), 9.40 (1H, m).

The filtrate was distilled to give a colorless oil of bp 62°, (CH₃)₃SiOEt. Yield, 0.4 g (17%).

2-Methyl-4H-pyrido[1,2-a]-sym-triazin-4-one (XIXb, R=CH₃)—Following the similar procedure as above, ethyl N-(2-pyridyl)acetimidate (Vb, R=CH₃) (3.3 g) was allowed to react with trimethylsilyl isocyanate (2.3 g) to give colorless needles (AcOEt) of mp 165° (XIXb, R=CH₃). Yield, 2.8 g (88%). *Anal.* Calcd. for C₈H₇ON₃ (XIXb): C, 59.62; H, 4.38; N, 26.07. Found: C, 59.54; H, 4.53; N, 25.71. IR $\nu_{\text{max}}^{\text{CHCl}_3}$ cm⁻¹: 1710, 1635. NMR (CDCl₃) ppm: 2.58 (3H, s), 7.26—8.30 (3H, m), 9.03 (1H, m).

2-Ethyl-4H-pyrido[1,2-a]-sym-triazin-4-one (XIXc, R=C₂H₅)—Following similar procedure described above, ethyl N-(2-pyridyl)propionimidate (Vc, R=C₂H₅) (3.6 g) was allowed to react with trimethylsilyl isocyanate (2.3 g) to give colorless leaves (AcOEt) of mp 148° (XIXc, R=C₂H₅). Yield, 2.6 g (74%). *Anal.* Calcd. for C₉H₉ON₃ (XIXc): C, 61.70; H, 5.18; N, 23.99. Found: C, 62.06; H, 5.35; N, 24.15. IR $\nu_{\text{max}}^{\text{CHCl}_3}$ cm⁻¹: 1710, 1638. NMR (CDCl₃) ppm: 1.36 (3H, t, J=7.8 Hz), 2.80 (2H, q, J=7.8 Hz), 7.22—8.28 (3H, m), 9.04 (1H, m).

2,3-Diphenyl-4H-pyrido[1,2-a]-sym-2,3-dihydrotriazin-4-one (XX)—A mixture of 2-benzylideneaminopyridine (VIa, R=H) (1.8 g) and phenyl isocyanate (1.2 g) was stirred at room temperature for 1 hr. Crystals separated were collected by suction. Purification by recrystallization from cold absolute MeOH gave yellow needles of mp 90°. Yield, 2.5 g (83%). *Anal.* Calcd. for C₁₉H₁₅ON₃ (XX): C, 75.73; H, 5.02;

8) A.L. Mndzhoyan and V.G. Afrikyan, *Ser. Khim. Nauk.*, **10**, 143 (1957).

N, 13.95. Found: C, 76.09; H, 5.23; N, 13.89. IR $\nu_{\text{max}}^{\text{CHCl}_3}$ cm^{-1} : 1700, 1650. NMR (CDCl_3) ppm: 5.70—7.50 (14H, m), 7.80 (1H, m).

Acknowledgement The authors are indebted to Mrs. A. Sato, Mrs. C. Koyanagi, Miss A. Ujiie of the Central Analysis Room of this Institute for elemental analysis and NMR spectral measurement. A part of the expences for this work was defrayed by the Grand-in-Aid for Scientific Research from the Ministry of Education, which is gratefully acknowledged.