[Chem. Pharm. Bull.
23(8)1869-1873(1975)]

UDC 547.831'233.057 : 615.28.015.11

Studies on Antibacterial Agents. III. Synthesis of N,N'-Bis(4-quinolyl-, 4-quinaldinyl-, 4-quinazolinyl-, or 9-acridinyl)polymethylenediamines and Their Sulfonamides

ASHOK KUMAR SINGH

Department of Chemistry, Faculty of Science, U.P. Postgraduate College¹⁾

(Received July 10, 1974)

Polymethylene bis-amines, with heterocycle as a carrier moiety, and their sulfonamides were synthesized and tested against *S. aureus*, *E. coli* and *M. tuberculosis* (H37Rv strain). Structure-activity relationship has been discussed.

Sulfonamides²⁻¹⁰⁾ are known to exhibit antibacterial properties. Though a large number of sulfonamides have been synthesized, compounds derived from bis-amines have not been reported so far. Our interest in synthetic antibacterials^{11,12)} prompted us to synthesize bis-amines (I) and their sulfonamides (II), and to evaluate their scope as antibacterial agents, particularly with reference to the heterocycle, *i.e.*, the carrier moiety and the size of the side chain.

Sulfonylation of the bis-amines with aromatic sulfonyl chloride in dry pyridine gave the corresponding sulfonamides which were characterized by elemental and spectral analyses.

Biological assay of the bis-amines and their sulfonamides were carried out *in vitro* against *S. aureus* and *E. coli* following Humphrey and Lightbrown's method,¹³⁾ and against *M. tuberculosis* (H37Rv strain) following the procedure of Youman.¹⁴⁾ The results are tabulated in Tables II and IV respectively.

- 1) Location: Varanasi-2, U.P., India.
- 2) S.F. Zakrzewski, A. Bloch, and C.A. Nichol, *J. Med. Chem.*, **11**, 885 (1968).
- 3) S. Wada, K. Kumaki, and I. Moriguchi, *Chem. Pharm. Bull.* (Tokyo), **17**, 2168 (1969).
- 4) L. Doub, U. Krolls, J.M. Vandeneelt, and M.W. Fisher, *J. Med. Chem.*, **13**, 242 (1970).
- 5) W.E. Kreighbaum, F.A. Grunwald, E.F. Harrison, J.A. Labudde, and A.A. Larsen, *J. Med. Chem.*, **13**, 247 (1970).
- 6) Z. Budesinsky, V. Valenta, V. Vosatko, and A. Simek, *Collect. Czech. Chem. Commun.*, **37**, 887 (1972).
- 7) N. Saito, T. Kurihara, S. Yasuda, K. Yamanaka, S. Tsuruta, T. Tanaka, and Y. Inamori, *Yakugaku Zasshi*, **88**, 1610 (1968).
- 8) N.D. Heindel, C.C.H. Ko, R.B. Birrer, and J.R. Merkel, *J. Med. Chem.*, **15**, 118 (1972).
- 9) G.H. Miller, P.H. Doukas, and J.K. Seydel, *J. Med. Chem.*, **15**, 700 (1972).
- 10) A. Lewis and R.G. Shepherd, "Medicinal Chemistry," Edt. A. Burger, Part I, Wiley-Interscience, New York, 1970, p. 409.
- 11) S.M. Deshpande and A.K. Singh, *Agv. Biol. Chem.* (Tokyo), **35**, 119 (1971).
- 12) S.M. Deshpande and A.K. Singh, *Chem. Pharm. Bull.* (Tokyo), **20**, 206 (1972).
- 13) J.H. Humphrey and J. Lightbrown, *J. Gen. Microbiol.*, **7**, 129 (1952).
- 14) G.P. Youman's, *Proc. Soc. Exp. Biol.*, **57**, 122 (1944).

Com- pound <i>n</i> No.	R	Formula ^{a)}	Yield (%)	mp ^{b)} (°C)	Calcd.			Found			
					C	H	N	C	H	N	
1	6	4-quinolyl	C ₂₄ H ₂₆ N ₄	78	195—196	77.82	7.02	15.13	78.00	7.00	14.98
2	7	4-quinolyl	C ₂₅ H ₂₈ N ₄	71	102—103	78.12	7.29	14.58	78.20	7.16	14.38
3	8	4-quinolyl	C ₂₆ H ₃₀ N ₄	77	164—165	78.39	7.53	14.07	78.18	7.48	13.69
4	9	4-quinolyl	C ₂₇ H ₃₂ N ₄	80	91—92	78.64	7.76	13.59	78.52	7.68	13.28
5	10	4-quinolyl	C ₂₈ H ₃₄ N ₄	82	156—157	78.87	7.98	13.14	78.66	7.89	13.00
6	12	4-quinolyl	C ₃₀ H ₃₈ N ₄	76	122—123	79.29	8.37	12.33	79.08	8.40	12.26
7	3	4-quinaldinyl	C ₂₃ H ₂₄ N ₄	80	254—255	77.52	6.74	15.81	77.30	6.66	15.66
8	6	4-quinaldinyl	C ₂₆ H ₃₀ N ₄	77	191—192	78.39	7.53	14.07	78.28	7.50	14.10
9	7	4-quinaldinyl	C ₂₇ H ₃₂ N ₄	71	137—138	78.64	7.78	13.59	78.48	7.72	13.38
10	8	4-quinaldinyl	C ₂₈ H ₃₄ N ₄	76	210—211	78.87	7.98	13.14	78.68	7.69	12.98
11	9	4-quinaldinyl	C ₂₉ H ₃₆ N ₄	70	172—173	79.09	8.18	12.72	79.00	8.06	12.48
12	10	4-quinaldinyl	C ₃₀ H ₃₈ N ₄	70	150—151	79.29	8.37	12.33	79.18	8.26	12.28
13	12	4-quinaldinyl	C ₃₂ H ₄₂ N ₄	75	115—116	79.66	8.71	11.61	79.44	8.70	11.46
14	9	4-quinazolinyl	C ₂₅ H ₃₀ N ₆	75	174—175	72.46	7.24	20.28	72.38	7.18	20.10
15	10	4-quinazolinyl	C ₂₆ H ₃₂ N ₆	82	69—70	76.47	7.84	20.58	76.38	7.68	20.42
16	8	9-acridinyl	C ₃₄ H ₃₄ N ₄	69	165—166	81.92	6.82	11.24	81.69	6.70	11.16

a) IR (Nujol) for all the compounds recorded absorption at 3300 cm⁻¹ (NH group).

b) crystallizationsolvent: chloroform

TABLE II^{a)}

No. from Table I	Antibacterial activity 0.1 ml of a 10 ⁻² M solution of the compound in water or alcohol by cup-plate method Diameter of cup is 10 mm		Antitubercular activity tested against <i>M. tuberculosis</i> (H37Rv strain) Growth in Youman's medium at the end of three weeks Concentration of drugs			
	<i>S. aureus</i>	<i>E. coli</i>	10 ⁻⁴ M	10 ⁻⁵ M	10 ⁻⁶ M	Rv control
1	26	31	—Ve	3+	4+	4+
2	28	28	—Ve	2+	4+	4+
3	24	27	—Ve	+	4+	4+
4	24	27	—Ve	Sl	4+	4+
5	22	26	—Ve	+	4+	4+
6	19	24	—Ve	+	4+	4+
7	28	27	Sl	3+	4+	4+
8	34	27	Sl	3+	4+	4+
9	27	27	—Ve	2+	4+	4+
10	27	28	—Ve	2+	4+	4+
11	27	34	—Ve	+	3+	4+
12	25	30	—Ve	—Ve	4+	4+
13	22	26	—Ve	—Ve	4+	4+
14	15	22	+	3+	4+	4+
15	14	20	+	2+	4+	4+
16	20	22	0	2+	4+	4+
5% Phenol control value	23	34	0	=no growth		
			+to 4+	=various stages of growth		
Ethanol	—Ve	—Ve	Sl	=slight growth		

a) All the compounds were tested *in vitro*.

—Ve: negative result

Sl : slight growth

TABLE III. $R'-SO_2-N-(CH_2)_n-N-SO_2-R'$
 $\begin{matrix} R' & & R \end{matrix}$

Com- pound No.	R	R'	Formula	Yield (%)	mp ^{a)} (°C)	Calcd.			Found			IR data ν cm ⁻¹ (m ^{b)})
						C	H	N	C	H	N	
1	6	4-quinolyl	$C_{38}H_{32}O_4N_4S_2Cl_2$	88	180—181	60.08	4.45	7.78	60.10	4.42	7.60	760(C-Cl), 1600(aromatic)
2	7	4-quinolyl	$C_{37}H_{34}O_4N_4S_2Cl_2$	76	194—195	60.57	4.63	7.63	60.46	4.48	7.48	750(C-Cl), 1600(aromatic)
3	8	4-quinolyl	$C_{38}H_{36}O_4N_4S_2Cl_2$	80	215—216	61.04	4.81	7.49	64.20	4.66	7.36	760(C-Cl), 1600(aromatic)
4	10	4-quinolyl	$C_{40}H_{40}O_4N_4S_2Cl_2$	82	209—210	61.93	5.16	7.22	61.68	5.08	7.08	760(C-Cl), 1600(aromatic)
5	12	4-quinolyl	$C_{42}H_{44}O_4N_4S_2Cl_2$	78	234—235	62.76	5.47	6.97	62.66	5.48	6.66	760(C-Cl), 1610(aromatic)
6	8	4-quinolyl	$C_{42}H_{44}O_6N_6S_2$	74	113—114	63.63	5.55	10.60	63.48	5.60	10.48	1600(aromatic), 1685(CO), 3320(NH)
7	10	4-quinolyl	$C_{44}H_{48}O_6N_6S_2$	76	91—92	64.39	5.85	10.24	64.18	5.66	10.33	1610(aromatic), 1680(CO), 3310(NH)
8	12	4-quinolyl	$C_{46}H_{52}O_6N_6S_2$	83	88—89	65.09	6.13	9.90	64.98	6.08	9.68	1610(aromatic), 1680(CO), 3320(NH)
9	6	4-quinolyl	$C_{38}H_{38}O_4N_4S_2$	80	134—135	67.25	5.60	8.25	66.98	5.45	8.08	1600(aromatic)
10	7	4-quinolyl	$C_{39}H_{40}O_4N_4S_2$	92	171—172	67.63	5.78	8.09	67.48	5.62	7.88	1600(aromatic)
11	8	4-quinolyl	$C_{40}H_{42}O_4N_4S_2$	91	185—186	67.98	5.94	7.93	67.67	5.78	7.80	1600(aromatic)
12	9	4-quinolyl	$C_{41}H_{44}O_4N_4S_2$	89	82—83	68.33	6.11	7.77	68.18	6.00	7.40	1600(aromatic)
13	10	4-quinolyl	$C_{42}H_{46}O_4N_4S_2$	85	170—171	68.66	6.26	7.62	68.40	6.30	7.28	1610(aromatic)
14	12	4-quinolyl	$C_{44}H_{50}O_4N_4S_2$	74	210—211	69.29	6.56	7.34	68.98	6.46	7.38	1610(aromatic)
15	6	4-quinolyl	$C_{38}H_{36}O_4N_4S_2Cl_2$	70	111—112	61.04	4.81	7.49	61.18	4.66	7.40	760(C-Cl), 1610(aromatic)
16	8	4-quinolyl	$C_{40}H_{40}O_4N_4S_2Cl_2$	82	115—116	61.93	5.16	7.22	61.68	5.08	7.30	750(C-Cl), 1610(aromatic)
17	9	4-quinolyl	$C_{41}H_{42}O_4N_4S_2Cl_2$	84	157—158	62.35	5.32	7.09	62.08	5.38	6.82	750(C-Cl), 1610(aromatic)
18	10	4-quinolyl	$C_{42}H_{44}O_4N_4S_2Cl_2$	75	204—205	62.76	5.47	6.97	62.60	5.45	6.66	760(C-Cl), 1610(aromatic)
19	6	4-quinolyl	$C_{42}H_{44}O_6N_6S_2$	80	310 (decomp.)	63.63	5.55	10.60	63.48	5.62	10.48	1615(aromatic), 1680(CO), 3310(NH)
20	8	4-quinolyl	$C_{44}H_{48}O_6N_6S_2$	72	229—230	64.39	5.85	10.24	64.40	5.66	10.08	1600(aromatic), 1680(CO), 3310(NH)
21	10	4-quinolyl	$C_{46}H_{52}O_6N_6S_2$	81	129—130	65.09	6.13	9.90	65.20	6.08	9.88	1600a(aromatic), 1680(CO), 3310(NH)
22	12	4-quinolyl	$C_{48}H_{56}O_6N_6S_2$	90	110—111	65.75	6.39	9.58	65.58	6.28	9.28	1610(aromatic), 1690(CO), 3340(NH)
23	3	4-quinolyl	$C_{37}H_{36}O_4N_4S_2$	74	122—123	66.86	5.42	8.43	66.66	5.33	8.18	1615(aromatic)
24	6	4-quinolyl	$C_{40}H_{42}O_4N_4S_2$	79	221—222	67.98	5.94	7.92	67.88	5.88	8.29	1600(aromatic)
25	7	4-quinolyl	$C_{41}H_{44}O_4N_4S_2$	75	139—140	68.33	6.11	7.77	68.30	6.08	7.47	1600(aromatic)
26	8	4-quinolyl	$C_{42}H_{46}O_4N_4S_2$	77	125—126	68.66	6.26	7.62	68.48	6.18	7.20	1600(aromatic)
27	9	4-quinolyl	$C_{43}H_{48}O_4N_4S_2$	84	101—102	68.98	6.41	7.48	68.88	6.28	7.54	1610(aromatic)
28	10	4-quinolyl	$C_{44}H_{50}O_4N_4S_2$	84	175—176	69.29	6.56	7.36	69.08	6.48	7.56	1610(aromatic)
29	12	4-quinolyl	$C_{46}H_{54}O_4N_4S_2$	91	208—209	69.87	6.83	7.08	69.66	6.66	6.75	1600(aromatic)
30	9	4-quinolyl	$C_{39}H_{40}O_4N_4S_2$	80	202—203	64.81	5.81	11.63	64.66	5.68	11.48	1610(aromatic)
31	10	4-quinolyl	$C_{40}H_{44}O_4N_4S_2$	76	229—230	65.21	5.97	11.41	65.18	5.78	11.18	1600(aromatic)
32	8	9-acridinyl	$C_{48}H_{46}O_4N_4S_2$	80	220—221	71.46	5.70	6.94	71.28	7.68	6.80	1615(aromatic)

a) crystallizationsolvent: hot water; b) m = medium intensity

TABLE IV^{a)}

No. from Table III	Antibacterial activity 0.1 ml of a 10 ⁻² M solution of the compound in water or alcohol by cup-plate method Diameter of cup is 10 mm		Antitubercular activity tested against <i>M. tuberculosis</i> (H37Rv strain) Growth in Youman's medium at the end of three weeks Concentration of drugs			
	<i>S. aureus</i>	<i>E. coli</i>	10 ⁻⁴ M	10 ⁻⁵ M	10 ⁻⁶ M	Rv control
1	19	18	-Ve	4+	4+	4+
2	16	16	-Ve	4+	4+	4+
3	16	20	-Ve	2+	4+	4+
4	13	19	-Ve	-Ve	4+	4+
5	15	18	-Ve	-Ve	4+	4+
6	18	20	-Ve	+	4+	4+
7	16	18	-Ve	2+	4+	4+
8	14	17	+	2+	4+	4+
9	20	22	-Ve	2+	4+	4+
10	20	16	0	3+	4+	4+
11	20	16	0	3+	4+	4+
12	17	20	-Ve	2+	4+	4+
13	16	16	0	3+	4+	4+
14	-Ve	-Ve	+	3+	4+	4+
15	21	18	Sl	4+	4+	4+
16	19	16	-Ve	4+	4+	4+
17	18	28	-Ve	4+	4+	4+
18	16	28	-Ve	4+	4+	4+
19	15	15	+	4+	4+	4+
20	17	17	-Ve	4+	4+	4+
21	15	18	-Ve	-Ve	4+	4+
22	16	17	-Ve	-Ve	4+	4+
23	17	18	+	2+	4+	4+
24	21	16	0	4+	4+	4+
25	20	16	0	4+	4+	4+
26	20	16	0	4+	4+	4+
27	18	16	0	4+	4+	4+
28	-Ve	-Ve	0	+	4+	4+
29	-Ve	-Ve	0	+	4+	4+
30	-Ve	-Ve	+	2+	4+	4+
31	-Ve	-Ve	3+	3+	4+	4+
32	-Ve	-Ve	0	+	4+	4+
5% Phenol control			0	=no growth		
value	26	20	+ to 4+	=various stages of growth		
Ethanol	-Ve	-Ve	Sl	=slight growth		

a) All the compounds were tested *in vitro*.

It is apparent from the results (Tables II and IV) that, with reference to carrier moiety, the activity increases in the following order:

quinazoline < acridine < quinoline < quinaldine.

The size of the side chain does not seem to influence the activity profoundly, and its effect is not of much significance. It is noteworthy that the bis-amines are more active against *S. aureus* and *E. coli* than against *M. tuberculosis*, however, the corresponding sulfonamides exhibit more antitubercular activity. The nature of the substitution in the aromatic sulfonyl group, in the case of sulfonamides, influences the antitubercular property and replacement of a hetero group, such as acetamido or chlorine in the *para* position by a methyl group enhances the antitubercular activity.

Experimental

Infrared spectrum was determined by Perkin-Elmer-257 spectrophotometer, using a nujol film. Melting points are uncorrected.

N,N'-Bis(4-quinolyl-, 4-quinaldinyl-, 4-quinazolinyl-, or 9-acridinyl)polymethylenediamine (I)—Bis-amine dihydrochloride^{11,12} was basified with 10–15% NaOH and the precipitated oil was taken up in chloroform, washed with distilled water, dried over anhydrous potassium carbonate, stripped of solvent to afford a pasty mass, which solidified on scratching. It was recrystallized from chloroform. Yields and melting points of the respective compounds are given in Table I. IR (Nujol): 3300 cm⁻¹ (NH).

N,N'-Bis(4-quinolyl-, 4-quinaldinyl-, 4-quinazolinyl-, 9-acridinyl-, or arenesulfonyl)polymethylenediamine (II)—Bis-amine and aromatic sulfonyl chloride, taken in the molar proportion of 1:2, were dissolved in dry pyridine using 10 ml of pyridine/g of the bis-amine, and the mixture was gently heated for 3–4 hr, at 130–135°. After the reaction was over, solvent was removed under reduced pressure, and the residue was triturated with cold water to get a solid, which on crystallization from hot water gave analytical sample. Yields and physical constants are given in Table III.

Acknowledgement Thanks are due to Dr. A.K. Mukerjee, Applied Chemistry Section, Institute of Technology, B.H.U., Varanasi, for his interest in this work. The biological screening of the compounds was carried out under the direction of Dr. T. Ramakrishnan, Indian Institute of Science, Bangalore, which is gratefully acknowledged. Competent technical assistance by Mr. V.N. Muley, Mr. P.S. Gurjar, and R.C. Bipin is appreciated.

[Chem. Pharm. Bull.
23(8)1873–1879(1975)]

UDC 547.867.04 : 547.823.057

Synthesis of Methylpyridine Derivatives. XXX.¹⁾ Reaction of Active Methylene Compounds with 6-Methyl-2-phenyl-4H-1,3-oxazin-4-one and 2-Ethoxy-2,6-dimethyl-2H-1,3-oxazin-4(3H)-one to give 3-Acetylpyridone Derivatives²⁾

TETSUZO KATO, YUTAKA YAMAMOTO, and MASATSUGU KONDO

Pharmaceutical Institute, Tohoku University³⁾

(Received September 27, 1974)

Reaction of 6-methyl-2-phenyl-4H-1,3-oxazin-4-one (I) with active methylene compounds, such as diethyl malonate (IIIa), acetylacetone (IIIb), cyclohexane-1,3-dione (IIIc), malononitrile (IIId), ethyl acetoacetate (IIIe), ethyl cyanoacetoacetate (IIIf), and cyanoacetophenone (IIIg) afforded pyridone derivatives, such as 3-acetyl-5-ethoxycarbonyl-4-hydroxy-6-phenyl-2(1H)-pyridone (IVa), 3,5-diacetyl-4-methyl-6-phenyl-2(1H)-pyridone (IVb), 4-acetyl-1-phenyl-6,7-dihydro-3,8-(2H, 5H)-isoquinolinedione (IVc), 3-acetyl-4-amino-5-cyano-6-phenyl-2(1H)-pyridone (IVd), 3-acetyl-5-ethoxycarbonyl-4-methyl-6-phenyl-2(1H)-pyridone (IVe), 3-acetyl-5-cyano-4-hydroxy-6-phenyl-2(1H)-pyridone (IVf), and 3-acetyl-5-cyano-4,6-diphenyl-2(1H)-pyridone (IVg), respectively.

Similarly, 2-ethoxy-2,6-dimethyl-2H-1,3-oxazin-4(3H)-one (II) reacted with IIIa–t under the same conditions to give the corresponding pyridone derivatives (Va–f).

We have previously reported that diketene reacted with imidates to give 1,3-oxazin-4-one derivatives.^{4,5)} The present paper reports a continuation of our study of those products related to the reaction with active methylene compounds to give pyridine derivatives.

Oxazines used in this reaction are 6-methyl-2-phenyl-4H-1,3-oxazin-4-one (I) and 2-ethoxy-2,6-dimethyl-2H-1,3-oxazin-4(3H)-one (II), prepared by the reaction of diketene with ethyl benzimidate and ethyl acetimidate, respectively.

- 1) Part XXIX: T. Kato and N. Niitsuma, *Yakugaku Zasshi*, **94**, 766 (1974).
- 2) This forms Part LXX of "Studies on Ketene and Its Derivatives," by T. Kato.
- 3) Location: Aobayama, Sendai, 980, Japan.
- 4) T. Kato and Y. Yamamoto, *Chem. Pharm. Bull.* (Tokyo), **15**, 1334 (1967).
- 5) T. Kato, H. Yamanaka, Y. Yamamoto, and M. Kondo, *Yakugaku Zasshi*, **92**, 886 (1972).