[Chem. Pharm. Bull.
28(1) 157—162 (1980)]

Introduction of Carbon Substituents into Pyrimidine and Purine Nucleosides by Sulfur Extrusion (Nucleosides and Nucleotides. XXX¹⁾)

AKIRA YAMANE, HIDEO INOUE, and TOHRU UEDA

Faculty of Pharmaceutical Sciences, Hokkaido University²⁾

(Received June 29, 1979)

Treatment of 4-thiouridine with phenacyl bromide, bromoacetone, and ethyl bromoacetate gave the corresponding 4-thioalkylcarbonyl derivatives. The sulfur extrusion reactions of these compounds afforded ribosides of 4-phenacylidene-2(3*H*)-pyrimidinone, 4-acetonylidene-2(3*H*)-pyrimidinone, and 2-pyrimidinone-4-acetic ester, respectively. The ease of sulfur extrusion depends on the electron-withdrawing ability of the carbonyl group attached to the 4-*S*-methylene group. Sulfur extrusion reactions starting from 6-thioinosine similarly gave the ribofuranosides of 6-phenacylpurine and 6-acetonylpurine. These purine ribosides exist mainly as the enol form rather than the keto form.

Keywords—sulfur extrusion reaction; thionucleosides; nucleosides; triphenylphosphine; UV; NMR

Few studies have been reported on the synthesis of nucleosides with carbon substituents in place of amino or oxo substituents on the base moieties of naturally occurring nucleosides. Procedures reported so far involve the glycosylation of 6-alkylpurines³⁾ or 4-methyl-2-pyrimidinone⁴⁾ and the photo-addition of methanol to ribofuranosylpurine.⁵⁾ 9- β -D-Ribofuranosyl-6-methylpurine and -1,6-dihydro-6-hydroxymethylpurine have been reported to be strong inhibitors of adenosine deaminases.⁵⁾ In preceding papers we have reported facile substitutions of a methylsulfonyl group with various carbon nucleophiles at the 2, 6, or 8 position of purine nucleosides to give the corresponding C-substituted derivatives.^{1,6,7)} We wish to report an alternative approach to the synthesis of C-substituted pyrimidine and purine nucleosides which utilizes the sulfur extrusion reaction established by Eschenmoser and co-workers.⁸⁾ A preliminary report has already appeared.⁹⁾ Essentially similar results were reported¹⁰⁾ after the completion of our experiments. As 4-thiouridine and 6-thioinosine contain a thioamide group, these nucleosides may serve as substrates for the conversion of a thioamide to a vinylogous amide, as reported in the case of 4-*p*-bromophenacylthiouracil, yielding 4-phenacyl-2-pyrimidinone.¹¹⁾ 4-Thiouridine (**1**), readily prepared by the sulfhydrolysis of cytidine,¹²⁾ was treated with phenacyl bromide, bromoacetone, and ethyl bromoacetate in aqueous solution in the presence of sodium hydrogen carbonate to give the corresponding alkylthio derivatives (**2**—**4**) in high yields. 2',3'-*O*-Isopropylidene-4-thiouridine (**1a**) also gave the 4-thioacetate derivative (**5**).

- 1) Part XXIX: A. Yamane, A. Matsuda, and T. Ueda, *Chem. Pharm. Bull.*, **28**, 150 (1980).
- 2) Location: *Kita-12, Nishi-6, Kita-ku, Sapporo, 060, Japan*.
- 3) J.A. Montgomery and K. Hewson, *J. Med. Chem.*, **11**, 48 (1967).
- 4) R.S. Klein, I. Wempen, K.A. Watanabe, and J.J. Fox, *J. Org. Chem.*, **35**, 2330 (1970).
- 5) B. Evans and R. Wolfenden, *J. Am. Chem. Soc.*, **92**, 4751 (1970).
- 6) A. Matsuda, Y. Nomoto, and T. Ueda, *Chem. Pharm. Bull.*, **27**, 183 (1979).
- 7) A. Yamane, Y. Nomoto, A. Matsuda, and T. Ueda, *Nucleic Acids Res. Special Publ.*, No. 5, s309 (1978), Nagoya.
- 8) M. Roth, P. Dubs, E. Götschi, and A. Eschenmoser, *Helv. Chim. Acta*, **54**, 710 (1971).
- 9) A. Yamane, H. Inoue, and T. Ueda, Abstract Paper 97th Meeting Pharm. Soc. Japan, Vol. II, Tokyo, 1977, p. 136.
- 10) H. Vorbrüggen and K. Krolkiewicz, *Angew. Chem. Internl. Ed.*, **15**, 689 (1976).
- 11) Footnote 9 of reference 8.
- 12) T. Ueda, K. Miura, M. Imazawa, and K. Odajima, *Chem. Pharm. Bull.*, **22**, 2377 (1974).

Chart 1

Heating the 4-phenacylthio derivative (2) in pyridine at reflux temperature gave a new product (6) which contained no sulfur. The ultraviolet (UV) absorption spectrum of the product exhibited a maximum at 375.5 nm, reflecting the elongation of the conjugation in the chromophore. In the nuclear magnetic resonance (NMR) spectrum an exchangeable proton at 13 ppm and a methine proton were observed instead of the methylene. Thus, the structure was assigned as 1-β-D-ribofuranosyl-4-phenacylidene-2(3H)-pyrimidinone. It should be noted that the sulfur extrusion in 2 occurred so readily that the addition of a sulfur trap, such as triethyl phosphite, was unnecessary. 4-Acetylthiouridine (3) was rather stable, and the elimination of sulfur took place on addition of triphenylphosphine and potassium *t*-butoxide in refluxing pyridine to give the 4-acetyliden-2(3H)-pyrimidinone (7). The spectroscopic characteristics were similar to those of 6. The difference in the ease of elimination of sulfur between 2 and 3 is explained by the difference of the acidity of the methylene group, that of the former compound being more acidic. This is further exemplified in the reaction of 4-thiouridine and diethyl bromomalonate. Treatment of 1a with diethyl bromomalonate in dimethylformamide (DMF) in the presence of potassium carbonate at room temperature afforded 4-bis(ethoxycarbonylmethyl)-2(3H)-pyrimidinone riboside (8). The reaction conditions of *S*-alkylation led to simultaneous sulfur extrusion in this instance.

The sulfur extrusion of the ethoxycarbonylmethylthio derivative (4) proceeded sluggishly, so the sugar hydroxyl group was protected prior to the reaction. Compound 5 was acetylated and treated with triphenylphosphine and sodium hydride in tetrahydrofuran (THF). The product 9 was obtained as an amorphous material. On saponification and mild acid treatment of 9, 1-(2,3-O-isopropylidene-β-D-ribofuranosyl)-4-methyl-2-pyrimidinone (10) was obtained in an amorphous form. The parent nucleoside is already known⁴⁾ and the spectral data of 10 were in good accord with those of 4-methyl-2-pyrimidinone riboside.⁴⁾

Treatment of 6-thiinosine with phenacyl bromide, bromoacetone, and ethyl bromoacetate gave the corresponding *S*-alkylated derivatives (11–13) in high yields. The extrusion reaction of the alkylthiopurines proceeded rather slowly. Treatment of 11 with triphenylphosphine did give the expected product, though it was difficult to isolate. Treatment of the benzoylated derivative (14) with triphenylphosphine and potassium *t*-butoxide in THF

Chart 2

gave the 6-phenacyl derivative (**15**). The debenzoylation of **15** with methanolic ammonia afforded 9- β -D-ribofuranosyl-6-phenacyl-purine (**16**) in high yield. The tautomeric form of **16** was assigned as the enol form rather than the keto form, since **16** gave intense color with ferric chloride solution. The NMR spectrum of **16** showed a methine proton signal at 5.56 ppm. Although the enol proton could not be detected due to the presence of the sugar hydroxyls, the absence of an N^1 -proton signal (usually appearing at >10 ppm) indicates the structure **16**.

The sulfur extrusion reaction of **12** with triphenylphosphine and potassium *t*-butoxide in pyridine gave the 6-acetonyl-purine riboside (**17**), though in low yield. Compound **17** also exhibited intense coloration with ferric chloride solution. The NMR spectrum of **17** indicated the presence of the other tautomer with an N^1 -proton (see "Experimental"). Attempted sulfur extrusion of **13** under various conditions met with little success. The desired product, 6-ethoxycarbonylmethyl-purine riboside, was obtained by the alternative route already reported.¹⁾

In conclusion, the extrusion of sulfur from alkylthio-nucleosides proceeded well when the carbonyl moiety attached to the thiomethylene group was strongly electron-withdrawing. The reduction of the keto group of the product should provide a new route for the preparation of alkylpyrimidine or -purine nucleosides. Reactions utilizing the keto group may yield various functionalized nucleosides. Studies along these lines are in progress. Biological tests of the newly synthesized C-substituted nucleosides are also under way, and preliminary tests of cytotoxic activity against leukemia L-5178Y cells indicated that 6-acetonyl-purines are moderately effective.¹³⁾

Experimental

Melting points were determined with a Yanaco MP-3 melting point apparatus and are uncorrected. UV spectra were recorded on a Shimadzu UV-300 recording spectrophotometer. NMR spectra were taken with a JEOL JNM-FX 100FT NMR spectrometer. Mass spectra were taken with a Hitachi RMU-7 mass spectrometer. IR spectra were measured on a Hitachi 215 spectrophotometer. Thin-layer chromatography was carried out on Merck TLC plates (silica gel 60F₂₅₄, pre-coated). Silica gel for column chromatography was Wakogel C-200. Ribonucleosides used in this work were purchased from Yamasa Shoyu Co., Ltd.

4-Phenacylthiouridine (2)—4-Thiouridine (**1**, 520 mg) and NaHCO₃ (184 mg, 1 eq) were dissolved in 3 ml of H₂O and phenacyl bromide (597 mg, 1.5 eq) in 3 ml of EtOH was added. The solution was stirred for 20 min while 20 ml of EtOH was added. The separated crystals were collected and recrystallized from MeOH to give 582 mg (77%) of **2**, mp 161.5–162.5°. *Anal.* Calcd for C₁₇H₁₈N₂O₆S: C, 53.97; H, 4.80; N,

13) The tests were performed by Dr. H. Yoshino and co-workers, Yamasa Shoyu Co.

7.41; S, 8.47. Found: C, 53.41; H, 4.78; N, 7.43; S, 8.52. UV $\lambda_{\max}^{\text{H}_2\text{O}}$ nm (ϵ): 300 (14900), 250 (15700). $\lambda_{\max}^{\text{H}_2\text{O}}$: 276 (11400), 230 (9700). NMR (DMSO- d_6) δ : 8.35 (d, 1, 6-H), 8.22–7.60 (m, 5, Ph), 6.60 (d, 1, 5-H, $J=9$ Hz), 5.78 (d, 1, 1'-H, $J=2.3$ Hz), 4.86 (s, 2, S-CH₂-).

4-Acetonylthiuridine (3)—A solution of 10.40 g of 1 and 3.67 g of NaHCO₃ in H₂O (150 ml)–EtOH (20 ml) was treated with 8.22 g of bromoacetone (80% purity, 1.2 eq), and the solution was stirred for 1 hr at room temperature. The solvent was concentrated *in vacuo* and the residue was crystallized from EtOH to give 10.50 g (83%) of 3, mp 142–144°. Anal. Calcd. for C₁₂H₁₆N₂O₆S: C, 45.57; H, 5.10; N, 8.86; S, 10.26. Found: C, 45.58; H, 5.12; N, 8.81; S, 10.36. UV $\lambda_{\max}^{\text{H}_2\text{O}}$ nm (ϵ): 301 (13300), 265 (sh, 8300). $\lambda_{\min}^{\text{H}_2\text{O}}$: 237 (2600). NMR (DMSO- d_6) δ : 8.32 (d, 1, 6-H), 6.55 (d, 1, 5-H, $J=8$ Hz), 5.80 (d, 1, 1'-H, $J=2.3$ Hz), 4.13 (s, 2, -SCH₂-), 2.29 (s, 3, COCH₃).

4-Ethoxycarbonylmethylthiuridine (4)—Ethyl bromoacetate (334 mg, 1.3 eq) was added to a solution of 390 mg of 1 and 138 mg of NaHCO₃ in H₂O (3 ml)–EtOH (2 ml), and the solution was stirred for 2 hr at room temperature. After adding 86 mg of the bromoacetate and stirring for another 2 hr, the solvent was removed *in vacuo* and the residue was taken up in a minimum amount of EtOH. The solution was applied to a column of silica gel (20 g) and the product was eluted with 8–10% MeOH in CHCl₃. The eluate was concentrated and the residue was crystallized from EtOH to give 278 mg (73%) of 4, mp 135–136.5°. Anal. Calcd. for C₁₃H₁₈N₂O₇S: C, 45.09; H, 5.24; N, 8.09; S, 9.26. Found: C, 44.91; H, 5.26; N, 7.86; S, 9.07. UV $\lambda_{\max}^{\text{H}_2\text{O}}$: 330 nm shoulder at 265 nm. $\lambda_{\min}^{\text{H}_2\text{O}}$: 237 nm. NMR (DMSO- d_6) δ : 8.46 (d, 1, 6-H), 6.96 (d, 1, 5-H, $J=7$ Hz), 6.15 (d, 1, 1'-H, $J=3$ Hz), 4.54 (q, 2, -OCH₂CH₃), 4.31 (s, 2, -SCH₂-), 1.53 (t, 3, -OCH₂CH₃, $J=7$ Hz).

2',3'-O-Isopropylidene-4-thiuridine (1a)—Cytidine (34.15 g) was suspended in 1400 ml of acetone, and 20 ml of 60% HClO₄ was added. The suspension was stirred for 2 hr and the clear solution was neutralized with anhydrous K₂CO₃. The filtrate was concentrated, and the residue was dissolved in 240 ml of H₂O. The solution (40 ml) was added to a mixture of 50 ml of pyridine and 80 ml of liquid H₂S in a tube under cooling with dry ice-acetone. The tube was sealed and kept for 3 days at 50°. After removal of H₂S by evaporation and bubbling the residue through with N₂ gas, the solution was concentrated and the residue was taken up in EtOH. The insoluble material was removed by filtration and the filtrate was concentrated. The residue was crystallized from EtOH to give 1a. Six runs of this reaction gave 1a in a yield of 31.85 g (75.7%), mp 182–183° (mp 170–172°¹⁴).

1-(2,3-O-Isopropylidene- β -D-ribofuranosyl)-4-ethoxycarbonylmethylthiuracil (5)—A solution of 4.50 g of 1a in 80 ml of DMF was treated with 1.38 g of NaHCO₃ in 10 ml of H₂O and 2.75 g of ethyl bromoacetate. The solution was stirred for 2 hr at room temperature. The solvent was removed *in vacuo* and the residue was dissolved in 100 ml of CHCl₃ and washed with H₂O. The organic layer was concentrated and the residue was crystallized from EtOH to give 5.21 g (90%) of 5, mp 178–180°. Anal. Calcd. for C₁₆H₂₂N₂O₇S: C, 49.74; H, 5.74; N, 7.25; S, 8.28. Found: C, 49.65; H, 5.70; N, 7.14; S, 8.20. NMR (DMSO- d_6) δ : 8.08 (d, 1, 6-H), 6.50 (d, 1, 5-H, $J=8.3$ Hz), 5.82 (s, 1, 1'-H), 4.12 (q, 2, OCH₂CH₃), 4.02 (s, 2, SCH₂), 1.48, 1.30 (s, 3+3, CMe₂), 1.22 (t, 3, OCH₂CH₃, $J=6.8$ Hz).

1- β -D-Ribofuranosyl-4-phenacylidene-2(3H)-pyrimidinone (6)—Compound 2 (3.78 g) was dissolved in 20 ml of pyridine and the solution was refluxed for 1 hr. The solvent was then removed *in vacuo* and the residue was taken up in H₂O and concentrated again. Concentration from H₂O was repeated several times and the final residue was taken up in a minimum amount of MeOH. Silica gel was added to the solution and the solvent was removed to leave a powder. This was applied to the top of a silica gel column (120 g) and the column was eluted with 10% MeOH in CHCl₃. The eluate containing the product was concentrated and the residue was crystallized from EtOH to give 1.60 g (46%) of 6, mp 186–187°. Anal. Calcd. for C₁₇H₁₈N₂O₆: C, 58.95; H, 5.24; N, 8.09. Found: C, 58.52; H, 5.32; N, 7.99. UV $\lambda_{\max}^{\text{H}_2\text{O}}$ nm (ϵ): 375.5 (15200), 262.5 (2800). $\lambda_{\min}^{\text{H}_2\text{O}}$: 300 (780), 242 (2100). $\lambda_{\max}^{\text{NaOH}}$ nm (ϵ): 388 (17000), 250 (4600). $\lambda_{\min}^{\text{NaOH}}$: 320 (1200), 234 (4000). NMR (DMSO- d_6) δ : 13.12 (bs, 1, N³-H), 7.82 (d, 1, 6-H), 8.10–7.45 (m, 5, Ph), 6.25 (s, 1, 4-CH=), 6.16 (d, 1, 5-H, $J=7.5$ Hz), 5.88 (d, 1, 1'-H, $J=4.5$ Hz). MS m/e : 346 (M⁺), 257 (M-89), 214 (B+1), 213 (B).

1- β -D-Ribofuranosyl-4-acetonilidene-2(3H)-pyrimidinone (7)—A solution of 1.58 g of 3 and 3.93 g of triphenylphosphine in 100 ml of pyridine was treated with 244 mg of *t*-BuOK in 20 ml of *t*-BuOH, and the solution was refluxed for 5 min. After cooling, the solution was neutralized by the addition of AcOH and the solvent was removed *in vacuo*. The residue was taken up in 100 ml of H₂O and washed with CHCl₃. The aqueous layer was concentrated and the residue was applied to a column of silica gel (60 g). The eluent with 10% MeOH in CHCl₃ was collected and concentrated, then the residue was crystallized from EtOH–AcOEt to give 717 mg (50%) of 7, mp 168–170°. Anal. Calcd. for C₁₈H₁₆N₂O₆: C, 50.70; H, 5.67; N, 9.86. Found: C, 50.71; H, 5.66; N, 9.83. UV $\lambda_{\max}^{\text{H}_2\text{O}}$ nm (ϵ): 370 (sh, 20700), 351 (265000), 338 (sh, 23400), 265 (3900). $\lambda_{\min}^{\text{H}_2\text{O}}$: 284 (3000). $\lambda_{\max}^{\text{NaOH}}$ nm (ϵ): 361 (38500), 280–260 (plateau, 5600). NMR (DMSO- d_6) δ : 12.60 (bs, 1, N³-H), 7.60 (d, 1, 6-H), 5.94 (d, 1, 5-H, $J=8.3$ Hz), 5.84 (d, 1, 1'-H, $J=4.5$ Hz), 5.38 (s, 1, 4-CH=), 2.06 (s, 3, CH₃). MS m/e : 284 (M⁺), 152 (B+1), 151 (B).

14) V.N. Shibaev, M.A. Grachev, and S.M. Spiridonova, "Synth. Proced. Nucleic. Acid. Chem.," Vol. 1, John Wiley and Sons, Inc., 1968, p. 503.

1-(2,3-*O*-Isopropylidene- β -D-ribofuranosyl)-4-bis(ethoxycarbonyl)methylene-2(3*H*)-pyrimidinone (8)—

A mixture of 2.0 g of **1a**, 966 mg of K_2CO_3 , and 1.91 g of diethyl bromomalonate in 20 ml of DMF was stirred for 1.5 hr at room temperature. AcOEt (80 ml) was added to the mixture and the solution was partitioned with 50 ml of H_2O . The organic layer was dried over Na_2SO_4 , concentrated *in vacuo*, and the residue was crystallized from EtOH–AcOEt to give 1.30 g (46%) of **8**, mp 123–124°. *Anal.* Calcd for $C_{19}H_{26}N_2O_9$: C, 53.51; H, 6.15; N, 6.57. Found: C, 53.42; H, 6.19; N, 6.43. UV $\lambda_{max}^{H_2O}$ nm (ϵ): 355 (sh, 14000), 336 (22800), 325 (sh, 22000), 237 (6300). $\lambda_{min}^{H_2O}$: 277 (2700). $\lambda_{max}^{0.1N NaOH}$ nm (ϵ): 344 (26700), 257 (7200). $\lambda_{min}^{0.1N NaOH}$: 296 (3600). NMR ($CDCl_3$) δ : 11.88 (bs, 1, N³-H), 7.06 (d, 1, 6-H), 6.38 (dd, 1, 5-H, $J=8.3$ and 1.5 Hz), 5.52 (d, 1, 1'-H), $J=2.3$ Hz), 4.22 (q, 4, $-OCH_2CH_3$), 1.54, 1.34 (s, 3+3, CH_3), 1.38 (t, 6, $-OCH_2CH_3$).

1-(5-*O*-Acetyl-2,3-*O*-isopropylidene- β -D-ribofuranosyl)-4-ethoxycarbonylmethyl-2-pyrimidinone (9)—

Compound **5** (100 mg) and Ac_2O (0.5 ml) in 3 ml of pyridine were kept overnight at room temperature. The solvent was then removed and the residue was taken up in $CHCl_3$, washed with H_2O , and dried over Na_2SO_4 . After removal of the solvent, the residue was dissolved in 5 ml of THF, then 136 mg of triphenylphosphine and 25 mg of 50% NaH were added and the mixture was heated under reflux for 1 hr. After neutralization of the solution with AcOH the solvent was removed *in vacuo* and the residue was applied to a column of silica gel (10 g). The eluent with 10% AcOEt in C_6H_6 was concentrated to leave 58 mg (56%) of **9** in an amorphous form. UV $\lambda_{max}^{H_2O}$ nm: 321, 267. $\lambda_{min}^{H_2O}$: 277, 227. $\lambda_{max}^{0.1N NaOH}$ nm: 338, 267. $\lambda_{min}^{0.1N NaOH}$: 272, 235. NMR ($CDCl_3$) δ : 10.86 (bs, N³-H), 6.68 (d, 1, 6-H), 5.58 (d, 1, 5-H, $J=7.5$ Hz), 5.52 (bs, 1, 1'-H), 5.1–4.8 (m, 2, 2', 3'-H), 4.71 (s, 1, 4-CH=), 4.14 (q, 2, $-OCH_2CH_3$), 4.28 (bs, 3, 4', 5'-H), 2.06 (s, 3, Ac), 1.54, 1.34 (s, 3+3, CH_3), 1.26 (t, 3, $-OCH_2CH_3$).

1-(2,3-*O*-Isopropylidene- β -D-ribofuranosyl)-4-methyl-2-pyrimidinone (10)—

Compound **9** (58 mg) was dissolved in a mixture of 0.1 *N* NaOH (2 ml) and EtOH (3 ml), and kept overnight at room temperature. The solution was acidified by the addition of 5 ml of 0.1 *N* HCl and kept overnight at 10°. The solution was then neutralized with 1 *N* NaOH and the solvent was evaporated off. The residue was applied to a column of silica gel (10 g). The eluent with 5% EtOH in $CHCl_3$ was concentrated to give an amorphous residue **10** (23 mg, 56%). UV $\lambda_{max}^{H_2O}$ 297 nm. $\lambda_{min}^{H_2O}$ 240 nm. $\lambda_{max}^{0.1N HCl}$ 309 nm. $\lambda_{min}^{0.1N HCl}$ 250 nm. NMR ($DMSO-d_6$) δ : 8.12 (d, 1, 6-H), 6.39 (d, 1, 5-H, $J=7.2$ Hz), 5.82 (d, 1, 1'-H, $J=1.5$ Hz), 2.28 (s, 3, 4- CH_3), 1.49, 1.29 (s, 3+3, CH_3).

6-Phenacylthioinosine (11)—

6-Thioinosine¹⁵ (2.84 g) was dissolved in H_2O (50 ml) and EtOH (10 ml) containing 924 mg of $NaHCO_3$ by warming. Phenacylbromide (2.39 g) was added to the cooled solution and the mixture was stirred for 2 days at room temperature. After removal of the solvent, the residue was taken up in a minimum amount of EtOH and applied to a column of silica gel (90 g). The eluent with 10–15% MeOH in $CHCl_3$ was collected and concentrated. The residue was crystallized from EtOH to give 3.06 g (76%) of **11**, mp 103–105°. *Anal.* Calcd for $C_{18}H_{18}N_4O_5S$: C, 53.73; H, 4.51; N, 13.93; S, 7.95. Found: C, 53.45; H, 4.43; N, 13.76; S, 8.13. UV $\lambda_{max}^{H_2O}$ nm (ϵ): 292 (sh, 20300), 284 (21500), 252 (14300), 226 (13700). $\lambda_{min}^{H_2O}$: 265 (11900), 236 (10100), 222 (13500). NMR ($DMSO-d_6$) δ : 8.80 (s, 1, 8-H), 8.68 (s, 1, 2-H), 8.3–7.6 (m, 5, Ph), 6.05 (d, 1, 1'-H, $J=5$ Hz), 5.12 (s, 2, SCH_2).

6-Acetylthioinosine (12)—

6-Thioinosine (3.41 g) was dissolved in H_2O (50 ml) and EtOH (5 ml) containing 1.11 g of $NaHCO_3$ by warming. Bromoacetone (2.41 g) was added to the solution and the mixture was stirred overnight. After work-up as described above, a crystalline product (**12**, 3.48 g, 85%) was obtained from EtOH, mp 138–140°. *Anal.* Calcd for $C_{13}H_{16}N_4O_5S$: C, 45.88; H, 4.74; N, 16.47; S, 9.40. Found: C, 45.82; H, 4.64; N, 16.32; S, 9.31. UV $\lambda_{max}^{H_2O}$ nm (ϵ): 281 (12700), 218 (8300). $\lambda_{min}^{H_2O}$: 240 (1800), 208 (7500). NMR ($DMSO-d_6$) δ : 8.82 (s, 1, 8-H), 8.76 (s, 1, 2-H), 6.08 (d, 1, 1'-H, $J=5.5$ Hz), 4.38 (s, 2, SCH_2), 2.34 (s, 3, $COCH_3$), IR (KBr): 1720 cm^{-1} ($COCH_3$).

6-Ethoxycarbonylmethylthioinosine (13)—

6-Thioinosine (4.26 g) and 3.0 g of ethyl bromoacetate were treated as described above and the product **13** (4.26 g, 77%) was crystallized from EtOH, mp 131.5–133°. *Anal.* Calcd for $C_{14}H_{18}N_4O_6S$: C, 45.40; H, 4.90; N, 15.13; S, 8.64. Found: C, 45.32; H, 4.86; N, 15.15; S, 8.71. UV $\lambda_{max}^{H_2O}$ nm (ϵ): 287 (sh, 11800), 280 (13000), 227 (8500). $\lambda_{min}^{H_2O}$: 239 (1600). NMR ($DMSO-d_6$) δ : 8.76 (s, 1, 8-H), 8.72 (s, 1, 2-H), 6.05 (d, 1, 1'-H, $J=5.3$ Hz), 4.26 (s, 2, SCH_2), 4.15 (q, 2, $-OCH_2CH_3$), 1.20 (t, 3, $-OCH_2CH_3$, $J=6.9$ Hz).

2',3',5'-Tri-*O*-benzoyl-6-phenacylthioinosine (14)—

2',3',5'-Tri-*O*-benzoyl-6-thioinosine¹⁵ (5.96 g) was dissolved in 60 ml of DMF, then 1.52 g of K_2CO_3 and 2.09 g of phenacyl bromide were added. The solution was stirred for 2 hr at room temperature and poured into ice-water. The precipitate was separated and dried to give 6.63 g (94%) of crude **14**, which was used directly for the next step.

9-(2,3,5-Tri-*O*-benzoyl- β -D-ribofuranosyl)-6-phenacylurine (15)—

Crude **14** (500 mg), triphenylphosphine (367 mg) and 50% NaH (67 mg) were taken up in 20 ml of THF. After heating under reflux for 2 hr, the solution was concentrated and the residue was taken up in a small amount of C_6H_6 . This solution was applied to a column of silica gel (20 g) and eluted with 10% AcOEt in C_6H_6 . The product **15** was obtained as a foam, 250 mg (74%). UV λ_{max}^{EtOH} : 373, 230 nm. λ_{min}^{EtOH} : 297 nm. This was used for the next step without purification.

15) J.J. Fox, I. Wempen, A. Hampton, and I.L. Doerr, *J. Am. Chem. Soc.*, **80**, 1669 (1958).

9- β -D-Ribofuranosyl-6-phenaclypurine (16)—Compound 15 (350 mg) was dissolved in 20 ml of MeOH saturated with NH_3 , and kept overnight at room temperature. Nitrogen was bubbled through the solution for 20 min while crystals separated. The precipitate was collected and washed well with MeOH- CHCl_3 to give 179 mg (93%) of 16, mp 238–240°. *Anal.* Calcd for $\text{C}_{18}\text{H}_{18}\text{N}_4\text{O}_5$: C, 58.37; H, 4.90; N, 15.13. Found: C, 58.37; H, 4.87; N, 15.04. UV $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ nm (ϵ): 386 (7500), 252 (3500). $\lambda_{\text{min}}^{\text{H}_2\text{O}}$: 305 (600). $\lambda_{\text{max}}^{0.1\text{N NaOH}}$: 384 (10500), 245 (3900). $\lambda_{\text{min}}^{0.1\text{N NaOH}}$: 305 (600), 240 (3800). NMR ($\text{DMSO}-d_6$) δ : 8.62 (s, 1, 8-H), 8.54 (s, 1, 2-H), 8.0–7.5 (m, 5, Ph), 6.56 (s, 1, 6-CH=), 6.02 (d, 1, 1'-H, $J=5.3$ Hz). MS m/e : 370 (M^+), 239 ($\text{B}+2$), 238 ($\text{B}+1$), 237 (B). Compound 16 gave a violet color with FeCl_3 solution.

9- β -D-Ribofuranosyl-6-acetonylpurine (17)—Compound 12 (1.70 g) and triphenylphosphine (3.93 g) were taken up in 100 ml of pyridine, then 560 mg of t -BuOK in 30 ml of t -BuOH was added. The solution was refluxed for 30 min, neutralized with AcOH, and concentrated. The residue was partitioned with H_2O and CHCl_3 . The aqueous layer was concentrated and the residue was applied to a column of silica gel (60 g). The eluent with 10% MeOH in CHCl_3 was concentrated and the residue was crystallized from EtOH- H_2O to give 445 mg (28%) of 17, mp 163–165°. *Anal.* Calcd for $\text{C}_{13}\text{H}_{16}\text{N}_4\text{O}_5 \cdot 1/4\text{H}_2\text{O}$: C, 49.91; H, 5.27; N, 17.90. Found: C, 49.86; H, 5.35; N, 17.80. UV $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ nm (ϵ): 360 (12500), 345 (sh, 9000), 268 (7200). $\lambda_{\text{min}}^{\text{H}_2\text{O}}$: 288 (1500), 226 (3800). $\lambda_{\text{max}}^{0.1\text{N NaOH}}$: 359 (25700), 241 (8800). $\lambda_{\text{min}}^{0.1\text{N NaOH}}$: 283 (2600), 226 (8200). NMR ($\text{DMSO}-d_6$) δ : 10.72 (bs, 0.3, $\text{N}^1\text{-H}$), 8.94, 8.56 (each s, 1, 8-H, ratio 1:2.5), 8.86, 8.47 (each s, 1, 2-H, ratio 1:2.5), 6.09, 5.90 (each d, 1, 1'-H, $J_{1',2'}=6.0$ and 5.3 Hz, ratio 1:2.5), 2.29, 2.11 (each s, 3, CH_3 , ratio 1:2.5). The presence of a minor tautomer resulted in complex signals for other protons, and assignment was impossible. MS m/e : 308 (M^+), 219 ($\text{M}-89$), 205 ($\text{B}+30$), 176 ($\text{B}+1$). Compound 17 showed a violet coloration with FeCl_3 solution.

Acknowledgement This work was supported in part by Grants-in-Aid for Scientific Research from the Ministry of Education, Science and Culture, and from the Ministry of Health and Welfare. The elemental analysis and NMR measurements were performed by the staff of the Central Analytical Room of this faculty, to whom our thanks are due.