Communications to the Editor

[Chem. Pharm. Bull.]
30(9)3424-3426(1982)

AN UNUSUAL TRANSFORMATION OF 5-AMINO-6-METHYL-3-PHENYL-4(3H)-
PYRIMIDONE WITH SELENIUM DIOXIDE: X-RAY CRYSTAL STRUCTURE OF
3-PHENYLCARBAMOYL-4-METHYL-1,2,5-SELENADIAZOLE

Taisei Ueda^a, Yoshitsugu Shibata^a, Jinsaku Sakakibara^{*,a},
Masatoshi Inoue^b, and Toshimasa Ishida^b

Faculty of Pharmaceutical Sciences, Nagoya City University^a,
Tanabe-dori, Mizuho-ku, Nagoya 467, Japan
Osaka College of Pharmacy^b, Kawai, Matsubara, Osaka 580, Japan

The reaction of 5-amino-6-methyl-3-phenyl-4(3H)-pyrimidone (1) with selenium dioxide in dioxane gave the unusual products: 6-phenyl-7(6H)-isoselenazolo[4,3-*d*]pyrimidone (2) and 3-phenylcarbamoyl-4-methyl-1,2,5-selenadiazole (3) in the ratio 1:2.

KEYWORDS—pyrimidine; pyrimidone; selenium dioxide; selenadiazole; isoselenazole; X-ray crystal structure; pyrazine

Selenium dioxide attacks a methyl group in the neighborhood of a ring nitrogen atom to give either an aldehyde or a carboxyl group. For instance, 2-methylpyridine gives a mixture of pyridine-2-carbaldehyde and pyridine-2-carboxylic acid.¹⁾ Recently, Yamanaka, *et al.*²⁾ reported the selective oxidations of 2,4-dimethyl-6-phenylpyrimidine, 4,6-dimethyl-2-phenylpyrimidine or 2,4-dimethylquinoline with selenium dioxide in dioxane to give 2-methyl-6-phenylpyrimidine-4-carbaldehyde, 6-methyl-2-phenylpyrimidine-4-carbaldehyde, or 4-methylquinoline-2-carbaldehyde. We now present an example of an unusual transformation of a pyrimidone by reaction with selenium dioxide. The reactions of some pyrimidones such as 6-methyl-3-phenyl-4(3H)-pyrimidone³⁾, 5-bromo-6-methyl-3-phenyl-4(3H)-pyrimidone³⁾, or 5-dimethylamino-6-methyl-3-phenyl-4(3H)-pyrimidone⁴⁾ with selenium dioxide in dioxane or xylene did not proceed, while the reaction of 5-amino-6-methyl-3-phenyl-4(3H)-pyrimidone⁵⁾ (1) with selenium dioxide gave two unusual products: 6-phenyl-7(6H)-isoselenazolo[4,3-*d*]pyrimidone (2) (mp 195-6°C) and 3-phenylcarbamoyl-4-methyl-1,2,5-selenazole (3) (mp 154-5°C). The structure of 2 was determined by ¹H-NMR and mass spectra. In the ¹H-NMR spectrum signals of C₆-methyl protons and C₅-NH₂ protons disappeared, and a new signal (δ 9.85) derived from =CH-Se- function was observed. The mass spectrum also indicated the presence of a selenium element in its molecule. Treatment of 2 with sodium borohydride in methanol gave 1. On the other hand, mass spectrum and elemental analyses of 3 led to the empirical formula C₁₀H₉N₃OSe. ¹³C-NMR and ¹H-NMR analyses of 3 suggested the elimination of C₂-carbon atom from 1. The reduction of 3 with sodium borohydride in methanol led to a compound of C₂₀H₁₈N₄O₂ whose structure was identified as 2,5-dimethyl-3,6-dianilidopyrazine(4). The structure of 3 was unambiguously determined by single-

crystal X-ray analysis as shown in Fig. 1, which also made possible a rationalization of its ^1H -NMR spectrum. The formation of 3 probably is explained as depicted in Chart 1. The selenium dioxide initially attacks the C_2 -methylene function of the resonant form of 1 to give labile aldehyde, then the elimination of the aldehyde group and the reaction of the resulting diimine with selenium dioxide occurs to give 3. This is the first known example of the transformation of a pyrimidone to a selenadiazole by reaction with selenium dioxide.

Chart 1

Table I. Yields (%) of 2 and 3 by the Reaction of 1 with Selenium Dioxide

Solvents	Reaction Temp. (°C)	SeO ₂ (mole)	Reaction Time (h)	<u>2</u>	<u>3</u>
Dioxane	60-65	1.5	1	31	60
Xylene	120-140	1.5	1	26	25
Ac ₂ O	120-140	1.5	1	0	0
Pyridine	110-116	1.5	1	3	5

Chart 2

Fig. 1. Perspective Drawing of 3

Crystal data: $\text{C}_{10}\text{H}_9\text{N}_3\text{OSe}$, $M=266.16$, orthorhombic, space group $Pbca$, $a=8.15(1)$, $b=9.402(1)$, $c=27.365(4)$ Å, $U=2099.0(5)$ Å³, $D_m=1.680(1)$, $D_g=1.684(1)$ g cm⁻³, $Z=8$. The structure was solved by direct methods (MULTAN 78⁶) and refined by least-squares to $R=0.088$ using 1787 independent reflections ($2\theta_{\text{max}} \leq 130^\circ$).

REFERENCES AND NOTES

- 1) N. Rabjohn, "Organic Reactions" Vol. V, ed. by R. Adams, John Wiley and Sons, Inc., London, 1949, p. 331; N. Rabjohn, "Organic Reactions" Vol. 24, ed. by W. G. Dauben, John Wiley and Sons, Inc., London, 1976, p. 261.
- 2) T. Sakamoto, T. Sakasai, and H. Yamanaka, *Chem. Pharm. Bull.*, 29, 2485 (1981).
- 3) S. Senda, K. Hirota, and O. Otani, *Yakugaku Zasshi*, 94, 571 (1974).
- 4) This compound was synthesized from 1 by the Eschweiler-Clark reaction.
- 5) T. Ueda, N. Oda, and I. Ito, *Heterocycles*, 8, 263 (1977).
- 6) P. Main, S. E. Hull, L. Lessinger, G. Germain, J. P. Declercq, and M. M. Woolfson, "A System of Computer Programs for the Automatic Solution of Crystal Structures from X-Ray Diffraction Data, MULTAN 78", University of York, 1978.

(Received July 15, 1982)