Communications to the Editor

[Chem. Pharm. Bull.]
 31(10)3769—3770(1983)

FORMATION OF PUMMERER'S KETONE FROM 4-METHYLPHENOXENIUM ION.
 AN ACID-CATALYZED SOLVOLYSIS OF N-p-TOLUENESULFONYL-O-(4-TOLYL)-
 HYDROXYLAMINE IN p-CRESOL

Yasuyuki Endo, Koichi Shudo* and Toshihiko Okamoto
 Faculty of Pharmaceutical Sciences, University of Tokyo, Hongo,
 Tokyo, 113, Japan

N-p-Toluenesulfonyl-O-(4-tolyl)hydroxylamine (1) reacts with
 p-cresol in the presence of trifluoroacetic acid to yield bisresol
 (2) and Pummerer's ketone (3).

KEYWORDS — phenoxenium ion; Pummerer's ketone; oxidation;
 oxidative coupling

Aryloxenium ions have received considerable interest in recent years particularly in connection with phenol oxidation¹⁾ and biosynthetic-type oxidative coupling reactions.²⁾ Previously, we have shown that aryloxenium ions can be generated in the acid-catalyzed solvolysis of N-acyl-O-arylhydroxylamines.³⁾ We now report an acid-catalyzed reaction of N-p-toluenesulfonyl-O-(4-tolyl)-hydroxylamine (1) with p-cresol: Pummerer's ketone is formed from 4-methylphenoxenium ion and cresol.

Reaction of (1) with p-cresol (25 equiv.) in the presence of trifluoroacetic acid (6.5 equiv.) at 20°C for 1 h gave 2,2'-dihydroxy-5,5'-dimethylbiphenyl (2) 24%, mp 152–153°C,⁴⁾ 2,9a-dimethyl-5a,6,7,9a-tetrahydro-7-oxodibenzofuran (Pummerer's ketone; (3), 9%, mp 124–125°C,⁵⁾ which was stable under the acidic conditions if the workup process is careful enough,⁶⁾ and 2-hydroxy-5-methylphenyl p-toluenesulfonimide (4)⁷⁾ (39%). Thus, bisresol (2) by ortho-ortho coupling and Pummerer's ketone (3) by ortho-para coupling was isolated. In this reaction, however, 2-hydroxy-4',5-dimethyldiphenylether was not produced by O-ortho coupling.

The mechanism for the reaction involves initial N-O bond heterolysis to the 4-methylphenoxenium ion (5), which attacks as an electrophilic species upon the aromatic nucleus of p-cresol.³⁾ We suggest that the reaction is a model for C-C bond formation in phenol oxidative coupling reactions. This suggestion is consistent with the Waters' view that PhO^+ participates in the C-C coupling reactions.^{1a)}

REFERENCES AND NOTES

- 1) (a) W.A.Waters, J.Chem.Soc.B, 1971, 2026; (b) K.Dimroth and H.Thomas, Chem. Ber. 100, 1967, 132; A.Ronlan and V.D.Parker, J.Chem.Soc.C, 1971, 3214; D.G.Hewitt, J.Chem.Soc.C, 1971, 1750; B.Speiser and A.Riekerm J.Chem.Res. [S], 1977, 314; J.Chem.Res. [M], 1977, 3601; A.Rieker, E.-L.Dreher, H.Geisel and M.H.Khalifa, Synthesis, 1978, 851.
- 2) D.H.R.Barton, Chem.Br., 3, 330 (1967); W.I.Taylor and A.R.Battersby, "Oxidative Coupling of Phenols," Marcel Dekker, New York, 1967; P.D.MacDonald and G.A.Hamilton in "Oxidation in Organic Chemistry," Part B, ed. by W.S.Trahanovsky, Academic Press, New York, 1973.
- 3) Y.Endo, K.Shudo and T.Okamoto, J.Am.Chem.Soc., 99, 7721 (1977); Y.Endo, K.Shudo and T.Okamoto, J.Am.Chem.Soc., 104, 6393 (1982); See also R.A.Abramovitch, M.Inbasekaran and S.Kato, J.Am.Chem.Soc., 95, 5428 (1973); R.A.Abramovitch, G.Alvernhe and M.Inbasekaran, Tetrahedron Lett., 1977, 1113; R.A.Abramovitch, G.Alvernhe, R.Bartnik, N.L.Dassanayake, N.Inbasekaran and S.Kato, J.Am.Chem. Soc., 103, 4558 (1981): for a generation of aryloxenium ions in the thermolysis of N-aryloxypyridinium salts.
- 4) C.G.Haynes, A.H.Turner and W.A.Waters, J.Chem.Soc., 1956, 2833.
- 5) K.M.Johnston, R.E.Jacobson and G.H.Williams, J.Chem.Soc.C, 1969, 1424.
- 6) The reaction mixture was poured into 5% aqueous sodium hydroxide at 0°C, and the solution was extracted with dichloromethane. The organic layer was dried and evaporated to give a sole product (3). The aqueous layer was acidified with concentrated hydrochloric acid and was extracted with dichloromethane. The dried dichloromethane extract was evaporated and p-cresol was removed under vacuum. The residue was chromatographed on a column of silica gel to give acidic products; (2) and (4).
- 7) The rearranged product (4) can be reasonably explained by acid-catalyzed [3,3]-sigmatropic-like rearrangement, which competes with the formation of the free aryloxenium ion. See Y.Endo, K.Shudo and T.Okamoto, Synthesis, 1980, 461.

(Received August 3, 1983)