[Chem. Pharm. Bull.]
31(6)1896—1901(1983)

Synthesis of 1,3-Dioxin-4-one Derivatives

MASAYUKI SATO, HIROMICHI OGASAWARA, KEIICHI OI and TETSUZO KATO*

Pharmaceutical Institute, Tohoku University Aobayama, Sendai 980, Japan

(Received November 15, 1982)

A facile and general synthesis of 2,2-dimethyl-1,3-dioxin-4-one derivatives (**1**) is reported. Treatment of β -keto acids with a mixture of acetone, acetic anhydride, and conc. sulfuric acid or with a mixture of isopropenyl acetate and conc. sulfuric acid gave 2,2-dimethyl-1,3-dioxin-4-ones (**1**). Similar treatment of *tert*-butyl esters of β -keto acids also gave **1**.

Keywords—2,2-dimethyl-1,3-dioxin-4-one derivative; diketene-acetone adduct; diketene; acylketene; β -keto acid; β -keto ester

2,2,6-Trimethyl-1,3-dioxin-4-one, so-called diketene-acetone adduct (**1a**), is easily prepared from diketene and acetone in the presence of an acidic catalyst.¹⁾ On heating, the adduct usually shows behavior similar to that of diketene itself.²⁾ For instance, the adduct reacts with compounds having a C=N or C $\equiv$ N bond to produce 1,3-oxazin-4-ones,³⁾ and with 1,3-dipoles such as isoquinolinium methylides to produce pyrrole derivatives.⁴⁾ The adduct also produces acetoacetyl derivatives on heating with alcohols, amines,¹⁾ and amides.⁵⁾

Such diketene-like reactivity of the adduct can be rationalized in terms of thermal fragmentation of the adduct to an acetylketene intermediate (**2a**).^{3a)} We recently obtained evidence that strongly supports the formation of the acetylketene intermediate (**2a**) in the reaction of the adduct.^{3c)}

Thus, 5- and 6-substituted 1,3-dioxin-4-ones, by analogy with adduct **1a**, may generate acylketenes **2**, which can be regarded as equivalent to mixed diketenes **3**. In the literature such mixed diketenes **3** are not easily accessible.⁶⁾ Though several references are available concerning synthesis of 1,3-dioxin-4-ones (**1**), most of the previous methods utilize 1,4-cycloaddition of ketones to acylketenes (**2**) prepared from acid halides,^{7a,b)} furanones,^{7c,d)} or diazo-ketones.^{7e)}

Chart 1

TABLE I. Synthesis of 2,2-Dimethyl-1,3-dioxin-4-ones (1b-j) from β -Keto Acids (5b-j)

Compd. No.	Substituent		Yield (%)	
	R ¹	R ²	Method A	Method B
1b	Me	Me	65	55
1c	Et	Me	68	77
1d	PhCH ₂	Me	68	63
1e	Ph	Me	89	74
1f	-(CH ₂) ₄ -		77	62
1g	Me	Ph	70	82
1h	H	Ph	59	74
1i	H	<i>p</i> -MeOPh	47	50
1j	H	<i>p</i> -NO ₂ Ph	48	53

We now report a facile and general synthesis of 1,3-dioxin-4-ones using β -keto acid derivatives.

Acetoacetic acids **5b–e**, 2-oxocyclohexanecarboxylic acid **5f**, and benzoylacetic acids **5g, h** were prepared in good yields by hydrolysis of the corresponding ethyl esters according to the literature. By applying Stiles's method,⁸⁾ *p*-anisoyl- and *p*-nitrobenzoylacetic acids (**5i, j**) were prepared by carboxylation of acetophenones with magnesium methyl carbonate (MMC).

Treatment of these acids with a mixture of acetone, acetic anhydride, and a catalytic amount of conc. sulfuric acid produced 1,3-dioxin-4-ones **1b–j** in good yields (method A). Treatment of the acids **5b–j** with a mixture of isopropenyl acetate and conc. sulfuric acid also produced 1,3-dioxin-4-ones **1b–j** (method B). The results are summarized in Table I.

When ketene was used in place of acetic anhydride or isopropenyl acetate, the acid **5h** gave **1h** in 62% yield (method C). Among these methods, method A seems to be most convenient in terms of the procedure, yield, and generality.

Use of *tert*-butyl esters of β -keto acids instead of ethyl esters eliminated the isolation of the acids, resulting in direct formation of 1,3-dioxin-4-ones. *tert*-Butyl 2-alkylacetoacetates **6b–d** were readily prepared from *tert*-butyl acetoacetate (**6a**). Treatment of *tert*-butyl esters **6b–d** with a mixture of acetone, acetic anhydride, and one equivalent of conc. sulfuric acid afforded the dioxinones **1b–d** in good yields. *tert*-Butyl 2,4-diphenylacetoacetate (**6k**) and 2-formylphenylacetate (**6l**) likewise cyclized to 1,3-dioxin-4-ones **1k, l**, respectively.

tert-Butyl benzoylacacetate (**6h**) has been synthesized from *tert*-butyl acetate in a low yield (25%).⁹⁾ We examined an alternative method based on the procedure given for ethyl benzoylacacetates.¹⁰⁾ Thus, *tert*-butyl acetoacetate (**6a**) was treated with benzoyl chloride in the presence of potassium *tert*-butoxide to give the diketo ester **7h** in a good yield. However, deacetylation of crude **7h** afforded **6h** in a low yield (27% from **6a**). *tert*-Butyl *p*-anisoylacacetate (**6i**) was

TABLE II. Physical, Analytical, and Spectral Data for 2,2-Dimethyl-1,3-dioxin 4-ones

Compd. No.	mp (°C) or bp (°C/mmHg)	Appearance (recrystn. solvent) ^{a)}	Formula	Analysis (%)		IR (CHCl ₃) cm ⁻¹	NMR (CDCl ₃) δ		
				Calcd (Found)	C H		2-Me	R ¹	R ²
1b	70/3	Oil	C ₈ H ₁₂ O ₃	61.52 (61.24)	7.75 (7.79)	1725 1622	1.60	1.75 (3H, s)	1.94 (3H, s)
1c	76/3	Oil	C ₉ H ₁₄ O ₃	63.51 (63.22)	8.29 (8.16)	1709 1642	1.62	1.03 (3H, t, <i>J</i> =7 Hz) 2.23 (2H, q, <i>J</i> =7 Hz)	1.95 (3H, s)
1d	43–44	Prisms (A)	C ₁₄ H ₁₆ O ₃	72.39 (72.09)	6.94 (7.11)	1712 1640	1.63	3.63 (2H, s) 7.22 (5H, s)	1.99 (3H, s)
1e	66–67	Prisms (B)	C ₁₃ H ₁₄ O ₃	71.54 (71.26)	6.47 (6.32)	1718 1630	1.76	7.30 (5H, s)	1.92 (3H, s)
1f	36–37	Prisms (C)	C ₁₀ H ₁₄ O ₃	65.91 (65.98)	7.74 (7.82)	1716 1650	1.66	1.65 (4H, m), 2.22 (4H, m)	
1g	86–87	Prisms (A)	C ₁₃ H ₁₄ O ₃	71.54 (71.74)	6.47 (6.56)	1710 1630	1.78	2.00 (3H, s)	7.48 (5H, s)
1h	62–63	Needles (B)	C ₁₂ H ₁₂ O ₃	70.57 (70.47)	5.79 (5.79)	1717 1622	1.80	5.88 (1H, s)	7.85 (5H, s)
1i	75–76	Leaves (D)	C ₁₃ H ₁₄ O ₄	66.65 (66.45)	6.02 (5.86)	1710 1610	1.79	5.79 (1H, s)	3.87 (3H, s) 6.83 (2H, d, <i>J</i> =13 Hz) 7.66 (2H, d, <i>J</i> =13 Hz) 7.88 (2H, d, <i>J</i> =13 Hz) 8.36 (2H, d, <i>J</i> =13 Hz)
1j	142 (dec.)	Needles (E)	C ₁₂ H ₁₁ NO ₅	57.83 (57.83)	4.45 ^{b)} (4.27)	1720 1620	1.85	6.04 (1H, s)	3.48 (2H, s) 7.25 (5H, m) 7.17 (1H, s)
1k	82–83	Needles (C)	C ₁₉ H ₁₈ O ₃	77.53 (77.78)	6.16 (6.10)	1720 1630	1.62	7.15 (5H, s)	
1l	38–39	Prisms (C)	C ₁₂ H ₁₂ O ₃	70.57 (70.32)	5.92 (5.96)	1722 1622	1.70	7.30 (5H, m)	
1m	65/1	Oil	C ₆ H ₈ O ₃ ^{c)}			1725 1622	1.70	5.30 (1H, d, <i>J</i> =9 Hz)	7.07 (1H, d, <i>J</i> =9 Hz)

a) A, hexane; B, hexane-ether; C, pentane; D, ether; E, ethyl acetate.

b) Calcd: N, 5.62; Found: N, 5.43.

c) Determined by high resolution mass spectrometry. Found: 128.0483; Calcd for C₆H₈O₃ (M⁺): 128.0473.

similarly prepared in 37% yield. Cyclization of the esters **6h** and **6i** gave 1,3-dioxin-4-ones **1h** and **1i** in 52 and 40% yields, respectively.

The cyclization of *tert*-butyl esters presumably proceeds through the keto acid **5**, since *tert*-butyl esters are easily converted to free acids under acidic conditions. Ethyl esters such as ethyl acetoacetate and benzoylacetate did not cyclize directly to 1,3-dioxin-4-ones under the same conditions.

Iwataki *et al.* have prepared compound **1a** and the 6-ethyl analogue (**1**: $R^1=H$, $R^2=Et$) by heating of acetyl Meldrum's acid (**8a**: $R^1=H$, $R^2=Me$) and the propionyl analogue (**8**: $R^1=H$, $R^2=Et$).¹¹⁾ Applying this procedure, the formyl compound **8m** ($R^1=R^2=H$) was heated without solvent to give a resinous product. However, heating of compound **8m** in a mixture of toluene and acetone gave 2,2-dimethyl-1,3-dioxin-4-one (**1m**) in 31% yield. Acetylation of methyl Meldrum's acid afforded 5-acetyl-2,2,5-trimethyl-1,3-dioxane-4,6-dione (**8b**: $R^1=R^2=Me$). This compound is relatively stable, and the thermal conversion had to be conducted in a sealed tube at 185°C to give **1b** in low yield.

The physical, analytical, and spectral data for 1,3-dioxin-4-ones **1b—m** are summarized in Table II. It should be mentioned that the mass spectra of all dioxinones showed the fragment ion peak ($M-58$)⁺, which corresponds to the ionized acylketene. This fragmentation resembles the thermal fragmentation of **1a**.

The results of this investigation indicate that the cyclization of β -keto acid derivatives with acetone is a very useful method for preparing a variety of 2,2-dimethyl-1,3-dioxin-4-ones. Thermal reactions of compounds **1b—m** with 1,2-, 1,3-, and 1,4-dipoles are under investigation.

Experimental

All melting and boiling points are uncorrected. Infrared (IR) spectra were measured with a JASCO A-102 spectrometer. Nuclear magnetic resonance (NMR) spectra were measured on a JEOL JNM-PMX 60 spectrometer and are reported as δ values (ppm) relative to tetramethylsilane as an internal standard. Mass spectra (MS) were obtained on a Hitachi M-52G mass spectrometer.

General Procedure for Preparation of β -Keto Acids **5b—h**—Compounds **5b**,¹²⁾ **5c**,¹³⁾ **5d**,¹²⁾ **5f**,¹⁴⁾ **5g**,¹⁵⁾ and **5h**¹⁶⁾ were prepared from the ethyl esters by slight modifications of the reported procedures. A mixture of a β -keto ester (0.1 mol) and 0.5 N aq. sodium hydroxide (200 ml) was stirred under ice-cooling for 3 h, and then left to stand at room temperature for 12 h. The reaction mixture was washed with ether and acidified with conc. hydrochloric acid under ice-cooling.

The precipitated acids (**5f—h**) were collected by suction, washed with water and then with carbon tetrachloride, and dried. The oily acids (**5b—d**) were extracted with ether from the acidified solution. The ether solution was washed with brine and dried over anhydrous magnesium sulfate. Removal of the ether *in vacuo* at room temperature gave the acids **5b—d** as oily substances.¹⁷⁾ 2-Phenylacetoacetic acid (**5e**) was prepared according to the literature.¹⁸⁾

***p*-Anisoylacetic Acid (**5i**)**—*p*-Methoxyacetophenone (3.0 g, 20 mmol) was heated with a 2 M solution of magnesium methyl carbonate⁹⁾ in dimethylformamide (40 ml, 80 mmol) at 110–120°C for 1 h. The reaction mixture was poured onto crushed ice, and acidified with conc. hydrochloric acid under ice-cooling. The solution was cooled at –10°C for 1 h. The separated crystals were collected by suction, washed with water and dissolved in 5% sodium carbonate. The solution was filtered and the filtrate was acidified with dil. hydrochloric acid. Separated crystals were collected by suction, washed with water, and dried to give **5i** as fine needles of mp 85–86°C (dec.) (lit.¹⁹⁾ mp 80°C (dec.)). Yield, 1.67 g (43%).

***p*-Nitrobenzoylacetic Acid (**5j**)**—*p*-Nitroacetophenone (3.30 g, 20 mmol) was treated in the same manner as described above, giving **5j** as fine prisms of mp 130°C (dec.) (lit.²⁰⁾ mp 135°C (dec.)). Yield, 2.2 g (53%).

General Procedure for Preparation of 2,2-Dimethyl-1,3-dioxin-4-ones (1b—j**) from Acids **5b—j****. **Method A**—Conc. sulfuric acid (0.01 mol) was added dropwise to a mixture of one of **5b—j** (0.05 mol), acetone (0.1 mol), and acetic anhydride (0.1 mol) with stirring below 5°C. The mixture was stirred under ice-cooling for 3 h, during which time crystalline **5** dissolved. After being kept in a refrigerator (*ca.* 0°C) for 12 h, the mixture was poured into 10% sodium carbonate solution (120 ml) under ice-cooling. The mixture was stirred at room temperature for 30 min to give the corresponding product.

Separated crystals were collected by suction, washed with water, dried, and recrystallized to give **1d—h**. Oily products **1b, c** were extracted with ether. The ether solution was washed with water, dried over magnesium sulfate and concentrated. Distillation under reduced pressure gave **1b, c**.

Method B—Conc. sulfuric acid (0.01 mol) was added dropwise to a mixture of **5** (0.05 mol) and isopropenyl acetate (0.1 mol) with stirring below 5°C. The mixture was stirred under ice-cooling for 3 h and then kept in a refrigerator for 12 h. The reaction mixture was worked up in the same manner as in method A, giving the corresponding **1**. Yields are shown in Table I. Physical, analytical, and spectral data are shown in Table II.

Method C—Conc. sulfuric acid (380 mg, 4 mmol) was added to a mixture of benzoylacetic acid (**5h**) (1.64 g, 10 mmol) and acetone (2.32 g, 40 mmol) under ice-cooling. Ketene (40 mmol) was introduced into the solution with stirring under ice-cooling. The mixture was kept in a refrigerator for 12 h, then worked up in the same manner as in method A, giving **1h**. Yield, 1.26 g (62%).

tert-Butyl 2,4-Diphenylacetoacetate (6k)—A mixture of *tert*-butyl phenylacetate²¹ (3.84 g) and sodium hydride (460 mg) was heated at 90°C with stirring for 2 h. The mixture was neutralized with dil. acetic acid, and precipitated crystals were collected by suction, washed with water, and dried. Recrystallization from pentane gave needles of mp 81–82°C. Yield, 1.9 g (61%). *Anal.* Calcd for C₂₀H₂₂O₃: C, 77.39; H, 7.14. Found: C, 77.12; H, 7.18. IR (CHCl₃): 1735, 1710 cm⁻¹. NMR (CCl₄) δ : 1.40 (9H, s), 3.62 (2H, s), 4.53 (1H, s), 6.90–7.40 (10H, m) (keto form); 1.30 (9H, s), 3.34 (2H, s), 6.90–7.40 (10H, m), 13.15 (1H, s) (enol form) (keto : enol = 1 : 1).

tert-Butyl 2-Formylphenylacetate (6l)—Ethyl formate (0.81 g) was added dropwise to a stirred and ice-cooled mixture of *tert*-butyl phenylacetate (1.92 g), potassium *tert*-butoxide (2.24 g), and dry ether (30 ml). After being stirred for 1 h under ice-cooling, the reaction mixture was neutralized with dil. acetic acid and extracted with ether. The ether solution was washed with water and dried over magnesium sulfate. Removal of the solvent *in vacuo* gave an oily substance, which was subjected to silica gel column chromatography. Elution with hexane–ethyl acetate (20 : 1) gave **6l** as prisms (from pentane) of mp 55–56°C. Yield, 1.02 g (46%). *Anal.* Calcd for C₁₃H₁₆O₃: C, 70.89; H, 7.32. Found: C, 70.82; H, 7.26. IR (CHCl₃): 1650 cm⁻¹. NMR (CCl₄) δ : 1.50 (9H, s), 7.17 (5H, s), 7.17 (1H, d, *J* = 13 Hz), 12.14 (1H, d, *J* = 13 Hz) (enol form).

tert-Butyl Benzoylacacetate (6h)—*tert*-Butyl acetoacetate (**6a**, 6.32 g) was added dropwise to a mixture of potassium *tert*-butoxide (9.88 g) and dry ether with stirring under ice-cooling. The mixture was stirred at room temperature for 30 min. Benzoyl chloride (6.18 g) was added dropwise to the solution at ca. 0°C, and stirring was continued at room temperature for 3 h. The mixture was acidified with 10% hydrochloric acid under ice-cooling and extracted with ether. The ether layer was washed with water, dried over magnesium sulfate, and concentrated to give crude *tert*-butyl 2-benzoylacetoacetate (**7h**) as an oil. Yield, 12 g. Crude **7h** (12 g) was stirred vigorously with a mixture of ammonium chloride (10.24 g), 28% ammonia (9.6 ml), and water (48 ml) at 42°C for 20 min. The mixture was extracted with ether. The ether layer was washed with water, dried over magnesium sulfate and concentrated. Distillation of the residue gave **6h** as an oil of bp 89–93°C (0.1 mmHg) (lit.⁹) bp 95°C (0.2 mmHg). Yield, 5.41 g (27% from compound **6a**).

tert-Butyl *p*-Anisoylacacetate (6i)—Following the procedure given for **6h**, compound **6a** (6.32 g) was treated with *p*-anisoyl chloride (7.48 g), giving crude **7i** (12 g) as an oil. Following the procedure for the preparation of **6h**, crude **7i** was deacetylated to give **6i** (6.4 g) as an oil. Purification by silica gel column chromatography using hexane–ethyl acetate (4 : 1) as an eluent gave **6i** as an oil. Yield, 4.0 g (37% from **6a**). IR (CHCl₃): 1730, 1680, 1600 cm⁻¹. NMR (CCl₄) δ : 1.20 (9H, s), 3.70 (2H, s), 3.83 (3H, s), 6.85 (2H, d, *J* = 13 Hz), 7.83 (2H, d, *J* = 13 Hz). Cu salt (recrystallized from ethanol): green needles of mp 205–208°C (dec.). *Anal.* Calcd for C₂₈H₃₆CuO₈: C, 59.61; H, 6.43. Found: C, 59.47; H, 6.14.

General Procedure for Preparation of Compound 1 from Compound 6—Conc. sulfuric acid (0.03 mol) was added dropwise to a mixture of **6** (0.03 mol), acetone (0.06 mol), and acetic anhydride (0.09 mol) with stirring below –5°C. The mixture was stirred under ice-cooling for 3 h, and then kept in a refrigerator for 12 h. The reaction mixture was poured into 10% sodium carbonate (120 ml) under ice-cooling. The mixture was worked up as described for method A to give **1b** (71%), **1c** (80%), **1d** (60%), **1h** (52%), **1i** (40%), **1k** (65%), or **1l** (90%). Physical, analytical, and spectral data are listed in Table II.

2,2-Dimethyl-1,3-dioxin-4-one (1m)—A mixture of **8m**²² (2.58 g), acetone (1.74 g) and xylene (60 ml) was heated under reflux for 30 min. The reaction mixture was subjected to silica gel column chromatography. Elution with hexane–ether (8 : 1) gave **1m** as an oil. Physical, analytical, and spectral data are listed in Table II.

5-Acetyl-2,2,5-trimethyl-1,3-dioxane-4,6-dione (8b)—Acetyl chloride (1.73 g) was added dropwise to a mixture of 2,2,5-trimethyl-1,3-dioxane-4,6-dione²³ (3.16 g), pyridine (3.48 g), and dichloromethane (50 ml) with stirring at –10––5°C. After being stirred for 1 h at room temperature, the mixture was washed with aq. 5% hydrochloric acid, and then with water. The organic layer was dried over magnesium sulfate and concentrated to dryness *in vacuo*. The residue was recrystallized from ether giving **8b** as prisms of mp 67–69°C. Yield, 3.3 g (83%). *Anal.* Calcd for C₉H₁₂O₅: C, 53.99; H, 6.04. Found: C, 53.73; H, 6.16. IR (CHCl₃): 1790, 1750 cm⁻¹. NMR (CDCl₃) δ : 1.80 (6H, s), 1.90 (3H, s), 2.33 (3H, s).

Conversion of Compound 8b to Compound 1b—Compound **8b** (2 g) and acetone (0.87 g) were heated in a sealed tube at 185°C for 1 h. The reaction mixture was distilled under reduced pressure to give 0.5 g (32%) of **1b**.

Acknowledgement The present work was supported in part by a Grant-in-Aid for Scientific Research

from the Ministry of Education, Science and Culture, Japan, which is gratefully acknowledged. Thanks are due to Mrs. C. Koyanagi for elemental analyses, and to Miss E. Kurosawa for mass spectral measurements.

References and Notes

- 1) M.F. Carroll and A.R. Bader, *J. Am. Chem. Soc.*, **75**, 5400 (1953); E.V. Dehmlow and A.R. Shamout, *Justus Liebigs Ann. Chem.*, **1982**, 1753.
- 2) For a review of the reactions of diketene, see: T. Kato, *Acc. Chem. Res.*, **7**, 267 (1974).
- 3) a) G. Jäger and J. Wenzelburger, *Justus Liebigs Ann. Chem.*, **1976**, 1689; b) M. Sato, H. Ogasawara, E. Yoshizumi, and T. Kato, *Heterocycles*, **17**, 297 (1982); c) *Idem*, *Chem. Pharm. Bull.*, **31**, 1902 (1983).
- 4) M. Sato, N. Kanuma, and T. Kato, *Chem. Pharm. Bull.*, **30**, 4359 (1982).
- 5) M. Sato, N. Kanuma, and T. Kato, *Chem. Pharm. Bull.*, **30**, 1315 (1982).
- 6) H.W. Moore and D.S. Wilbur, *J. Org. Chem.*, **45**, 4483 (1980).
- 7) a) G. Jäger, *Chem. Ber.*, **105**, 137 (1972); b) G. Jäger and R.J. Singer, *Justus Liebigs Ann. Chem.*, **1977**, 1799; c) G. Kollenz, E. Ziegler, W. Ott, and G. Kriwetz, *Z. Naturforsch., B: Anorg. Chem., Org. Chem.*, **32**, 701 (1977) [*Chem. Abstr.*, **87**, 201432n (1977)]; d) Y.S. Andreichikov, L.F. Gein, and G.D. Plakhina, *Zh. Org. Khim.*, **16**, 2336 (1980) [*Chem. Abstr.*, **94**, 156840a (1981)]; e) H. Stetter and K. Kiehs, *Chem. Ber.*, **98**, 2099 (1965).
- 8) M. Stiles, *J. Am. Chem. Soc.*, **81**, 2598 (1959).
- 9) A. Treibs and K. Hintermeier, *Chem. Ber.*, **87**, 1163 (1954).
- 10) R.L. Shriner, A.G. Schmidt, and L.J. Roll, "Organic Syntheses," Coll. Vol. 2, ed. by A.H. Blatt, John Wiley and Sons, Inc., New York, 1943, p. 266.
- 11) I. Iwataki, S. Makisawa, and K. Saito, Jpn. Patent 79106478 (1979) [*Chem. Abstr.*, **92**, 41966n (1980)].
- 12) M. Ceresole, *Chem. Ber.*, **15**, 1871 (1882).
- 13) C. Mannich and M. Bauroth, *Chem. Ber.*, **57**, 1108 (1924).
- 14) H.D. Gardner, W.H. Perkin, and H. Watson, *J. Chem. Soc.*, **97**, 1764 (1910).
- 15) E. Hope and W.H. Perkin, *J. Chem. Soc.*, **95**, 2045 (1909).
- 16) E. Beckmann and T.V. Paul, *Justus Liebigs Ann. Chem.*, **1891**, 16.
- 17) Compound **5d**, reported to be an oil, crystallized in a refrigerator; mp 47–50°C (dec.).
- 18) A.J. Attwood, A. Stevenson, and J.F. Thorpe, *J. Chem. Soc.*, **1923**, 1762.
- 19) C.G. Swain, R.F. Bader, R.M. Esteve, and R.N. Griffin, *J. Am. Chem. Soc.*, **83**, 1951 (1961).
- 20) C. Bulow and E. Hailer, *Chem. Ber.*, **35**, 915 (1902).
- 21) W.G. Kenyon, R.B. Meyer, and C.R. Hauser, *J. Org. Chem.*, **28**, 3108 (1963).
- 22) G.A. Bihlmayer, G. Derflinger, J. Derkosch, and O.E. Polansky, *Monatsh. Chem.*, **98**, 564 (1967).
- 23) K. Pihlaja and M. Seilo, *Acta Chem. Scand.*, **22**, 3053 (1968).