[Chem. Pharm. Bull.]
32(4)1624—1627(1984)

Ethanol Enhancement of Cytochrome P-450 Content in Yeast, *Saccharomyces cerevisiae* D7

TAMOTSU MORITA* and ICHII MIFUCHI

Shizuoka College of Pharmacy, 2-2-1 Oshika, Shizuoka 422, Japan

(Received August 8, 1983)

When *Saccharomyces cerevisiae* D7 was cultured in yeast extract-proteose peptone-2% glucose (YPG) liquid medium supplemented with 1.5 and 3% ethanol, the maximum cytochrome P-450 contents in the whole cell suspension were 1.5 and 1.3 times higher, respectively, than that in YPG medium without ethanol. The addition of 6% ethanol to YPG medium decreased the growth rate and final cell population of the yeast markedly, while the cellular content of cytochrome P-450 was similar to that of cells grown in YPG medium. Though the logarithmic phase was extended under glucose-repressed conditions in YPG medium containing 10% glucose, there was no increase in the cytochrome P-450 content compared with the cells grown on 2% glucose. In the medium in which 2% ethanol was the sole carbon and energy source (YPE medium), the growth rate and final cell population of the yeast showed a marked decrease compared with those of cells grown on glucose. Cytochrome P-450 was not detectable in the cells grown on ethanol.

Keywords—cytochrome P-450 content; ethanol enhancement; reduced CO-difference spectrum; yeast; *Saccharomyces cerevisiae*

Saccharomyces cerevisiae has a microsomal cytochrome P-450¹⁾ which is in many respects similar to those found in mammalian microsomes.²⁾ Though little is known about the metabolic role of yeast cytochrome P-450, yeast has been used as a screening organism for promutagens and procarcinogens without mammalian activation systems such as S-9 fraction and host mediated assay.³⁾

Yeast cytochrome P-450 is formed in the cells grown under various conditions, such as in respiratory-repressed cells grown on high glucose concentration (up to 20%),^{2a,4)} in semi-anaerobically grown cells^{2a,5)} and in aerobically grown cells on relatively low concentration of glucose.^{3a,6,7)} In every case, the cytochrome P-450 content in batch-cultured yeast cells was growth-related and reached maximum at the end of the logarithmic phase of growth. Therefore, it was suggested that the formation of cytochrome P-450 in yeast cells is affected by a common metabolite of glucose or an effector generated by the metabolite.^{6a)} Ethanol is a common metabolite in cultures of yeast grown on glucose.⁸⁾ It is also used as a solvent and/or a disinfectant of the compounds tested in mutagenicity assays. The aim of the present paper was therefore to study the effect of exogenously added ethanol on cytochrome P-450 formation in *S. cerevisiae* D7, which was constructed for mutagenicity assay.⁹⁾

Experimental

Yeast Strain and Culture Conditions—*Saccharomyces cerevisiae* D7 (*a/x*, *ade2-40/ade2-119*, *trp5-12/trp5-27*, *ilv1-92/ilv1-92*)⁹⁾ was obtained from Dr. F. K. Zimmermann. Preincubated yeast cells were inoculated into 4 l of each liquid medium at 5×10^4 cells/ml and incubated with vigorous aeration (1.2 l/min) at 30°C. After various time intervals, samples were withdrawn and cell population and cytochrome P-450 contents were determined. Cell population was counted with a hemocytometer after supersonication (40 W, 29 kHz, 8 s) to remove cell clumps and unseparated daughter cells.

Media—Standard YPG medium contained 1% Difco yeast extract, 2% Difco proteose peptone No. 3 and 2% glucose. YPE medium contained 2% ethanol as a carbon and energy source instead of glucose in YPG medium. YPG

medium was also supplemented with 0.5, 1.5, 3 and 6% ethanol. Glucose was sterilized separately and ethanol was added after cooling the autoclaved medium.

Measurement of Cytochrome P-450 Content in Whole Cell Suspensions of Yeast—Cytochrome P-450 was measured principally as described by Omura and Sato.¹⁰⁾ Harvested yeast cells were washed twice in cold distilled water and suspended in 0.1 M phosphate buffer (pH 7.2) at 1.0×10^9 cells/ml. About 0.3 mg/ml of sodium dithionite ($\text{Na}_2\text{S}_2\text{O}_4$) was added and CO gas was bubbled through the sample cuvette for 30 s. The reduced CO-difference spectra were recorded with a Hitachi EPS-3T spectrophotometer. The approximate contents of cytochrome P-450 were calculated as described by Kowal *et al.*¹¹⁾ by multiplying the absorption differences between 460 and 490 nm by a factor of 2.3 (see Results) and expressed as nmol of cytochrome P-450/ 10^9 cells. An extinction coefficient of 92/mm/cm¹²⁾ was used.

Results

Measurement of Cytochrome P-450 Content and Contribution of Cytochrome Oxidase

In the quantitation of cytochrome P-450 in the whole cell suspensions of *S. cerevisiae* D7, the absorption peak of the cytochrome shifted from 450 to 460 nm after the end of the logarithmic phase. This shift mainly resulted from a deep trough around 443 nm due to the reduced-CO spectrum of cytochrome oxidase (cytochrome a_3). Thus, it was not possible to quantitate the cytochrome P-450 content in the whole cell suspensions on the basis of the absorbancy at 450 nm. We eliminated the contribution of cytochrome oxidase by the method of Kowal *et al.*¹¹⁾ in which the absorption change between 460 and 490 nm ($A_{460-490}$) was used for calculating the cytochrome P-450 content in bovine liver microsomes.

Table I shows the ratio of $A_{460-490}$ to $A_{450-490}$ in the reduced CO-difference spectra of whole cell suspension and microsome fraction of *S. cerevisiae* D7 harvested before the late logarithmic phase, when little cytochrome oxidase had been formed. The ratio was essentially constant at 43.3 ± 1.9 ($n = 13$). The whole cell suspension from stationary phase culture, which contained a high level of cytochrome oxidase, did not show any absorption at 460 nm (data were not shown). Thus, the approximate contents of cytochrome P-450 in whole cell suspensions were calculated by multiplying $A_{460-490}$ by the factor of 2.3 throughout the

TABLE I. The Ratio of $A_{460-490}$ to $A_{450-490}$ in the Reduced CO-Difference Spectra of Whole Cell Suspension and Microsome Fraction of *S. cerevisiae* D7

Medium	Incubation time (h)	Concentration	$\frac{A_{460-490}}{A_{450-490}}$ (%)
YPG	14	0.5×10^9 cells	45.5
	14	1.0×10^9 cells	44.8
	14	1.0×10^9 cells	44.8
	14	1.7×10^9 cells	42.7
	16	1.0×10^9 cells	45.2
	16	1.0×10^9 cells	44.9
	14	33 mg protein	42.2
	14	55 mg protein	45.1
YPG with 1.5% ethanol	12	1.0×10^9 cells	40.0
	14	1.0×10^9 cells	42.2
	16	1.0×10^9 cells	41.1
YPG with 3% ethanol	14	1.0×10^9 cells	44.3
	16	1.0×10^9 cells	40.0
Average $\pm$ SD			43.3 ± 1.9

The percent ratios of $A_{460-490}$ to $A_{450-490}$ in the reduced CO-difference spectra are shown. Concentrations are given as cells/ml for whole cell suspension and as mg protein/ml for microsome fraction. Other conditions were the same as in Figs. 1 and 2.

Fig. 1. Growth Curve of *S. cerevisiae* D7 in Each Medium

Media used: yeast extract-peptone-2% glucose (YPG, ●), YPG with 1.5% ethanol (○); with 3% ethanol (▲) and with 6% ethanol (△); YPG with 10% glucose (□); yeast extract-peptone-2% ethanol (YPE, ■). Yeast cells were inoculated into each medium at 5×10^4 cells/ml and incubated at 30 °C aerobically.

Fig. 2. Time Course of Change of Cellular Cytochrome P-450 Content in *S. cerevisiae* D7 in Each Medium

Media used: Same as in Fig. 1. Samples were withdrawn from each medium, washed and suspended in phosphate buffer at 1.0×10^9 cells/ml. Cytochrome P-450 content was calculated from the reduced CO-difference spectrum as described in the text. Results for YPG and YPG with 1.5% ethanol are averages $\pm$ S.D. of 4 and 2 independent experiments, respectively.

experiment.

The Effect of Ethanol on Growth and Cellular Cytochrome P-450 Content of Yeast

Figures 1 and 2 show the time course of growth and cytochrome P-450 content of *S. cerevisiae* D7 in each medium. The addition of 0.5% ethanol to YPG medium did not affect the growth or cytochrome P-450 formation of the yeast (data were not shown). Though the growth and the final cell population were almost the same in YPG medium (control) and that containing 1.5% ethanol, the maximum cellular content of cytochrome P-450 was about 50% higher in the latter. In YPG medium containing 3% ethanol, the growth was somewhat repressed but the maximum cytochrome P-450 content was still 33% higher than the control. Though the growth and final cell population decreased markedly in the presence of 6% ethanol, the cytochrome P-450 content was still equal to that of the control.

In YPG medium containing 10% glucose, the logarithmic phase was extended and the final cell population was almost 2.6 times higher than that in 2% glucose-YPG medium. Though the time course of change of cytochrome P-450 content was rather gradual in 10% glucose-YPG medium, there was no decrease in the maximum content of cellular cytochrome P-450 (Fig. 2). When the yeast was grown on 2% ethanol as a sole carbon and energy source, the growth rate and the final cell population decreased markedly. There was no detectable cytochrome P-450 in the whole cell suspension from the culture grown on ethanol.

Discussion

It has been reported that in aerobically grown yeast cells, cytochrome P-450 is formed only in a medium with high glucose concentration,^{2a,4)} but in the present experiment, the

cytochrome P-450 contents were the same in *S. cerevisiae* D7 grown on 2 and 10% glucose. Callen *et al.*^{3a)} and Wiseman *et al.*⁷⁾ have also shown that almost the same amount of cytochrome P-450 is formed in yeast cells grown on 2 or 10% glucose and on 1 or 20% glucose, respectively.

No detectable cytochrome P-450 was formed in *S. cerevisiae* D7 grown on ethanol as a sole carbon and energy source. Cytochrome P-450 was also not formed in other yeast genera, *Candida* and *Torulopsis*, during their growth on ethanol.¹³⁾ In contrast, ethanol increased the cytochrome P-450 contents in higher plant tissues,¹⁴⁾ cultured hepatocytes,¹⁵⁾ rats¹⁶⁾ and hamsters.¹⁷⁾ The present experiment showed that the addition of 1.5 or 3% ethanol to the medium could increase the cytochrome P-450 content in yeast cells grown on glucose. Though the mechanism of this increase in cytochrome P-450 content is not clear from the present data, it is suggested that ethanol may not be able to induce cytochrome P-450 but may enhance the formation or inhibit the degradation of cytochrome P-450 in the yeast cells grown on glucose. Detailed studies on the formation of cytochrome P-450 and on the activation of promutagens and/or procarcinogens should facilitate the application of the yeast as a screening organism.

Acknowledgement The authors are grateful to Dr. F. K. Zimmermann of Technische Hochschule, Darmstadt, for the supply of *S. cerevisiae* D7.

References

- 1) A. Lindenmayer and L. Smith, *Biochim. Biophys. Acta*, **93**, 445 (1964).
- 2) a) K. Ishidate, K. Kawaguchi, K. Tagawa and B. Hagihara, *J. Biochem. (Tokyo)*, **65**, 375 (1969); b) Y. Yoshida and H. Kumaoka, *J. Biochem. (Tokyo)*, **71**, 915 (1972); c) Y. Yoshida and H. Kumaoka, *J. Biochem. (Tokyo)*, **78**, 785 (1975); d) A. Wiseman and T.-K. Lim, *Biochem. Soc. Trans.*, **3**, 974 (1975); e) L. F. J. Woods and A. Wiseman, *Biochim. Biophys. Acta*, **613**, 52 (1980).
- 3) a) D. F. Callen and R. M. Philpot, *Mutat. Res.*, **45**, 309 (1977); b) D. Siebert, U. Bayer and H. Marquardt, *Mutat. Res.*, **67**, 145 (1979); c) V. F. Simmon, *J. Natl. Cancer Inst.*, **62**, 901 (1979); d) V. W. Mayer and C. J. Goin, *Mutat. Res.*, **78**, 243 (1980); e) P. Wilcox and J. M. Parry, *Carcinogenesis*, **2**, 1201 (1981); f) E. R. Nestmann, E. R. Stephen, D. J. Kowbel and A. Nasim, *Can. J. Genet. Cytol.*, **24**, 771 (1982); g) R. D. Mehta and R. C. von Borstel, *Mutat. Res.*, **92**, 49 (1982).
- 4) A. Wiseman, C. McCloud and T.-K. Lim, *Biochem. Soc. Trans.*, **4**, 685 (1976).
- 5) a) Y. Yoshida, H. Kumaoka and R. Sato, *J. Biochem. (Tokyo)*, **75**, 1201 (1974); b) Y. Yoshida, H. Kumaoka and R. Sato, *J. Biochem. (Tokyo)*, **75**, 1211 (1974); c) M. Trinn, O. Käppeli and A. Fiechter, *Eur. J. Appl. Microbiol. Biotech.*, **15**, 64 (1982).
- 6) a) S. O. Kärenlampi, E. Marin and O. O. P. Hänninen, *Biochem. J.*, **194**, 407 (1981); b) Y. Aoyama, T. Okikawa and Y. Yoshida, *Biochim. Biophys. Acta*, **665**, 596 (1981).
- 7) A. Wiseman, T.-K. Lim and C. McCloud, *Biochem. Soc. Trans.*, **3**, 276 (1975).
- 8) E. S. Polakis, W. Bartley and G. A. Meek, *Biochem. J.*, **90**, 369 (1964).
- 9) F. K. Zimmermann, R. Kern and H. Rasenberger, *Mutat. Res.*, **28**, 381 (1975).
- 10) T. Omura and R. Sato, *J. Biol. Chem.*, **239**, 2370 (1964).
- 11) J. Kowal, E. R. Simpson and R. W. Estabrook, *J. Biol. Chem.*, **245**, 2438 (1970).
- 12) Y. Yoshida, Y. Aoyama, H. Kumaoka and S. Kubota, *Biochem. Biophys. Res. Commun.*, **78**, 1005 (1977).
- 13) A. P. Itchenko, S. Mauersberger, R. N. Matyashova and A. B. Lozinov, *Mykrobiologiya*, **49**, 452 (1980).
- 14) D. Reichhart, J. P. Salaün, I. Benveniste and F. Durst, *Arch. Biochem. Biophys.*, **196**, 301 (1979).
- 15) J. F. Sinclair, P. R. Sinclair, E. L. Smith, W. J. Bement, J. Pomeroy and H. Bonkowsky, *Biochem. Pharmacol.*, **30**, 2805 (1981).
- 16) a) E. Rubin, C. S. Lieber, A. P. Alvarez, W. Levin and R. Kuntzman, *Biochem. Pharmacol.*, **20**, 229 (1971); b) J.-P. Villeneuve, P. Mavie and J.-G. Joly, *Biochem. Biophys. Res. Commun.*, **70**, 723 (1976).
- 17) G. D. McCoy, C. B. Chen, S. S. Hecht and E. C. McCoy, *Cancer Res.*, **39**, 793 (1979).