[Chem. Pharm. Bull.
33(11)5108—5109(1985)]

A Convenient Procedure for the Synthesis of Phenothiazine Drugs

HERVÉ GALONS,^a MARCEL MIOCQUE,^{*,a} CLAUDE COMBET-FARNOUX,^a
YOUNES BENSÂÏD,^a GUY DECODTS,^b
and GEORGES BRAM^b

*Laboratoire de Chimie Organique, Faculté de Pharmacie, U.A. CNRS 496,^a Avenue J.B. Clément,
92290 Châtenay-Malabry, France and Laboratoire de Chimie des Réactions
Sélectives sur Support, U.A. CNRS 478, Université Paris-Sud,^b
Bâtiment 410, 91405 Orsay, France*

(Received March 8, 1985)

Alkylation of phenothiazines to prepare chlorpromazine and analogous compounds was achieved by solid-liquid phase-transfer catalysis in the presence of Aliquat 336 in the absence of any organic solvent. This simple procedure is easy and rapid.

Keywords—phenothiazine; chlorpromazine; alkylation; phase-transfer catalysis

The study of the biological activity of phenothiazine derivatives led to the discovery of the first neuroleptic agents and to the development of major antihistaminic and anti-parkinsonian drugs. These compounds are in most cases prepared by alkylation of the parent heterocycle. Since the proton on the nitrogen is only weakly acidic, the anionic activation requires strong bases such as alkali metal amides¹⁻³⁾ or organometallic compounds.^{4,5)}

Very recently, phase-transfer catalysis was used successfully in the preparation of chlorpromazine⁶⁾ and alkylphenothiazines.⁷⁾ We have previously shown that anionic alkylation of indole,⁸⁾ purine and pyrimidine⁹⁾ anions can be performed with simple halides in the presence of a catalytic quantity of tetra-alkyl ammonium salts without any organic solvent. When we attempted to apply this approach to the alkylation of phenothiazine, we found that a very simple procedure worked efficiently, *i.e.*, phenothiazine was directly heated with the commercially available hydrochloride of the alkylating agent, potassium hydroxide and a catalytic quantity of Aliquat 336.¹⁰⁾ The reaction also proceeded at room temperature, but the yields were lower. Neither fresh distillation of the chloroalkylamine nor formation of the anion of the heterocycle was necessary before adding the alkylating agent. In contrast to previous observations¹¹⁾ bulky substituents such as an isopropyl group on the nitrogen did not affect the yield.

This technique is characterized by a very easy work-up, and it avoids the use of any expensive and polluting solvent. It requires smaller amounts of catalyst, a lower reaction temperature and a much shorter reaction time (1 h instead of 18 h) as compared to the most recently reported method.⁶⁾

Experimental

Melting points were taken on a Kofler apparatus and are uncorrected. Proton nuclear magnetic resonance (¹H-NMR) spectra were recorded on a Varian EM 390 spectrometer.¹²⁾

2-Chloro-10-(3-dimethylaminopropyl)phenothiazine (1a, Chlorpromazine)—2-Chlorophenothiazine (2.33 g, 0.01 mol), KOH (1.40 g, 0.025 mol), 3-chlorodimethylaminopropyl hydrochloride (1.62 g, 0.01 mol) and Aliquat 336 (0.120 g, 0.0003 mol) were heated for 1 h at 80 °C. After the mixture had cooled to room temperature 1 g of silica gel was added to retain the catalyst. The mixture was extracted three times with 50 ml of diethyl ether. The crude chlorpromazine was purified by column chromatography (silica gel, ethyl acetate), or the hydrochloride was directly obtained by bubbling hydrogen chloride gas into the ethereal solution. The salt, which separated as an oil, was crystallized by trituration in ethyl acetate. Yield 3.02 g (85%), mp 170 °C. *Anal.* Calcd for C₁₇H₂₀Cl₂N₂S: C, 57.46; H, 5.63; N, 7.89. Found: C, 57.31; H, 5.55; N, 8.12. ¹H-NMR (CDCl₃) δ: 1.90 (2H, m, *J* = 5 Hz), 2.20 (6H, s), 2.35 (2H, t, *J* = 5 Hz), 3.80 (2H, t, *J* = 5 Hz), 6.63–7.15 (7H, m).

The following compounds were prepared in a similar fashion.

10-(2-Dimethylaminoethyl)phenothiazine (1b)—Yield: 78% or 71% after 24 h at 20 °C. Hydrochloride, mp 200 °C. *Anal.* Calcd for C₁₆H₁₉ClN₂S: C, 62.64; H, 6.20; N, 9.14. Found: C, 62.53; H, 6.12; N, 9.21. ¹H-NMR (CDCl₃) δ: 2.25 (6H, s), 2.60 (2H, t, *J* = 5 Hz), 3.95 (2H, t, *J* = 5 Hz), 6.60–7.15 (8H, m).

10-(2-Dimethylamino-2-methylethyl)phenothiazine (1c, Promethazine)¹³⁾—Yield: 71 %. Hydrochloride, mp 230 °C. *Anal.* Calcd for C₁₇H₂₁ClN₂S: C, 63.65; H, 6.24; N, 8.73. Found: C, 63.55; H, 6.10; N, 8.91. ¹H-NMR (CDCl₃) δ: 1.03 (3H, d, *J* = 4 Hz), 2.20 (6H, s), 3.00 (1H, m), 3.61 (1H, m, *J*_{AB} = 8 Hz, *J*_{AX} = 6 Hz), 3.95 (1H, m, *J*_{AB} = 8 Hz, *J*_{BX} = 2 Hz), 6.66–6.81 (8H, m).

10-(2-Dimethylamino-1-methylethyl)phenothiazine (1d, Isopromethazine)—Yield: 12%. Hydrochloride, mp 190 °C. *Anal.* Calcd for C₁₇H₂₁ClN₂S: C, 63.65; H, 6.24; N, 8.73. Found: C, 63.61; H, 6.11; N, 8.91. ¹H-NMR (CDCl₃) δ: 1.58 (3H, d, *J* = 5 Hz), 2.03 (6H, s), 2.66 (1H, m, *J*_{AB} = 8 Hz, *J*_{AX} = 4 Hz).

10-(2-Diisopropylaminoethyl)phenothiazine (1e)—Yield: 84% or 50% after 24 h at 20 °C. Hydrochloride, mp 197 °C. *Anal.* Calcd for C₂₀H₂₇ClN₂S: C, 66.20; H, 7.45; N, 7.72. Found: C, 66.05; H, 7.31; N, 7.78. ¹H-NMR (CDCl₃) δ: 1.03 (6H, d, *J* = 5 Hz), 2.83 (2H, t, *J* = 5.5 Hz), 3.08 (1H, m, *J* = 5 Hz), 3.96 (2H, t, *J* = 5.5 Hz), 6.80–7.26 (8H, m).

2-Chloro-10-(2-diisopropylaminoethyl)phenothiazine (1f)—Yield: 79%. Hydrochloride, mp 170 °C. *Anal.* Calcd for C₂₀H₂₆Cl₂N₂S: C, 64.72; H, 5.84; N, 6.29. Found: C, 64.65; H, 5.75; N, 6.37. ¹H-NMR (CDCl₃) δ: 1.05 (6H, d, *J* = 5 Hz), 2.76 (2H, t, *J* = 5.5 Hz), 3.03 (1H, m, *J* = 5 Hz), 3.80 (2H, t, *J* = 5.5 Hz), 6.70–7.13 (7H, m).

References and Notes

- 1) H. Gilman and D. A. Shirley, *J. Am. Chem. Soc.*, **66**, 888 (1944).
- 2) P. Charpentier, *C. R. Acad. Sci.*, **225**, 306 (1947).
- 3) P. Charpentier and R. Ducrot, *C. R. Acad. Sci.*, **232**, 415 (1951).
- 4) J. B. Ragland and V. J. Kinross-Wright, *Justus Liebigs Ann. Chem.*, **36**, 1356 (1964).
- 5) G. Cauquil, A. Casadevall, and E. Casadevall, *Bull. Soc. Chim. Fr.*, **1960**, 1049.
- 6) S. J. Scholka and H. Zimmer, *Synthesis*, **1984**, 29.
- 7) I. Gozlan, D. Ladkani, M. Halpern, M. Rabinowitz, and D. Avnir, *J. Heterocycl. Chem.*, **21**, 613 (1984).
- 8) J. Barry, G. Bram, G. Decodts, A. Loupy, P. Pigeon, and J. Sansoulet, *Tetrahedron*, **39**, 2669 (1983).
- 9) G. Bram, G. Decodts, Y. Bensaïd, C. Combet-Farnoux, H. Galons, and M. Miocque, *Synthesis*, **1985**, 543.
- 10) Aliquat 336 consists essentially of trioctylmethyl ammonium chloride. C. M. Starks, *J. Am. Chem. Soc.*, **93**, 195 (1971).
- 11) J. B. Wright, E. H. Lincoln, R. V. Heinzelmann, and J. H. Hunter, *J. Am. Chem. Soc.*, **72**, 3536 (1950).
- 12) The ¹H-NMR spectra of the bases are reported.
- 13) Compounds **1c** (promethazine) and **1d** (isopromethazine) were obtained in the same reaction and were separated by column chromatography (silica gel, ethyl acetate).