[Chem. Pharm. Bull.]
[35(8)3502—3506(1987)]

Biological Activities of 3,3'-Dihydroxy- α,β -diethylstilbene and Its Isomer, Diethylstilbestrol

YOSHIHIKO INAMORI,*^a MAYURI KUBO,^a MASAFUMI OGAWA,^a
HIROSHI TSUJIBO,^a YASUYOSHI MIKI^b
and SHOJI TAKEMURA^b

Osaka University of Pharmaceutical Sciences,^a Kawai, Matsubara-shi, Osaka 580,
Japan and Faculty of Pharmaceutical Sciences, Kinki University,^b
3-4-1, Kowakae, Higashiosaka-shi, Osaka 577, Japan

(Received January 19, 1987)

3,3'-Dihydroxy- α,β -diethylstilbene (I) and its isomer, diethylstilbestrol (II), showed antifungal activity against plant-pathogenic fungi, phyto-growth-inhibitory activity and a hypotensive effect on rats. First, I exhibited rather strong antifungal activity against plant-pathogenic fungi; the minimal inhibitory concentrations against *Fusarium oxysporum* f. sp. *lycopersici*, *Botryotinia fuckeliana* and *Pyrenophora graminea* were 4 $\mu\text{g/ml}$. On the other hand, the activity of II was weaker than that of I. Secondly, both compounds inhibited the growth of roots of two plants species even at low concentration of 50 ppm. Thirdly, I and II showed a hypotensive effect on rats (I, -23.33 ± 5.03 mmHg; II, -41.00 ± 2.40 mmHg; 10 mg/kg, i.v.). Unlike II, I has virtually no hormonal activity. It is noteworthy that in spite of the loss of hormonal character in going from II to I, the above-mentioned activities are fully retained.

Keywords—3,3'-dihydroxy- α,β -diethylstilbene; diethylstilbestrol; biological activity; antifungal activity; plant-pathogenic fungus; phyto-growth-inhibitory activity; hypotensive effect; hormonal side effect; phospholipidosis

Various oxystilbene compounds such as 3,3',4,5'-tetrahydroxystilbene (III, Chart 1),^{2,3)} 3,4-*O*-isopropylidene-3,3',4,5'-tetrahydroxystilbene (IV, Chart 1)^{4,5)} and diethylstilbestrol (II, Chart 1)^{6,7)} have been reported to have the following biological activities by the authors; antifungal activity, phyto-growth-inhibitory activity, coronary vasodilator action on the isolated guinea-pig heart, hypotensive effect on rats and ichthyotoxicity. Among these

Chart 1

compounds, II showed the strongest activities, but it had a hormonal side effect. Furthermore, Coralgil (hexestrol bis(β -diethylaminoethyl ether) dihydrochloride, V, Chart 1) a coronary vasodilator having the same basic skeleton as I and II, was found to have the severe side effects of phospholipidosis of the liver⁸ and lung⁹ cells, and foam cell syndrome.¹⁰ Therefore, since 1970 its clinical use has been discontinued.

Subsequently, we synthesized various oxystilbene compounds in order to obtain more active substances. As a result, in the previous paper,¹¹ we reported that 3,3'-dihydroxy- α,β -diethylstilbene (I, Chart 1), an isomer of II, had rather strong ichthyotoxicity and coronary vasodilator action on the isolated guinea-pig heart. However, in contrast to II, I shows almost no side effect.¹² In this respect, the biological activities of I are noteworthy. Therefore, the antifungal activity on plant-pathogenic fungi, the phytogrowth-inhibitory activity and the hypotensive effect on rats of I were examined in detail in comparison with those of II. The results are presented here.

Materials and Methods

Chemicals—3,3'-Dihydroxy- α,β -diethylstilbene (I) and diethylstilbestrol (II, Aldrich Chemical Co., Ltd.) were used for the biological activity tests. Compound I was synthesized according to the method of Dodds *et al.*¹³ I: mp 176–178 °C (reported value: mp 176 °C). ¹H-NMR (CDCl₃, δ ppm): 0.77 (6H, t, J = 8 Hz, $2 \times -CH_3$), 2.13 (4H, q, J = 8 Hz, $2 \times -CH_2-$), 6.65–7.25 (8H, m, arom H), 4.65 (2H, s, $2 \times -OH$). Sodium 2,4-dichlorophenoxyacetate (Tokyo Kasei Co., Ltd.) was used as a standard for the phytogrowth-inhibitory activity test.

Organisms—The plant-pathogenic fungi were as follows: *Botryotinia fuckeliana* IFO-9760, *Pyrenophora graminea* IFO-6633, *Rhizoctonia solani* IFO-30464, *Cochliobolus miyabeanus* IFO-4870, *Ceratocystis fimbriata* IFO-4864, *Fusarium oxysporum* f. sp. *lycopersici* IFO-6531 and *Aureobasidium pullulans* IFO-4464. The plants used were *Brassica rapa* and *Raphanus sativus* L. var. *raphanistroides* MAKINO. The animals used for the experiments on blood pressure were male Wistar strain rats weighing 305–355 g (3 rats/group).

Biological Activity Tests—1) Antifungal Activity Test⁵: Antifungal activity was tested by the agar dilution method. The media used were as follows: potato sucrose agar in all cases except for *Fusarium oxysporum* f. sp. *lycopersici* (potato dextrose agar: Eiken Chemical Co., Ltd.). The test fungi were applied to media containing various concentrations of I and II. The plates were incubated at 27 °C for 5 d and the growth was observed with the naked eye.

2) Phytogrowth-Inhibitory Activity Test¹⁴: Aliquots (1 ml) of acetone solutions of I, II and sodium 2,4-dichlorophenoxyacetate were each diluted in 100 ml of sterilized agar (0.8%, Difco Chemical Co., Ltd.) to the concentration of 50 ppm. The agar containing chemicals or acetone alone (control) was poured into a 500 ml sterilized beaker covered with aluminum foil. Then, 20 seeds of each plant sterilized with 70% EtOH and 1% NaClO were put on the agar and left for 7 d under 600 lux illumination. The lengths of the roots were measured and averaged. The phytogrowth-inhibitory activity was expressed as the ratio of the length of roots to that of the control (1.00).

3) Measurement of Blood Pressure⁵: Systemic blood pressure was measured with a pressure transducer (Nihon Kohden Kogyo Co., Ltd. P-23 ID, AP-601 G) following cannulation of the carotid artery in ras under anesthesia with sodium pentobarbital (40 mg/kg, i.p.). Compounds I and II were suspended in 5% acacia and administered *via* the femoral vein. It was shown that 5% acacia has no effect on blood pressure.

Results

Antifungal Activities of 3,3'-Dihydroxy- α,β -diethylstilbene (I) and Diethylstilbestrol (II) on Plant-Pathogenic Fungi

The antifungal activities of 3,3'-dihydroxy- α,β -diethylstilbene (I) and diethylstilbestrol (II) were investigated by the agar dilution method. The results are summarized in Table I. Compounds I and II showed antifungal activities against all plant-pathogenic fungi tested except for *Ceratocystis fimbriata*. The activity of I was stronger than that of II. The minimal inhibitory concentrations (MIC) of I for *Fusarium oxysporum* f. sp. *lycopersici*, *Botryotinia fuckeliana* and *Pyrenophora graminea* were 4 μ g/ml.

Phytogrowth-Inhibitory Activities of 3,3'-Dihydroxy- α,β -diethylstilbene (I) and Diethylstilbestrol (II)

The inhibitory effects of 3,3'-dihydroxy- α,β -diethylstilbene (I) and diethylstilbestrol (II)

TABLE I. Antifungal Activities of 3,3'-Dihydroxy- α,β -diethylstilbene (I) and Diethylstilbestrol (II) on Plant-Pathogenic Fungi

Fungi	MIC ($\mu\text{g/ml}$)	
	I	II
<i>Fusarium oxysporum</i> f. sp. <i>lycopersici</i> IFO-6531	4.0	7.0
<i>Botryotinia fuckeliana</i> IFO-9760	4.0	50.0
<i>Pyrenophora graminea</i> IFO-6633	4.0	10.0
<i>Aureobasidium pullulans</i> IFO-4464	25.0	25.0
<i>Cochliobolus miyabeanus</i> IFO-4870	25.0	50.0
<i>Rhizoctonia solani</i> IFO-30464	25.0	50.0
<i>Ceratocystis fimbriata</i> IFO-4864	1000.0	> 1000.0

Culture conditions: 27°C, 5 d. Media: potato sucrose agar (*Fusarium oxysporum* f. sp. *lycopersici* IFO-6531, potato dextrose agar). Method: agar dilution method.

TABLE II. Inhibitory Effects of 3,3'-Dihydroxy- α,β -diethylstilbene (I) and Diethylstilbestrol (II) on Plant Growth

Plant	Growth (ratio) ^{a)}		
	I	II	2,4-D ^{b)}
<i>Brassica rapa</i> L.	0.46	0.43	0.06
<i>Raphanus sativus</i> L. var. <i>rahanistroides</i> MAKINO	0.83	0.65	0.10

- a) Growth in control experiments after 7 d was taken as 1.00.
Concentration: 50 ppm.
Quantity of light: 600 lux.
Experimental size: 20 seeds/group, 2 groups.
b) Sodium 2,4-dichlorophenoxyacetate.

TABLE III. Effects of 3,3'-Dihydroxy- α,β -diethylstilbene (I) and Diethylstilbestrol (II) on Blood Pressure in Rats

Dose (mg/kg)	Mean arterial blood pressure (mmHg)	
	I	II
10	-23.33 $\pm$ 5.03	-41.00 $\pm$ 2.40
20	-45.67 $\pm$ 2.52	-48.00 $\pm$ 14.0
30	-52.66 $\pm$ 0.58	-51.00 $\pm$ 12.0

Each value represents the mean $\pm$ S.D. of 3 rats. Route: Intravenous injection.

on two plant species were examined according to the previous paper.¹⁴⁾ As shown in Table II, compounds I and II inhibited the growth of roots both kinds of plants even at the low concentration of 50 ppm, but their activities were weaker than that of sodium 2,4-dichlorophenoxyacetate used as a standard.

Effect of 3,3'-Dihydroxy- α,β -diethylstilbene (I) and Diethylstilbestrol (II) on Blood Pressure in Rats

The effects of 3,3'-dihydroxy- α,β -diethylstilbene (I) and diethylstilbestrol (II) on blood pressure in rats were investigated. The results are summarized in Table III. Both compounds showed hypotensive action. In the group given 10 mg/kg, the action of II was stronger than that of I. However, at the doses of 20 and 30 mg/kg, I showed almost the same activity as II. The hypotensive actions of both compounds were transient and the blood pressure recovered to the original level within 5 min. The patterns of the hypotensive actions of both compounds were similar to those of III³⁾ and IV.⁵⁾

Discussion

Like other oxystilbene compounds,²⁻⁵⁾ 3,3'-dihydroxy- α,β -diethylstilbene (I) and diethylstilbestrol (II) showed antifungal activity on plant-pathogenic fungi, as well as

phytogrowth-inhibitory activity and hypotensive action on rats.

First, compounds I and II showed rather strong antifungal activity against plant-pathogenic fungi (Table I). The antifungal activities of both compounds on plant-pathogenic fungi might be intrinsic to oxystilbene compounds. This view is supported by the findings that 3,4-*O*-isopropylidene-3,3',4,5'-tetrahydroxystilbene (IV),⁵⁾ which has the same skeleton as I, exhibited strong antifungal activity against plant-pathogenic fungi, and pinosylvine¹⁵⁾ and pterostilbene,¹⁶⁾ which also have the same skeleton as I, were reported to be phytoalexins. Further studies on the antifungal activities of many oxystilbene compounds against plant-pathogenic fungi should be fruitful.

Secondly, compounds I and II inhibited the growth of roots of two species of plants even at low concentration of 50 ppm (Table II). As regards phytogrowth-inhibitory activity of oxystilbene compounds, 3,3',4,5'-tetrahydroxystilbene (III)²⁾ was reported to have rather strong activity. Plant growth regulators such as caffeic acid,¹⁷⁾ ferulic acid¹⁷⁾ and chlorogenic acid,¹⁷⁾ which have a partial structure in common with I, have been isolated from higher plants. Batatasin III¹⁸⁾ and lunularin,¹⁹⁾ having the same basic skeleton as I and II, were found to have phytogrowth-inhibitory activities. These findings suggest that oxystilbene compounds generally have phytogrowth-inhibitory activity. Further studies on the phytogrowth-inhibitory activity of many compounds having the same partial structures as I are in progress.

Thirdly, compounds I and II had a hypotensive effect on rats (Table III). The hypotensive effects of both compounds on rats were considered to be intrinsic to oxystilbene compounds, because II,²⁰⁾ III³⁾ and IV⁵⁾ also show similar effects. In addition to the above-mentioned oxystilbene compounds, stilbamidine²¹⁾ and dimethylstilbamidine,²¹⁾ antiprotozoal substances which are also stilbene compounds, show potent hypotensive activity. We are now investigating the hypotensive effects of many oxystilbene compounds.

It is very interesting that the above-mentioned activities of I were retained in spite of the loss of the hormonal side effect on shifting the phenolic hydroxyl group to the *m*-position (from II to I).¹²⁾ However, as stated above, Coralgil (V), in spite of its strong coronary vasodilator action, is no longer used clinically because of the severe side effects of phospholipidosis of the liver⁸⁾ and lung⁹⁾ cells, and foam cell syndrome.¹⁰⁾ Thus, I and other oxystilbene compounds (II—IV) need to be tested carefully for similar side effects. Such a study on I and other oxystilbene compounds (II—IV) is in progress.

Acknowledgement The authors thank Professor Mitsugi Kozawa, Osaka University of Pharmaceutical Sciences, for valuable advice during this project.

References and Notes

- 1) This work was presented at the 36th Kinki Regional General Meeting of the Japanese Society of Pharmacy, Osaka, November 1986.
- 2) Y. Inamori, Y. Kato, M. Kubo, M. Yasuda, K. Baba and M. Kozawa, *Chem. Pharm. Bull.*, **32**, 213 (1984).
- 3) Y. Inamori, M. Kubo, Y. Kato, M. Yasuda, K. Baba and M. Kozawa, *Yakugaku Zasshi*, **104**, 819 (1984).
- 4) Y. Inamori, M. Kubo, Y. Kato, M. Yasuda, K. Baba and M. Kozawa, *Chem. Pharm. Bull.*, **32**, 801 (1984).
- 5) Y. Inamori, M. Kubo, Y. Kato, M. Yasuda, H. Tsujibo, K. Baba and M. Kozawa, *Chem. Pharm. Bull.*, **33**, 2904 (1985).
- 6) Y. Inamori, M. Kubo, M. Ogawa, M. Moriwaki, H. Tsujibo, K. Baba and M. Kozawa, *Chem. Pharm. Bull.*, **33**, 420 (1985).
- 7) Y. Inamori, M. Kubo, M. Ogawa, M. Moriwaki, H. Tsujibo, K. Baba and M. Kozawa, *Chem. Pharm. Bull.*, **33**, 4478 (1985).
- 8) A. Yamamoto, S. Adachi, T. Kitani, T. Nasu, Y. Shinji and M. Nishikawa, *Saishin-Igaku*, **26**, 2209 (1971).
- 9) T. Shikata, Y. Endo, T. Orita, T. Kanetaka and Z. Takahashi, *Acta Hepatologica Japonica*, **12**, 256 (1971).
- 10) Y. Onozawa and M. Komiya, *The Japanese Society of Internal Medicine*, **59**, 1228 (1970).
- 11) Y. Inamori, M. Kubo, H. Tsujibo, M. Ogawa, Y. Saito, Y. Miki and S. Takemura, *Chem. Pharm. Bull.*, **35**, 887 (1987).

- 12) D. Adam and H. Schonenberger, *Arzneim.-Forsch.*, **16**, 738 (1966).
- 13) E. C. Dodds, L. Goldberg, W. Lawson and R. Robinson, *Nature* (London), **141**, 247 (1938).
- 14) Y. Inamori, M. Kubo, H. Tsujibo, M. Ogawa, K. Baba, M. Kozawa and E. Fugita, *Chem. Pharm. Bull.*, **34**, 3928 (1986).
- 15) L. Shain, *Phytopathology*, **57**, 1034 (1967).
- 16) H. Orth and P. Forschner, *Holzforschung*, **19**, 111 (1970).
- 17) R. del Moral and C. H. Muller, *Am. Midl. Nat.*, **83**, 254 (1970).
- 18) Y. Hashimoto, K. Hasegawa and A. Kawarada, *Planta* (Berl.), **108**, 369 (1972).
- 19) J. Gorham, *Phytochemistry*, **16**, 249 (1977).
- 20) K. Kondo, T. Okuno, T. Eguchi, T. Yasui, H. Suzuki, S. Nagahama and T. Saruta, *Endocrinol. Jpn.*, **27**, 307 (1980).
- 21) J. N. Ashley, H. J. Barber, A. J. Ewins, G. Newbery and A. D. H. Self, *J. Chem. Soc.*, **1942**, 103.