Communications to the Editor

[Chem. Pharm. Bull.]
36(2) 834-836 (1988)

BOERAVINONE A AND B, NEW ROTENOID ANALOGUES FROM BOERHAAVIA DIFFUSA LINN

Shigetoshi Kadota, Nzunzu Lami, Yasuhiro Tezuka, and Tohru Kikuchi*

Research Institute for Wakan-Yaku (Oriental Medicines), Toyama Medical
and Pharmaceutical University, 2630 Sugitani, Toyama 930-01, Japan

Two new rotenoid analogues, boeravinone A and B, have been isolated from the roots of Boerhaavia diffusa Linn. (Nyctaginaceae) and their structures have been determined based on the 2-D NMR spectroscopy including INADEQUATE and ^1H - ^{13}C long range COSY.

KEYWORDS — Boerhaavia diffusa; Nyctaginaceae; rotenoid; boeravinone A; boeravinone B

The roots of Boerhaavia diffusa Linn. (Nyctaginaceae) are used as a folk medicine for treatment of asthma and abdominal tumor in India.¹⁾ From this plant new rotenoid analogues have been isolated and their structures elucidated.

The ether extract from the dried roots (1.2 kg) of B. diffusa, collected in Mannang, Nepal, was separated by silica gel column chromatography followed by repeated preparative TLC to give three new rotenoid analogues, named boeravinone A (125 mg), B (41 mg), and C (129 mg), together with known sterols, sterol glucosides, and fatty acids. This paper deals with the structure elucidation of boeravinone A (1) and B (2).

Boeravinone A (1), mp 215-217°C, yellow needles (from MeOH-CH₂Cl₂), $[\alpha]_D^{20}$ (acetone), has the molecular formula C₁₈H₁₄O₆ and showed IR (KBr) absorption at 3250 (OH), 1650 (conj. CO), and 1620 cm⁻¹ and UV (EtOH) bands at 340, 300, 276, and 217 nm (log ϵ : 3.39, 4.07, 4.37, and 4.39). Its mass spectrum exhibited the M⁺ peak at m/z 326 and fragment ion peaks at m/z 295 (M⁺-OCH₃) and 267 (M⁺-OCH₃-CO).

The ^1H - and ^{13}C -NMR spectra of 1 indicated the presence of a carbonyl (δ_{C} 180.4), five aromatic CH groups (δ_{H} 6.56, 7.16, 7.20, 7.29, 9.10; δ_{C} 93.53, 122.91, 117.48, 129.06, 127.48), a hydroxyl (δ_{H} 13.44), a methoxy group (δ_{H} 3.45; δ_{C} 55.73), and a t-methyl group (δ_{H} 2.39; δ_{C} 7.92) (Table I). Each carbon signal except the quarternary one was assigned based on the ^1H - ^{13}C COSY spectral data.²⁾

- 1: R₁ = CH₃, R₂ = H
2: R₁, R₂ = H
3: R₁, R₂ = CH₃

Methylation of **1** with diazomethane gave a methyl ether (**3**), $C_{19}H_{16}O_6$ (amorphous powder). The 1H -NMR spectrum of **3** showed a singlet due to a newly introduced methoxy group (δ 3.91), together with a one-proton singlet (δ 12.94) which could be ascribed to a phenolic OH, strongly hydrogen-bonded with the 12-ketone group (Table I).

Boeravinone A was found to have a number of quarternary carbons (Table I). So the 2-D INADEQUATE spectrum²⁾ of **1** was measured to clarify the sequence of carbon atoms in the molecule. The result indicated the presence of partial structures (A, B, and C) as depicted in **4**.

Next, we measured the 1H - ^{13}C long range COSY³⁾ of **1** in order to confirm the connectivities of the partial structures and substituent groups. As shown in the formula **5**, the carbon signal at δ 110.12 (C-12a) is correlated with the protons at δ 5.89 (6-H) and 9.10 (1-H), and the signal at δ 95.33 (C-6) is correlated with the proton at δ 3.45 (OCH_3). Also, the carbon signals at δ 109.12 (C-10), 149.05 (C-5a), and 160.54 (C-11) are correlated with the proton signals at δ 2.39 (CH_3), 6.56 (8-H), and 13.44 (11-OH), at δ 5.89 (6-H), 7.20 (4-H), 7.29 (3-H), and 9.10 (1-H), and at δ 2.39 (CH_3), and 13.44 (11-OH), respectively. Some of the other significant long range correlations observed are also shown by arrows.

Although connectivity could not be detected between carbons 12 and 11a (each quarternary carbon), there must be a chemical bonding between them since the 11-OH is internally hydrogen-bonded with the 12-ketone group. Also, in view of the composition, an ether linkage should exist between quarternary carbons 6a and 7a.

On the basis of the above results, the structure of boeravinone A was determined to be **1**.

Boeravinone B (**2**) was obtained as an amorphous yellow powder, $[\alpha]_D^{20}$ 0° (acetone), and has the molecular formula $C_{17}H_{12}O_6$. Its 1H -NMR spectrum showed a pattern very similar to that of **1**, except for the absence of a singlet due to the methoxy group (Table I), indicating that the structure is **2**.

Treatment of boeravinone B (**2**) with p-TsOH in MeOH gave a methyl acetal, mp 207–209°C, which was found to be identical with boeravinone A (**1**) by IR and 1H -NMR comparisons. Therefore, the structure of boeravinone B was established to be **2**.

The biological activities of **1** and **2** are currently under investigation.

Rotenoids,⁴⁾ which are shown to possess an insecticidal activity and other interesting biological activities,⁵⁾ have so far been isolated only from the Leguminosae and Stemonaceae plants. Our present result provided the first example of the isolation of rotenoid analogues from a plant of the Nyctaginaceae family. Distribution of rotenoid analogues in other Nyctaginaceae plants and their biological activities would be a problem of particular interest.

Table I. ^1H (400 MHz) and ^{13}C (100 MHz) NMR Spectra of Boeravinone A (1), B (2), and the Dimethyl Ether Derivative (3)

Position	Boeravinone A (1) (pyridine d_5)		Boeravinone B (2) (pyridine d_5)		Dimethyl ether (3) (CDCl_3)	
	$\delta \text{H}^{a,b}$	$\delta \text{C}^{c,d}$	$\delta \text{H}^{a,b}$	$\delta \text{C}^{c,d}$	δH^a	δC^e
1	9.10 dd (8.0,1.3)	127.48 d	9.14 dd (8.0,1.2)	127.44 d	8.85 dd (8.0,1.5)	127.23
1a	—	117.77 s	—	118.00 s	—	117.10
2	7.16 ddd (8.0,7.6,1.3)	122.91 d	7.16 ddd (8.0,6.0,3.0)	122.53 d	7.17 ddd (8.0,7.5,1.5)	122.98
3	7.29 ddd (8.0,7.6,1.3)	129.06 d	7.30 m	128.96 d	7.30 ddd (8.0,7.5,1.5)	128.96
4	7.20 dd (8.0,1.3)	117.48 d	7.30 m	118.00 d	7.12 dd (8.0,1.5)	116.97
5a	—	149.05 s	—	150.06 s	—	148.37
6	5.89 s	95.33 d	6.52 s	89.53 d	5.80 s	95.06
6-OMe	3.45 s	55.73 q	—	—	3.60 s	55.95
6a	—	154.82 s	—	155.19 s	—	154.40
7a	—	155.06 s	—	155.77 s	—	155.00
8	6.56 s	93.53 d	6.65 s	93.58 d	6.45 s	89.35
9	—	164.11 s	—	164.08 s	—	163.64
9-OMe	—	—	—	—	3.91 s	56.04
10	—	109.12 s	—	109.00 s	—	109.73
10-Me	2.39 s	7.92 q	2.42 s	8.08 q	2.12 s	7.37
11	—	160.54 s	—	160.59 s	—	159.21
11a	—	105.32 s	—	105.28 s	—	106.16
11-OH	13.44 s	—	13.67 s	—	12.94 s	—
12	—	180.40 s	—	180.69 s	—	180.28
12a	—	110.12 s	—	109.47 s	—	110.54

δ values in ppm. a) Coupling constants in Hz. b) ^1H - ^1H Correlation spectra were measured. c) The multiplicities of carbon signals were determined by means of the DEPT method, and are indicated as s, d, t, and q. d) ^1H - ^{13}C Correlation spectra were measured. e) Only the complete decoupling spectrum was measured.

ACKNOWLEDGEMENT This work was supported in part by a Grant-in-Aid for Overseas Scientific Survey (No. 58041031), and a Grant-in-Aid for Scientific Research (No. 61470147) from the Ministry of Education, Science and Culture of Japan.

REFERENCES AND NOTES

- 1) A. N. Misra and H. P. Tiwari, *Phytochemistry*, **10**, 3318 (1971) and references therein.
- 2) A. Bax, "Two-Dimensional NMR in Liquids," D. Reidal Publishing Co., Dordrecht, Holland, (1982); R. Benn and H. Gunther, *Angew. Chem., Int. Ed. Engl.*, **22**, 350 (1983).
- 3) C. Francisco, B. Banaigs, and J. Teste, *J. Org. Chem.*, **51**, 1115 (1986).
- 4) J. B. Harbone and T. J. Mabry, "The Flavonoids: Advances in Research," Chapman and Hall (1982).
- 5) For example, crocodile repellent and piscicidal activities have been reported. See N. Finch and W. D. Ollis, *Proc. Chem. Soc.*, **1960**, 176; W. D. Ollis, C. A. Rhodes and I. O. Sutherland, *Tetrahedron*, **23**, 4741 (1967).

(Received November 9, 1987)