Stereoselective Reactions. XVI.¹⁾ Total Synthesis of (–)-β-Bourbonene by Employing Asymmetric (2+2) Photocycloaddition Reaction of Chiral Butenolide

Kiyoshi TOMIOKA, Masahide TANAKA and Kenji KOGA*

Faculty of Pharmaceutical Sciences, University of Tokyo, Hongo, Bunkyo-ku, Tokyo 113, Japan. Received September 28, 1988

Total synthesis of (–)-β-bourbonene (3) was achieved by employing the (2+2) photocycloaddition reaction of (S)-γ-pivaloyloxymethyl-β-methyl-γ-butenolide (5e) with cyclopentenone ethyleneacetal (15). The major photoadduct (16) was readily converted to the *cis,anti,cis*-tricyclo[5.3.0.0^{2,6}]decane system (22) in 70% yield. Further transformation provided the optically pure title compound (3).

Keywords photocycloaddition; β-bourbonene; spatol; tricyclo[5.3.0.0^{2,6}]decane; stereoselectivity; regioselectivity; cytotoxic diterpene; butenolide; asymmetric synthesis; total synthesis

The *cis,anti,cis*-tricyclo[5.3.0.0^{2,6}]decane ring system is found in spatane diterpenes (1,2)²⁾ and bourbonene sesquiterpenes (3,4).³⁾ Spatol (1), a representative of spatane diterpenes isolated from brown marine algae of the family Dictyotaceae, is known to be endowed with remarkable biological properties including a potent inhibition of cell replication.^{2c,d)} The uniqueness of the carbon framework of these terpenes coupled with their biological activities has attracted much effort aimed at total synthesis of these terpenes.^{4,5)} For the purpose of total synthesis of spatol and related spatane diterpenes,⁶⁾ we have initiated studies on the development of new methodology for the construction of the optically active *cis,anti,cis*-tricyclo[5.3.0.0^{2,6}]decane ring system. We describe here the details of our approach and total syntheses of (–)-β-bourbonene (3) and (–)-norbourbonene (4), representatives of bourbonene sesquiterpenes.⁷⁾

Synthetic Design The general approach under consideration for the synthesis of the requisite tricyclodecane ring system in optically active form involves the diastereoselective (2+2) photocycloaddition reaction^{8,9)} of an optically active γ-hydroxymethyl-β-methyl-γ-butenolide (5b)^{10,11)} with a cyclopentenone derivative (Chart 1). It is possible to predict the stereochemistry at the newly created chiral centers of B on the basis of the least hindered approach.^{11c)} The γ-lactone part of the photoadduct (B) is

Chart 1

then converted to the cyclopentenone portion of C. It is noteworthy that A is used not only as a stereocontrolling factor but also as a building block of cyclopentenone, the so-called chiral synthon.

Stereochemistry of the Cycloaddition Photocycloaddition reaction of γ-butyrolactone with 1,1-dimethoxyethylene has been reported by Uda *et al.* to give two diastereomers in an almost non-stereocontrolled manner.¹²⁾ Our studies began with irradiation of a solution of 5a (R = CH₂OCPh₃)^{10a)} and cyclopentenone in acetonitrile using a low pressure mercury lamp at 15°C under argon. The products were isolated by column chromatography or by recrystallization. The ratio of the two diastereomers was determined by nuclear magnetic resonance (NMR) spectroscopy and by isolation (Table I).

The stereochemistry was assigned based on the NMR characteristics of 6a and 7a. The *cis,anti,cis* arrangement of adducts was ascertained by NMR spectroscopy. The coupling constants of the carbonyl α-proton signal appearing at 2.11 (minor) and at 2.30 (major) ppm as a doublet were 4 Hz, corresponding to the desired arrangement.^{12,13)} Furthermore, the signals of angular methyl carbon of the major and minor products appeared at 12.8 and 19.4 ppm, respectively (Table II). Judging from steric compression

TABLE I. Diastereoface Differentiation in the (2+2) Photoaddition

Run	5	R	Yield (%) ^{a)}	6:7 ^{b)}
1 ^{c)}	5a	CH ₂ OC(Ph) ₃	29 (48)	60:40
2	5b	CH ₂ OH	86	56:44 ^{d)}
3	5c	CH ₂ OCH ₂ OCH ₃	80	50:50 ^{d)}
4	5d	CH ₂ OCOCH(CH ₃) ₂	62	62:38
5	5e	CH ₂ OCO(CH ₃) ₃	81	71:29
6 ^{e)}	5e	CH ₂ OCO(CH ₃) ₃	30 (95)	79:21
7	5f	CH ₂ OSi(CH ₃) ₂ C(CH ₃) ₃	64	68:32
8	5g	C(CH ₃) ₂ OH	57	65:35

a) Yields refer to isolated 6 and 7. Numbers in parentheses are corrected yields based on the consumed 5. b) The ratios were determined based on the isolation of 6 and 7, except runs 2 and 3. c) The reaction was carried out for 16 h. d) The ratios were determined by the ¹H-NMR spectroscopy (angular methyl proton). e) The reaction was carried out at –30°C.

TABLE II. ^{13}C -NMR Spectroscopic Data for **6** and **7**^{a)}

	6a	7a ^{b)}	6d	7d	6e	7e	6f	7f	6g	7g
C1, C2 and C7	47.57 d	38.70 d	41.33 d	38.75 d	41.47 d	38.74 d	42.01 d	38.70 d	40.26 d	38.84 d
	47.61 d	41.23 d	47.66 d	41.38 d	47.62 d	41.38 d	47.57 d	41.47 d	47.86 d	40.94 d
	48.98 d	48.59 d	48.74 d	48.59 d	48.74 d	48.64 d	49.27 d	48.93 d	50.44 d	50.00 d
C3	179.4 s	179.2 s	178.6 s	179.0 s	178.7 s	179.0 s	179.4 s	179.4 s	178.4 s	173.8 s
C5	89.09 d	87.38 d	87.48 d	87.17 d	87.29 d	86.07 d	89.68 d	88.41 d	95.67 d	94.21 d
C6	41.04 s	40.11 s	41.04 s	40.35 s	40.99 s	40.35 s	41.13 s	40.31 s	42.99 s	40.69 s
C8, C9 and C10	26.07 t	25.83 t	25.98 t	25.88 t	26.03 t	25.88 t	26.37 t	25.98 t	25.78 t	26.27 t
	27.49 t	27.88 t	27.44 t	27.93 t	27.49 t	27.93 t	27.68 t	27.93 t	26.66 t	27.73 t
	32.29 t	32.41 t	32.26 t	32.51 t	32.26 t	32.55 t	32.36 t	32.65 t	31.92 t	32.51 t
C11	12.82 q	19.35 q	13.06 q	19.10 q	13.06 q	19.20 q	12.87 q	19.98 q	14.57 q	21.20 q
C12	62.77 t	62.82 t	63.21 t	63.45 t	63.26 t	63.41 t	62.29 t	61.99 t	71.93 s	71.59 s
C on R	87.14 s	87.09 s	18.86 q	18.91 q	27.10 q	27.15 q	-5.60 q	-5.56 q	25.15 q	23.54 s
	127.1 d		33.92 q	33.87 d	38.70 s	38.74 s	18.03 s	-5.41 q	28.95 q	31.00 q
	127.8 d		176.5 s	176.7 s	177.8 s	179.0 s	25.73 q	18.13 s		
	128.6 d							25.78 q		
	143.5 s									

a) Expressed in ppm relative to internal tetramethylsilane in CDCl_3 . b) Data obtained from the spectrum of a mixture of **6a** and **7a**.

effects, these lead to the assignment of **6a** and **7a** as the major and minor adducts, respectively. It is important to note that the diastereofacial differentiation in photocycloaddition of chiral butenolide (**5a**) is consistent with the least hindered approach, giving **6a** as a major product. However, efficiency was poorer than we had expected.^{11c)}

Photocycloaddition of several butenolides (**5**) with substituents of various steric sizes was then studied with regard to diastereofacial differentiation (Table I). The ^{13}C -NMR data are summarized in Table II. Chiral butenolides with a variety of side chains were prepared from the corresponding alcohol (**5b**)¹⁴⁾ and from epoxide (**8**) (Chart 2). Efficiency in the diastereofacial differentiation correlates to the bulkiness of the substituents; **5b, c** showed almost no selection and **5e** with the pivaloyl moiety gave the highest ratios of 71:29 at 15°C and 79:21 at -30°C. Compared to a complete stereoselection in the enolate alkylation of **5a**,^{11c)} the poorer efficiency is probably attributable to the intermediacy of diradical species **11**—**14**.⁸⁾ Since **13** and **14** would not be more stable than **11** and **12**, the distribution of

products would be governed by the intermediacy of **11** and **12**. Least hindered approach of cyclopentene and **5** would lead to the predominant formation of **11** rather than **12** at the first bond-forming stage. However, the severe steric repulsion between methyl and R groups in the *cis* arrangement would destabilize **11** at the second bond forming stage and **11** would revert to the olefin and **5**, giving rise to formation of both **11** and **12**.

Having established the stereochemistry in (2+2) photocycloaddition of the chiral butenolide (**5**), we turned our attention to the synthesis of β -bourbonene (**3**).

Photocycloaddition of 5e with 15 A (2+2) photocycloaddition of **5e** with **15**¹⁵⁾ under the same conditions described above afforded the four products **16**—**19** in a ratio of 3.1:1.3:1:1. Purification of the crude products by column chromatography afforded an inseparable mixture of **16** and **17** in 44% yield as well as **18** and **19** in 10 and 9.7% yields, respectively. The major product **16** was nicely isolated in 25% yield by recrystallization of the mixture.

The angular methyl carbon signals of these isomers **16**—**19** appeared at 13.5, 15.1, 19.8, and 21.2 ppm, respectively, indicating that **16** and **17** are the products arising from the least-hindered approach as in **6**, and **18** and **19** are those arising from the approach at the opposite face as in **7**. The ^{13}C -NMR data are summarized in Table III.

Furthermore **16** was converted to (-)- β -bourbonene (**3**) of natural configuration via the intermediate **22** (*vide infra*), indicating that major photoadduct is **16**. By the same transformation, the three isomers **17**—**19** were converted to **24**, **26**, and **28**, respectively.

These chemical correlations and ^{13}C -NMR spectroscopic analyses allowed the assignment of product structures.

Preferred formation of **16** over **17**, a regioisomer due to ethylene acetal group, is probably attributable to the lower steric congestion caused by interaction with a linear ar-

TABLE III. ¹³C-NMR Spectroscopic Data for 16–19^{a)}

	16	17	18	19
C1, C2 and C7	44.50 d 45.52 d	49.08 d 52.49 d	37.04 d 44.69 d 45.32 d	38.45 d 44.30 d 48.98 d
C3, C15	177.4 s 177.6 s	178.2 s	178.0 s	178.0 s 178.4 s
C5	86.80 d	87.29 d	85.97 d	86.60 d
C6	40.89 s	41.38 s	40.79 s	40.79 s
C8, C9	23.93 t 34.31 t	117.9 s 29.37 t	24.37 t 34.36 t	118.4 s 29.53 t
C10	117.0 s	34.41 t	117.4 s	34.21 t
C11, C12 and C14	62.92 t 63.99 t 64.62 t	64.33 t 65.10 t	63.16 t 64.33 t 64.97 t	63.06 t 65.60 t
C13	13.45 q	15.06 q	19.84 q	21.15 q
C16	38.40 s	38.40 s	38.79 s	38.75 s
C17	26.80 q	27.10 q	27.15 q	27.10 q

a) Expressed in ppm relative to internal tetramethylsilane in CDCl₃.

rangement of ethyleneacetal, methyl, and pivaloyloxy-methyl groups as shown in 29.

Synthesis of Tricyclo[5.3.0.0^{2,6}]decane Ring The major product 16 was transformed to 22 in a sequence of reactions without any problem. Sequential treatment of 16 with methyl lithium in tetrahydrofuran (THF) (to afford 20), sodium methoxide in methanol, and sodium metaperiodate in water–ethyl acetate (AcOEt) furnished a keto-aldehyde 21, which was finally treated with sodium hydroxide in water–ether to afford the desired product 22 in 70% overall yield from 16.

The other three isomers 17–19 were also converted to 24, 26 and 28 in 52, 73, and 58% yields, respectively. Excepting the sign of the optical rotation, 22 and 26, 24 and 28 were identical with each other in spectroscopic comparisons.

Synthesis of (–)-β-Bourbonene (3) Having accomplished the construction of the desired optically active tricyclo[5.3.0.0^{2,6}]decane system (22) according to the our

original design, the remaining steps to 3 were uneventful manipulations. Then 22 was hydrogenated to 30 and reduced with sodium borohydride stereoselectively to give 31 as a single isomer by the attack of hydride from the less hindered side of the carbonyl group in 30. Tosylation followed by substitution of the tosyloxy group of 32 with sodio diethyl malonate proceeded with inversion of configuration at the carbon atom to give 33, in which the requisite carbons for the isopropyl group were introduced in the correct stereochemistry. Reduction of the malonic ester moiety to the isopropyl group of 36 and transacetalization gave norbourbonone (4)^{3a)} as a colorless solid. The melting point and specific rotation of synthetic 4 were in good agreement with those reported.^{3a)}

Finally, methylenation of 4 with methylene triphenyl-

phosphorane completed the synthesis to give (–)-3 as an oil. The ¹H-NMR, infrared (IR), and mass (MS) spectra data and specific rotation of synthetic (–)-3 were in good agreement with those reported for natural 3 isolated from *Geranium Bourbon*,^{3b} and the IR and MS spectra obtained were superimposable on those of 3 provided by Dr. Yoko Naya.

Conclusion

By (2 + 2) photocycloaddition of optically active butenolides with cyclopentene as a key step, construction of the optically pure tricyclo[5.3.0.0^{2,6}]decane ring system was accomplished, and this was successfully applied to the first total synthesis of optically pure β-bourbonene. This success, combined with the ready availability of both enantiomers of butenolides (5),¹⁴ made it possible to access the optically pure tricyclo[5.3.0.0^{2,6}]decanes in both enantiomers, and was applied to the first total synthesis of optically pure spatane diterpenes (1, 2).⁶

Experimental¹⁶

(±)-4-Hydroxymethyl-3-methyl-2-buten-4-olide (5b) A solution of (±)-5a (2.04 g, 5.5 mmol) in methanol (MeOH) (180 ml) containing 12 N HCl (18 ml) was stirred at room temperature for 3 h. After concentration under reduced pressure, the residue was purified by silica gel column chromatography (ether) to give (±)-5b (0.59 g, 84%) as colorless prisms of mp 59–60 °C (from benzene). IR $\nu_{\text{max}}^{\text{KBr}}$ cm^{–1}: 3370 (OH), 1730 (C=O), 1640 (C=C). ¹H-NMR (CDCl₃) and MS spectra were identical with those of (–)-5b.

(±)-4-Methoxymethoxymethyl-3-methyl-2-buten-4-olide (5c) A solution of (±)-5b (105 mg, 0.82 mmol), diisopropylethylamine (0.16 ml, 0.92 mmol), and chloromethyl methyl ether (0.07 ml, 0.92 mmol) in CH₂Cl₂ (2 ml) was stirred at room temperature for 5 h and then diluted with CH₂Cl₂ (20 ml). The whole was washed successively with water, 10% HCl, water, and satd. NaCl, then dried over MgSO₄. Concentration and purification by silica gel column chromatography (ether) afforded (±)-5c (117 mg, 83%) as a colorless oil. IR $\nu_{\text{max}}^{\text{neat}}$ cm^{–1}: 1755 (C=O), 1640 (C=C). ¹H-NMR (CDCl₃) δ: 2.12 (3H, s, CH₃C), 3.34 (3H, s, CH₃O), 3.75 (1H, dd, J=5, 11 Hz, OCH₂CHOCO), 3.90 (1H, dd, J=4, 11 Hz, OCH₂CHOCO), 4.60 (2H, s, OCH₂O), 4.82–4.98 (1H, m), 5.80–5.86 (1H, m, CH₃C=CHCO). MS *m/z*: Calcd for C₈H₁₂O₄ (M⁺): 172.0736. Found: 172.0738.

(±)-4-Isobutyryloxymethyl-3-methyl-2-buten-4-olide (5d) A solution of (±)-5b (201 mg, 1.57 mmol), isobutyryl chloride (0.25 ml, 2.4 mmol), and pyridine (0.2 ml, 2.5 mmol) in CH₂Cl₂ (5 ml) was stirred at room temperature for 23.5 h. After addition of water (10 ml), the organic layer was washed with 10% HCl, water, satd. NaHCO₃, and satd. NaCl successively, then dried over MgSO₄. Concentration and purification by silica gel column chromatography (ether) afforded (±)-5d as a pale yellow oil (300 mg, 97%). IR $\nu_{\text{max}}^{\text{neat}}$ cm^{–1}: 1760 (lactone C=O), 1740 (ester C=O), 1645 (C=C). ¹H-NMR (CDCl₃) δ: 1.15 (6H, d, J=7 Hz, (CH₃)₂CH), 2.10 (3H, s, CH₃C), 2.52 (1H, septet, J=7 Hz, (CH₃)₂CH), 4.28 (1H, dd, J=5, 12 Hz, OCH₂CHOCO), 4.44 (1H, dd, J=4, 12 Hz, OCH₂CHOCO), 4.89–5.04 (1H, m, OCH₂CHOCO), 5.80–5.90 (1H, m, CH₃C=CHCO). MS *m/z*: Calcd for C₁₀H₁₄O₄ (M⁺): 198.0890. Found: 198.0870.

(±)-3-Methyl-4-pivaloyloxymethyl-2-buten-4-olide (5e) Prepared according to the procedure described for (–)-5e, as a pale yellow oil. IR $\nu_{\text{max}}^{\text{neat}}$ cm^{–1}: 1760 (lactone C=O), 1728 (ester C=O), 1645 (C=C). ¹H-NMR (CDCl₃) and MS spectra were identical with those of (–)-5e.

(±)-4-tert-Butyldimethylsilyloxymethyl-3-methyl-2-buten-4-olide (5f) A solution of (±)-5b (105 mg, 0.82 mmol), imidazole (168 mg, 2.47 mmol), and tert-butyldimethylsilyl chloride (177 mg, 1.18 mmol) in dimethylformamide (DMF) (2 ml) was stirred at room temperature for 6 h. After addition of benzene (20 ml), the whole was washed with water, and satd. NaCl, and then dried over MgSO₄. Concentration and purification by silica gel column chromatography (ether) afforded (±)-5f (196 mg, 99%) as a colorless oil. IR $\nu_{\text{max}}^{\text{neat}}$ cm^{–1}: 1760 (C=O), 1645 (C=C). ¹H-NMR (CDCl₃) δ: 0.08 (6H, s, (CH₃)₂Si), 0.88 (9H, s, (CH₃)₃CSi), 2.11 (3H, s, CH₃C), 3.91 (2H, d, J=4, OCH₂CHOCO), 4.72–4.84 (1H, m, OCH₂CHOCO), 5.72–5.82 (1H, m, CH₃C=CHCO). MS *m/z*: Calcd for C₁₂H₂₂O₃Si (M⁺): 242.1338. Found: 242.1395.

(±)-4-(2-Hydroxyprop-2-yl)-3-methyl-2-buten-4-olide (5g) *n*-Butyllithium (*n*-BuLi) (1.43 M in hexane solution, 43.8 ml, 63 mmol) was added to a solution of diisopropylamine (9 ml, 62 mmol) in THF (60 ml) at –78 °C under Ar, then AcOEt (6.1 ml, 62 mmol) was added at –78 °C, and the whole was stirred at –78 °C for 15 min. 3,4-Epoxy-4-methylpentan-2-one¹⁷ (7.15 ml, 63 mmol) was added and the whole was stirred at –78 °C for 30 min, and then satd. NH₄Cl was added. After addition of satd. NaCl, the resultant mixture was extracted with AcOEt (50 ml × 3). The organic layer was washed with satd. NaCl and dried over MgSO₄. After concentration, the residue was dissolved in benzene, and the insoluble material was removed by filtration. The filtrate was again concentrated to give (±)-ethyl 4,5-epoxy-3-hydroxy-3,5-dimethylhexanoate (11.1 g, 88%) as a colorless oil. This was used directly without further purification. IR $\nu_{\text{max}}^{\text{neat}}$ cm^{–1}: 3480 (OH), 1730 (C=O). ¹H-NMR (CDCl₃) δ: 1.29 (3H, t, J=8 Hz, CH₂CH₃), 1.33 (3H, s, (CH₃)₂C), 1.38 (3H, s, (CH₃)₂C), 1.54 (3H, s, CH₃COH), 2.63 (2H, s, CH₂CO), 2.80 (1H, s, C(O)CH), 3.35 (1H, s, OH), 4.24 (2H, q, J=8 Hz, CH₂CH₃). MS *m/z*: 202 (M⁺). A solution of the epoxy ester prepared above (9.91 g) and 10% H₂SO₄ (7 ml) in acetone (70 ml) was stirred at room temperature for 2 h, and NaHCO₃ (solid) was added to the reaction mixture, then the whole was stirred vigorously. The insoluble material was filtered off, and the filtrate was concentrated *in vacuo*. Crystallization of the residue from AcOEt afforded (±)-3-hydroxy-4-(2-hydroxyprop-2-yl)-3-methylbutan-4-olide (3.4 g, 40%) as colorless needles of mp 125.5–127 °C. IR $\nu_{\text{max}}^{\text{KBr}}$ cm^{–1}: 3350 (OH), 3200 (OH), 1780 (C=O). ¹H-NMR (CDCl₃) δ: 1.40 (3H, s, (CH₃)₂COH), 1.53 (3H, s, (CH₃)₂COH), 1.44 (3H, s, CH₃COH), 2.60 (1H, d, J=18 Hz, CH₂CO), 2.70 (1H, d, J=18 Hz, CH₂CO), 3.99 (1H, s, CHOCO). MS *m/z*: 156 (M⁺ – H₂O). Anal. Calcd for C₈H₁₄O₄: C, 55.16; H, 8.10. Found: C, 55.07; H, 8.07. A solution of the lactone prepared above (1.84 g, 10.6 mmol) and diazabicycloundecene (DBU) (1.7 ml, 11.3 mmol) in THF (10 ml) was heated under reflux for 5.3 h and cooled. Concentration and purification of the residue by column chromatography (silica gel, benzene–acetone 9:1) afforded (±)-5g (525 mg, 32%) as a colorless oil. IR $\nu_{\text{max}}^{\text{neat}}$ cm^{–1}: 3440 (OH), 1740 (C=O), 1635 (C=C). ¹H-NMR (CDCl₃) δ: 1.10 (3H, s, (CH₃)₂COH), 1.38 (3H, s, (CH₃)₂COH), 2.20 (3H, s, CH₃C=C), 3.25 (1H, br, OH), 4.69 (1H, s, CHOCO), 5.78–5.92 (1H, m, CH₂C=CHCO). MS *m/z*: Calcd for C₈H₁₁O₃ (M⁺ – H): 155.0706. Found: 155.0699.

General Procedure for the Photocycloaddition of Butenolides with Cyclopentene A 0.1 M solution of butenolide in acetonitrile containing 10 eq of cyclopentene was internally irradiated with a 10 W low pressure mercury lamp at 15 °C for 9 h. Concentration and purification of the residue as follows gave the purified adducts (Table I).

(±)-(1R*,2S*,5R*,6R*,7S*)-6-Methyl-5-trityloxymethyl-4-oxatricyclo[5.3.0.0^{2,6}]decan-3-one (6a) and (±)-(1R*,2S*,5S*,6R*,7S*)-6-Methyl-5-trityloxymethyl-4-oxatricyclo[5.3.0.0^{2,6}]decan-3-one (7a) Purification by column chromatography (silica gel, AcOEt–hexane 1:5) gave a mixture of (±)-6a and (±)-7a as an oil and recovered (±)-5a as a solid. Mixture of (±)-6a and (±)-7a: IR $\nu_{\text{max}}^{\text{neat}}$ cm^{–1}: 1770 (C=O), 1600 (benzene ring). ¹H-NMR (CDCl₃) δ: 0.76 (1.8H, s, CH₃C), 1.00 (1.2H, s, CH₃C), 1.05–1.86 (6H, m, (CH₂)₃), 2.11 (0.4H, d, J=3 Hz, CHCO of 6a), 2.22 (0.6H, d, J=4 Hz, CHCO of 7a), 2.30–2.60 (1H, m), 2.64–2.92 (1H, m), 3.03 (0.6H, dd, J=4, 10 Hz, OCH₂CHOCO of 6a), 3.20 (0.4H, dd, J=6, 10 Hz, OCH₂CHOCO of 7a), 3.29 (0.6H, dd, J=4, 10 Hz, OCH₂CHOCO of 6a), 3.60 (0.4H, dd, J=6, 10 Hz, OCH₂CHOCO of 7a), 4.26 (0.4H, t, J=6 Hz, OCH₂CHOCO of 7a), 4.43 (0.6H, t, J=4 Hz, OCH₂CHOCO of 6a), 6.90–7.56 (15H, m, PhH). MS *m/z*: 438 (M⁺). Crystallization of the mixture of (±)-6a and (±)-7a from AcOEt–hexane gave (±)-6a as colorless prisms of mp 162–163 °C. IR $\nu_{\text{max}}^{\text{KBr}}$ cm^{–1}: 1760 (C=O). ¹H-NMR (CDCl₃) δ: 0.84 (3H, s, CH₃C), 1.16–1.84 (6H, m, (CH₂)₃), 2.30 (1H, d, J=4 Hz, CHCO), 2.58 (1H, t, J=7 Hz), 2.80–3.00 (1H, m), 3.13 (1H, dd, J=4, 10 Hz, OCH₂CHOCO), 3.33 (1H, dd, J=4, 10 Hz, OCH₂CHOCO), 4.57 (1H, t, J=4 Hz, OCH₂CHOCO), 7.00–7.56 (15H, m, benzene ring). MS *m/z*: 438 (M⁺). Anal. Calcd for C₃₀H₃₀O₃: C, 82.16; H, 6.90. Found: C, 81.90; H, 7.15.

(±)-(1R*,2S*,5R*,6R*,7S*)-5-Hydroxymethyl-6-methyl-4-oxatricyclo[5.3.0.0^{2,6}]decan-3-one (6b) and (±)-(1R*,2S*,5S*,6R*,7S*)-5-Hydroxymethyl-6-methyl-4-oxatricyclo[5.3.0.0^{2,6}]decan-3-one (7b) Purification by column chromatography (silica gel, AcOEt–hexane 1:2) gave a mixture of (±)-6b and (±)-7b as a colorless oil. IR $\nu_{\text{max}}^{\text{neat}}$ cm^{–1}: 3400 (OH), 1760 (C=O). ¹H-NMR (CDCl₃) δ: 1.05 (1.7H, s, CH₃C of 6b), 1.12 (1.3H, s, CH₃C of 7b), 1.16–2.00 (6H, m, (CH₂)₃), 2.04–2.28 (1H, m), 2.45–3.00 (2H, m), 3.32 (1H, br), 3.64–4.13 (2H, m), 4.31 (0.44H, dd, J=4, 7 Hz, OCH₂CHOCO of 6b), 4.58 (0.56H, t, J=5 Hz, OCH₂CHOCO of 7b). MS *m/z*: 196 (M⁺).

($\pm$)-(1*R**,2*S**,5*R**,6*R**,7*S**)-5-Methoxymethoxymethyl-6-methyl-4-oxatricyclo[5.3.0.0^{2,6}]decan-3-one (6c) and ($\pm$)-(1*R**,2*S**,5*S**,6*R**,7*S**)-5-Methoxymethoxymethyl-6-methyl-4-oxatricyclo[5.3.0.0^{2,6}]decan-3-one (7c) Purification by column chromatography (silica gel, AcOEt-hexane 1:3) gave a mixture of ($\pm$)-6c and ($\pm$)-7c as a colorless oil. IR $\nu_{\text{max}}^{\text{neat}}$ cm⁻¹: 1770 (C=O). ¹H-NMR (CDCl₃) δ : 1.06 (1.5H, s, CH₃C), 1.12 (1.5H, s, CH₃C), 1.28–2.10 (6H, m, (CH₂)₃), 2.18–2.30 (1H, m), 2.52–3.04 (2H, m), 3.36 (1.5H, s, OCH₃), 3.39 (1.5H, s, CH₃O), 3.64–3.96 (2H, m), 4.33 (1H, t, *J*=6 Hz, OCH₂CHOCO), 4.61 (1H, s, OCH₂O), 4.66 (1H, s, OCH₂O). MS *m/z*: Calcd for C₁₃H₂₀O₄Si (M⁺): 240.1361. Found: 240.1371.

($\pm$)-(1*R**,2*S**,5*R**,6*R**,7*S**)-5-Isobutyryloxymethyl-6-methyl-4-oxatricyclo[5.3.0.0^{2,6}]decan-3-one (6d) and ($\pm$)-(1*R**,2*S**,5*S**,6*R**,7*S**)-5-Isobutyryloxymethyl-6-methyl-4-oxatricyclo[5.3.0.0^{2,6}]decan-3-one (7d) Purification by column chromatography (silica gel, ether-hexane 1:2) gave ($\pm$)-6d and ($\pm$)-7d as colorless oils.

($\pm$)-6d: IR $\nu_{\text{max}}^{\text{neat}}$ cm⁻¹: 1770 (lactone C=O), 1735 (ester C=O). ¹H-NMR (CDCl₃) δ : 1.04 (3H, s, CH₃C), 1.16 (6H, d, *J*=7 Hz, (CH₃)₂CH), 1.28–2.02 (6H, m, (CH₂)₃), 2.25 (1H, d, *J*=4 Hz, CHCO of lactone), 2.32–2.80 (2H, m), 2.80–3.04 (1H, m), 4.11 (1H, dd, *J*=6, 12 Hz, OCH₂CHOCO), 4.33 (1H, dd, *J*=4, 12 Hz, OCH₂CHOCO), 4.67 (1H, dd, *J*=4, 6 Hz, OCH₂CHOCO). MS *m/z*: Calcd for C₁₅H₂₂O₄Si (M⁺): 266.1516. Found: 266.1498.

($\pm$)-7d: IR $\nu_{\text{max}}^{\text{neat}}$ cm⁻¹: 1775 (lactone C=O), 1740 (ester C=O). ¹H-NMR (CDCl₃) δ : 1.14 (3H, s, CH₃C), 1.18 (6H, d, *J*=7 Hz, (CH₃)₂CH), 1.28–1.98 (6H, m, (CH₂)₃), 2.29 (1H, d, *J*=3 Hz, CHCO of lactone), 2.40–2.82 (3H, m), 4.36 (3H, s, OCH₂CHOCO). MS *m/z*: Calcd for C₁₅H₂₂O₄Si (M⁺): 266.1515. Found: 266.1509.

($\pm$)-(1*R**,2*S**,5*R**,6*R**,7*S**)-6-Methyl-5-pivaloyloxymethyl-4-oxatricyclo[5.3.0.0^{2,6}]decan-3-one (6e) and ($\pm$)-(1*R**,2*S**,5*S**,6*R**,7*S**)-6-Methyl-5-pivaloyloxymethyl-4-oxatricyclo[5.3.0.0^{2,6}]decan-3-one (7e) Purification by column chromatography (silica gel, ether-hexane 1:2) gave ($\pm$)-6e and ($\pm$)-7e as colorless oils.

($\pm$)-6e: IR $\nu_{\text{max}}^{\text{neat}}$ cm⁻¹: 1765 (lactone C=O), 1730 (ester C=O). ¹H-NMR (CDCl₃) δ : 1.04 (3H, s, CH₃C), 1.20 (9H, s, (CH₃)₃CCO), 1.40–2.00 (6H, m, (CH₂)₃), 2.26 (1H, d, *J*=4 Hz, OCH₂CHCO), 2.52–2.80 (1H, m), 2.80–3.00 (1H, m), 4.05 (1H, dd, *J*=4, 12 Hz, OCH₂CHOCO), 4.32 (1H, dd, *J*=4, 12 Hz, OCH₂CHOCO), 4.65 (1H, t, *J*=4 Hz, CH₂CHOCO). MS *m/z*: Calcd for C₁₆H₂₄O₄ (M⁺): 280.1674. Found: 280.1710.

($\pm$)-7e: IR $\nu_{\text{max}}^{\text{neat}}$ cm⁻¹: 1770 (lactone C=O), 1725 (ester C=O). ¹H-NMR (CDCl₃) δ : 1.14 (3H, s, CH₃C), 1.22 (9H, s, (CH₃)₃CCO), 1.40–1.96 (6H, m, (CH₂)₃), 2.28 (1H, d, *J*=3 Hz, OCH₂CHCO), 2.68–2.80 (2H, m), 4.33 (3H, s, OCH₂CHOCO). MS *m/z*: Calcd for C₁₆H₂₄O₄ (M⁺): 281.1751. Found: 281.1724.

($\pm$)-(1*R**,2*S**,5*R**,6*R**,7*S**)-5-*tert*-Butyldimethylsilyloxymethyl-6-methyl-4-oxatricyclo[5.3.0.0^{2,6}]decan-3-one (6f) and ($\pm$)-(1*R**,2*S**,5*S**,6*R**,7*S**)-5-*tert*-Butyldimethylsilyloxymethyl-6-methyl-4-oxatricyclo[5.3.0.0^{2,6}]decan-3-one (7f) Purification by column chromatography (silica gel, ether-benzene 1:20) gave ($\pm$)-6f and ($\pm$)-7f as colorless solids.

($\pm$)-6f: mp 58–60°C (colorless prisms from hexane). IR $\nu_{\text{max}}^{\text{KBr}}$ cm⁻¹: 1750 (C=O). ¹H-NMR (CDCl₃) δ : 0.06 (6H, s, (CH₃)₂Si), 0.87 (9H, s, (CH₃)₃CSi), 1.13 (3H, s, CH₃C), 1.32–1.96 (6H, m, (CH₂)₃), 2.24 (1H, d, *J*=4 Hz, OCH₂CHCO), 2.58–2.94 (2H, m), 3.78 (1H, dd, *J*=7, 10 Hz, OCH₂CHOCO), 3.97 (1H, dd, *J*=5, 10 Hz, OCH₂CHOCO), 4.21 (1H, dd, *J*=5, 7 Hz, OCH₂CHOCO). MS *m/z*: 295 (M⁺–CH₃). Anal. Calcd for C₁₇H₃₀O₃Si: C, 65.79; H, 9.74. Found: C, 65.84; H, 9.90.

($\pm$)-7f: mp 71.5–73°C (colorless needles from hexane). IR $\nu_{\text{max}}^{\text{KBr}}$ cm⁻¹: 1755 (C=O). ¹H-NMR (CDCl₃) δ : 0.07 (6H, s, (CH₃)₂Si), 0.87 (9H, s, (CH₃)₃CSi), 1.13 (3H, s, CH₃C), 1.28–1.97 (6H, m, (CH₂)₃), 2.22 (1H, d, *J*=3 Hz, OCH₂CHCO), 2.65 (1H, t, *J*=7 Hz), 2.74–2.97 (1H, m), 3.78 (2H, d, *J*=4 Hz, OCH₂CHOCO), 4.42 (1H, t, *J*=4 Hz, OCH₂CHOCO). MS *m/z*: 311 (M⁺+1). Anal. Calcd for C₁₇H₃₀O₃Si: C, 65.79; H, 9.74. Found: C, 65.62; H, 10.03.

($\pm$)-(1*R**,2*S**,5*R**,6*R**,7*S**)-5-(2-Hydroxyprop-2-yl)-6-methyl-4-oxatricyclo[5.3.0.0^{2,6}]decan-3-one (6g) and ($\pm$)-(1*R**,2*S**,5*S**,6*R**,7*S**)-5-(2-Hydroxyprop-2-yl)-6-methyl-4-oxatricyclo[5.3.0.0^{2,6}]decan-3-one (7g) Purification by column chromatography (silica gel, acetone-benzene 1:19) gave ($\pm$)-6g and ($\pm$)-7g as a colorless solid and a colorless oil, respectively.

($\pm$)-6g: mp 99.5–100°C (colorless needles from AcOEt-hexane). IR $\nu_{\text{max}}^{\text{KBr}}$ cm⁻¹: 3480 (OH), 1760 (C=O). ¹H-NMR (CDCl₃) δ : 1.20 (3H, s, CH₃C), 1.32 (3H, s, CH₃COH), 1.39 (3H, s, CH₃COH), 1.44–2.00 (6H, m, (CH₂)₃), 2.10 (1H, br, OH), 2.18 (1H, d, *J*=5 Hz, CHCO), 2.53 (1H, t, *J*=5 Hz), 2.92–3.17 (1H, m, CHCHCO), 4.39 (1H, s, CHOCO). MS *m/z*:

207 (M⁺–OH). Anal. Calcd for C₁₃H₂₀O₃: C, 68.61; H, 8.99. Found: C, 68.89; H, 8.98.

($\pm$)-7g: IR $\nu_{\text{max}}^{\text{neat}}$ cm⁻¹: 3460 (OH), 1755 (C=O). ¹H-NMR (CDCl₃) δ : 1.14 (3H, s, CH₃C), 1.25 (3H, s, CH₃COH), 1.39 (3H, s, CH₃COH), 1.44–2.06 (6H, m, (CH₂)₃), 2.26 (1H, br, OH), 2.26 (1H, d, *J*=4 Hz, CHCO), 2.62–2.86 (1H, m), 3.80 (1H, t, *J*=8 Hz), 3.96 (1H, s, CHOCO). MS *m/z*: Calcd for C₁₃H₁₉O₂ (M⁺–OH): 207.1395. Found: 207.1402.

(–)-4-Hydroxymethyl-3-methyl-2-buten-4-olide (5b) According to the procedure described for ($\pm$)-5b, (–)-5b was prepared from (–)-5a^{10a} (86%) as colorless needles of mp 60–62.5°C (from benzene). [α]_D²⁰ –27.4° (*c*=1.05, CHCl₃). IR $\nu_{\text{max}}^{\text{KBr}}$ cm⁻¹: 3400 (OH), 1720 (C=O), 1635 (C=C). ¹H-NMR (CDCl₃) δ : 2.07 (3H, s, CH₃C), 3.32 (1H, br, OH), 3.68 (1H, dd, *J*=4, 13 Hz, OCH₂CHOCO), 3.94 (1H, dd, *J*=3, 13 Hz, OCH₂CHOCO), 4.77 (1H, br, OCH₂CHOCO), 5.60–5.76 (1H, m, CH₃C=CHCO). MS *m/z*: 129 (M⁺+1). Anal. Calcd for C₆H₈O₃: C, 56.24; H, 6.29. Found: C, 55.97; H, 6.21.

(–)-3-Methyl-4-pivaloyloxymethyl-2-buten-4-olide (5e) A solution of (–)-5b (3.3 g, 26 mmol), pivaloyl chloride (4.5 ml, 37 mmol), pyridine (3 ml, 37 mmol), and 4-dimethylaminopyridine (DMAP) (0.5 g, 4 mmol) in CH₂Cl₂ (60 ml) was stirred at room temperature for 23 h. After addition of water (10 ml), the organic layer was washed with 10% HCl, water, satd. NaHCO₃, and satd. NaCl, successively, then dried over MgSO₄. Concentration afforded (–)-5e (3.95 g, 72%) as colorless needles of mp 65–67°C (from AcOEt-hexane). [α]_D²⁰ –69.1° (*c*=0.90, CHCl₃). IR $\nu_{\text{max}}^{\text{KBr}}$ cm⁻¹: 1760 (lactone C=O), 1720 (ester C=O), 1640 (C=C). ¹H-NMR (CDCl₃) δ : 1.18 (9H, s, (CH₃)₃CO), 2.12 (3H, s, CH₃C), 4.36 (1H, dd, *J*=4, 12 Hz, OCH₂CHOCO), 4.43 (1H, dd, *J*=3, 12 Hz, OCH₂CHOCO), 5.01 (1H, br, OCH₂CHOCO), 5.82–5.96 (1H, m, CH₃C=CHCO). MS *m/z*: 212 (M⁺). Anal. Calcd for C₁₁H₁₆O₄: C, 62.25; H, 7.60. Found: C, 62.26; H, 7.60.

Cycloaddition of (–)-5e with 15 A solution of (–)-5e (1.74 g, 8.2 mmol) and 15 (90 ml, 80 mmol) in acetonitrile was internally irradiated with a 30 W low pressure mercury lamp at 0°C for 92 h under Ar. After concentration, the residual oil was separated by medium pressure column chromatography (silica gel, AcOEt-hexane 1:4). The first fraction gave (+)-19 (0.27 g, 9.7%) as colorless prisms of mp 74.5–76°C (from AcOEt-hexane). [α]_D²⁰ +70.8° (*c*=1.13, CHCl₃). IR $\nu_{\text{max}}^{\text{KBr}}$ cm⁻¹: 1770 (lactone C=O), 1720 (ester C=O). ¹H-NMR (CDCl₃) δ : 1.23 (9H, s, (CH₃)₃CCO), 1.39 (3H, s, CH₃C), 1.72–2.14 (4H, m, (CH₂)₂), 2.40–2.54 (2H, m), 2.60–2.84 (1H, m), 3.78–3.94 (4H, m, O(CH₂)₂O), 4.34 (3H, s, OCH₂CHOCO). MS *m/z*: 338 (M⁺). Anal. Calcd for C₁₈H₂₆O₆: C, 63.89; H, 7.74. Found: C, 64.17; H, 7.84. The second fraction gave (+)-18 (0.29 g, 10%) as colorless needles of mp 110–112°C (from AcOEt-hexane). [α]_D²⁰ +91.2° (*c*=0.50, CHCl₃). IR $\nu_{\text{max}}^{\text{KBr}}$ cm⁻¹: 1765 (lactone C=O), 1720 (ester C=O). ¹H-NMR (CDCl₃) δ : 1.21 (9H, s, (CH₃)₃CCO), 1.24 (3H, s, CH₃C), 1.68–2.09 (4H, m, (CH₂)₂), 2.47 (1H, dd, *J*=4, 8 Hz, CHCHCO), 2.75 (1H, d, *J*=4 Hz, CHCO), 2.68–2.86 (1H, m), 3.80–4.02 (4H, m, O(CH₂)₂O), 4.34 (3H, s, OCH₂CHOCO). MS *m/z*: 338 (M⁺). Anal. Calcd for C₁₈H₂₆O₆: C, 63.89; H, 7.74. Found: C, 63.69; H, 7.80. The third fraction gave a mixture of (–)-16 and 17 as a colorless oil (1.67 g). Crystallization of this oil from AcOEt-hexane gave (–)-16 as colorless needles (0.68 g, 25%) of mp 126–128°C. [α]_D²⁰ –22.5° (*c*=0.32, CHCl₃). IR $\nu_{\text{max}}^{\text{KBr}}$ cm⁻¹: 1775 (lactone C=O), 1725 (ester C=O). ¹H-NMR (CDCl₃) δ : 1.14 (3H, s, CH₃C), 1.20 (9H, s, (CH₃)₃CCO), 1.66–2.12 (4H, m, (CH₂)₂), 2.52–2.76 (3H, m), 3.76–3.98 (4H, m, O(CH₂)₂O), 4.08 (1H, dd, *J*=4, 12 Hz, OCH₂CHOCO), 4.32 (1H, dd, *J*=4, 12 Hz, OCH₂CHOCO), 4.61 (1H, t, *J*=4 Hz, OCH₂CHOCO). MS *m/z*: 338 (M⁺). Anal. Calcd for C₁₈H₂₆O₆·1/4H₂O: C, 63.00; H, 7.79. Found: C, 62.98; H, 7.74. The mother liquor (0.74 g) of the third fraction contained (–)-16 and 17, but all attempts to separate these compounds were unsuccessful and this mixture was used without further purification.

(–)-(1*S*,2*R*,5*S*,6*S*,7*R*)-10-Ethylenedioxy-3,6-dimethyl-5-pivaloyloxymethyl-4-oxatricyclo[5.3.0.0^{2,6}]decan-3-ol (20) A stirred solution of (–)-16 (594 mg, 1.75 mmol) in THF (35 ml) was treated with MeLi (1.16 M in ether, 2 ml, 2.32 mmol) at –78°C under Ar, and the whole mixture was stirred at –78°C for 50 min, then quenched with satd. NH₄Cl (30 ml). The resultant mixture was extracted with CH₂Cl₂ (50 ml × 3), and the organic layers were combined, washed with satd. NaCl and dried over MgSO₄. After evaporation of the solvents, the residue was chromatographed (silica gel, benzene-acetone 40:1) to afford epimeric mixture of (–)-20 (593 mg, 95%) as colorless needles of mp 94.5–95°C (from AcOEt-hexane). [α]_D²⁰ –21.0° (*c*=1.37, CHCl₃). IR $\nu_{\text{max}}^{\text{KBr}}$ cm⁻¹: 3440 (OH), 1720 (ester C=O). ¹H-NMR (CDCl₃) δ : 1.08–1.32 (12H, m), 1.42 (1.5H, s, OC(OH)CH₃), 1.60 (1.5H, s, OC(OH)CH₃), 1.68–2.96 (4H, m), 2.11–2.48 (3H, m), 2.69–2.83 (0.5H, m), 3.06 (0.5H, s, OH), 3.64–4.00 (5.5H, m), 4.04 (1H, dd, *J*=4, 9 Hz, OCH₂CHOCO), 4.74 (0.5H, dd, *J*=11, 9 Hz,

CHOC(OH)CH_3). MS m/z : 354 (M^+). Anal. Calcd for $\text{C}_{19}\text{H}_{30}\text{O}_6$: C, 64.38; H, 8.53. Found: C, 64.42; H, 8.68.

(+)-(1*S*,2*R*,5*S*,6*S*,7*R*)-8-Ethylenedioxy-3,6-dimethyl-5-pivaloyloxymethyl-4-oxatricyclo[5.3.0.0^{2,6}]decan-3-ol (23) and (-)-(1*S*,2*R*,5*S*,6*S*,7*R*)-10-Ethylenedioxy-3,6-dimethyl-5-pivaloyloxymethyl-4-oxatricyclo[5.3.0.0^{2,6}]decan-3-ol (20) A stirred solution of (-)-16 and 17 (obtained as the mother liquor after crystallization of pure (-)-16, 737 mg) in THF (50 ml) was treated with MeLi (1.16 M in ether, 2.1 ml, 2.4 mmol) at -78 °C under Ar, and the whole mixture was stirred at -78 °C for 1 h, then treated as described above. The resultant residue was separated by chromatography (silica gel, benzene-acetone 10:1). The first fraction gave (+)-23 (303 mg, 10% from (-)-5e) as an oil. $[\alpha]_D^{20} + 8.82^\circ$ ($c = 1.52$, CHCl_3). IR $\nu_{\text{max}}^{\text{CHCl}_3} \text{cm}^{-1}$: 3440 (OH), 1720 (ester C=O). $^1\text{H-NMR}$ (CDCl_3) δ : 1.12–1.32 (12H, m), 1.36–1.54 (3H, m), 2.68–2.20 (6H, m), 2.32–2.65 (1H, m), 2.88–3.20 (1H, m), 3.64–4.24 (6.5H, m), 4.73 (0.5H, dd, $J = 9, 11$ Hz, CHOC(OH)CH_3). MS m/z : Calcd for $\text{C}_{19}\text{H}_{30}\text{O}_6$ (M^+): 354.2039. Found: 354.2017. The second fraction gave (-)-20 (119 mg, 4.1% from (-)-5e) as colorless needles.

(-)-(1*S*,2*R*,6*S*,7*R*)-10-Ethylenedioxy-6-methyltricyclo[5.3.0.0^{2,6}]decan-3-one (22) Sodium methoxide (3.5 M in MeOH, 0.8 ml, 2.8 mmol) was added to a solution of (-)-20 (593 mg, 1.68 mmol) in MeOH (40 ml), and the whole was heated under reflux for 30 min and then cooled to room temperature. After addition of water (10 ml) and concentration, the residue was extracted with CH_2Cl_2 (30 ml $\times$ 5), and the organic layers were combined and dried over K_2CO_3 . Concentration left the crude diol (0.4 g, quant.) as an oil. This oil was dissolved in AcOEt (50 ml), a solution of NaIO_4 (0.89 g, 4.2 mmol) in water (35 ml) was added, and the whole was stirred vigorously at room temperature for 1.2 h. AcOEt (100 ml) was added, and the organic layer of the resultant mixture was washed successively with 10% $\text{Na}_2\text{S}_2\text{O}_3$ and satd. NaCl, then dried over MgSO_4 . Evaporation of the solvent afforded 21 (338 mg) as an oil. IR $\nu_{\text{max}}^{\text{CHCl}_3} \text{cm}^{-1}$: 2620 (C–H of aldehyde), 1720 (C=O of aldehyde and ketone). $^1\text{H-NMR}$ (CDCl_3) δ : 1.29 (3H, s, CH_3C), 1.72–1.99 (4H, m), 2.07 (3H, s, CH_3CO), 2.58–2.92 (3H, m), 3.60–3.96 (4H, m, $\text{O}(\text{CH}_2)_2\text{O}$), 9.65 (1H, s, CHO). MS m/z : 238 (M^+). A solution of 1 N NaOH (2 ml, 2 mmol) was added to a solution of crude 21 (338 mg) in ether (10 ml) and the whole was stirred at room temperature for 1.2 h. After dilution with CH_2Cl_2 (20 ml), the resultant mixture was washed with water (10 ml) and dried over MgSO_4 . Evaporation of the solvent and purification by column chromatography (silica gel, AcOEt-hexane 1:2) afforded (-)-22 (271 mg, 74% from (-)-20) as colorless prisms of mp 49–50.5 °C (from AcOEt-hexane). $[\alpha]_D^{20} - 179^\circ$ ($c = 0.40$, CHCl_3). IR $\nu_{\text{max}}^{\text{KBr}} \text{cm}^{-1}$: 1695 (C=O), 1570 (C=C). $^1\text{H-NMR}$ (CDCl_3) δ : 1.16 (3H, s, CH_3C), 1.68–2.16 (5H, m), 2.40–2.64 (2H, m), 3.84–4.00 (4H, m, $\text{O}(\text{CH}_2)_2\text{O}$), 6.18 (1H, d, $J = 5$ Hz, $\text{CH}=\text{CHCO}$), 7.51 (1H, d, $J = 5$ Hz, $\text{CH}=\text{CHCO}$). MS m/z : 220 (M^+). Anal. Calcd for $\text{C}_{13}\text{H}_{16}\text{O}_3$: C, 70.89; H, 7.32. Found: C, 70.98; H, 7.41.

(-)-(1*R*,2*R*,6*R*,7*R*)-8-Ethylenedioxy-6-methyltricyclo[5.3.0.0^{2,6}]decan-3-one (24) According to the procedure described for (-)-22, (-)-24 was prepared from 23 as a colorless oil (52%) of $[\alpha]_D^{20} - 181^\circ$ ($c = 1.18$, CHCl_3). IR $\nu_{\text{max}}^{\text{CHCl}_3} \text{cm}^{-1}$: 1690 (C=O), 1580 (C=C). $^1\text{H-NMR}$ (CDCl_3) δ : 1.28 (3H, s, CH_3C), 1.68–2.06 (4H, m, $\text{CCCH}_2\text{CH}_2\text{C}$), 2.08–2.28 (2H, m), 2.32–2.56 (1H, m), 3.70–4.12 (4H, m, $\text{O}(\text{CH}_2)_2\text{O}$), 6.09 (1H, d, $J = 5$ Hz, $\text{CH}=\text{CHCO}$), 7.47 (1H, d, $J = 5$ Hz, $\text{CH}=\text{CHCO}$). MS m/z : Calcd for $\text{C}_{13}\text{H}_{16}\text{O}_3$ (M^+): 220.1096. Found: 220.1089.

(+)-(1*R*,2*S*,5*S*,6*R*,7*S*)-10-Ethylenedioxy-3,6-dimethyl-5-pivaloyloxymethyl-4-oxatricyclo[5.3.0.0^{2,6}]decan-3-ol (25) According to the procedure described for (-)-20, (-)-25 was prepared from (+)-18 as colorless needles (93%) of mp 133.5–135.5 °C (from benzene-hexane). $[\alpha]_D^{20} + 50.9^\circ$ ($c = 1.12$, CHCl_3). IR $\nu_{\text{max}}^{\text{KBr}} \text{cm}^{-1}$: 3440 (OH), 1725 (C=O). $^1\text{H-NMR}$ (CDCl_3) δ : 1.18 (3H, s, CH_3C), 1.20 (9H, s, $(\text{CH}_3)_3\text{CO}$), 1.48 (3H, s, $\text{CH}_3\text{C}(\text{OH})\text{O}$), 1.52–2.25 (7H, m), 2.31–2.52 (1H, m), 3.63–3.98 (4H, m, $\text{O}(\text{CH}_2)_2\text{O}$), 4.05–4.29 (3H, m, OCH_2CHO). MS m/z : 354 (M^+). Anal. Calcd for $\text{C}_{19}\text{H}_{30}\text{O}_6$: C, 64.38; H, 8.53. Found: C, 64.29; H, 8.45.

(-)-(1*R*,2*S*,6*R*,7*S*)-10-Ethylenedioxy-6-methyltricyclo[5.3.0.0^{2,6}]decan-3-one (26) According to the procedure described for (-)-22, (-)-26 was prepared from (+)-25 as colorless prisms (73%) of mp 50.0–51.0 °C (from AcOEt-hexane). $[\alpha]_D^{20} + 177^\circ$ ($c = 1.02$, CHCl_3). The $^1\text{H-NMR}$ and IR spectra were identical with those of (-)-22.

(+)-(1*R*,2*S*,5*S*,6*R*,7*S*)-8-Ethylenedioxy-3,6-dimethyl-5-pivaloyloxymethyl-4-oxatricyclo[5.3.0.0^{2,6}]decan-3-ol (27) According to the procedure described for (-)-20, (+)-27 was prepared from (+)-19 as a colorless oil (quant.). $[\alpha]_D^{20} + 16.3^\circ$ ($c = 0.87$, CHCl_3). IR $\nu_{\text{max}}^{\text{neat}} \text{cm}^{-1}$: 1730 (C=O). $^1\text{H-NMR}$ (CDCl_3) δ : 1.20 (9H, s, $(\text{CH}_3)_3\text{CO}$), 1.32 (3H, s, CH_3C), 1.48 (3H, s, $\text{CH}_3\text{C}(\text{OH})\text{O}$), 1.58–2.48 (8H, m), 3.60–3.88 (4H, m, $\text{O}(\text{CH}_2)_2\text{O}$), 3.89–4.08 (1H, m, OCH_2CH), 4.14–4.30 (2H, m,

OCH_2CHO). MS m/z : Calcd for $\text{C}_{19}\text{H}_{30}\text{O}_6$ (M^+): 354.2042. Found: 354.2047.

(-)-(1*S*,2*S*,6*S*,7*S*)-8-Ethylenedioxy-6-methyltricyclo[5.3.0.0^{2,6}]decan-3-one (28) According to the procedure described for (-)-22, (-)-28 was prepared from (+)-27 as a colorless oil (58%) of $[\alpha]_D^{20} + 186^\circ$ ($c = 0.88$, CHCl_3). The IR and $^1\text{H-NMR}$ were identical with those of (-)-24.

(-)-(1*S*,2*R*,6*S*,7*R*)-10-Ethylenedioxy-6-methyltricyclo[5.3.0.0^{2,6}]decan-3-one (30) A solution of (-)-22 (271 mg) in AcOEt (30 ml) was hydrogenated over 10% Pd-C (29 mg) under H_2 for 3 h. The catalyst was filtered off and the filtrate was evaporated to give (-)-30 (263 mg, 96%) as colorless needles of mp 95–96 °C (from hexane). $[\alpha]_D^{20} - 176^\circ$ ($c = 0.91$, CHCl_3). IR $\nu_{\text{max}}^{\text{KBr}} \text{cm}^{-1}$: 1720 (C=O). $^1\text{H-NMR}$ (CDCl_3) δ : 1.13 (3H, s, CH_3C), 1.60–2.08 (7H, m), 2.12–2.36 (2H, m), 2.38–2.77 (2H, m), 3.68–4.06 (4H, m, $\text{O}(\text{CH}_2)_2\text{O}$). MS m/z : 222 (M^+). Anal. Calcd for $\text{C}_{13}\text{H}_{18}\text{O}_3$: C, 70.24; H, 8.16. Found: C, 70.04; H, 8.21.

(-)-(1*S*,2*R*,3*S*,6*S*,7*R*)-10-Ethylenedioxy-6-methyltricyclo[5.3.0.0^{2,6}]decan-3-ol (31) NaBH_4 (78 mg, 2 mmol) was added to a solution of (-)-30 (263 mg, 1.18 mmol) in ethanol (EtOH) (10 ml) at -78 °C and the whole was stirred at -78 °C for 2 h and at room temperature for 1.5 h. After evaporation of the solvent, the residue was taken up into satd. NaCl (20 ml), and extracted with CH_2Cl_2 (30 ml $\times$ 3), and combined extract was dried over MgSO_4 . Evaporation of the solvent gave (-)-31 (279 mg, quant.) as a colorless oil. $[\alpha]_D^{20} - 35.0^\circ$ ($c = 1.13$, CHCl_3). IR $\nu_{\text{max}}^{\text{CHCl}_3} \text{cm}^{-1}$: 3450 (OH). $^1\text{H-NMR}$ (CDCl_3) δ : 0.98 (3H, s, CH_3C), 1.03–2.39 (12H, m), 3.62–3.95 (4H, m, $\text{O}(\text{CH}_2)_2\text{O}$), 4.32 (1H, dt, $J = 9, 7$ Hz, CHOH). MS m/z : Calcd for $\text{C}_{13}\text{H}_{20}\text{O}_3$ (M^+): 224.1416. Found: 224.1419.

(-)-(1*S*,2*R*,3*S*,6*S*,7*R*)-10-Ethylenedioxy-6-methyltricyclo[5.3.0.0^{2,6}]decan-3-yl *p*-Toluenesulfonate (32) A solution of (-)-31 (279 mg, 1.18 mmol) and *p*-toluenesulfonyl chloride (1.4 g, 7.3 mmol) in pyridine (10 ml) was stirred at room temperature for 20.5 h. Water (10 ml) was added and resultant mixture was stirred for 30 min, and then extracted with CH_2Cl_2 (30 ml $\times$ 3). The combined extracts were washed with water, 10% HCl, water, and satd. NaHCO_3 , successively, then dried over MgSO_4 . Concentration gave (-)-32 (452 mg, quant.) as colorless needles of mp 71–72.5 °C (from hexane). $[\alpha]_D^{20} - 53.2^\circ$ ($c = 1.17$, CHCl_3). IR $\nu_{\text{max}}^{\text{KBr}} \text{cm}^{-1}$: 1595, 1360, 1185. $^1\text{H-NMR}$ (CDCl_3) δ : 0.96 (3H, s, CH_3C), 1.17–2.30 (11H, m), 2.45 (3H, s, CH_3 -aromatic), 3.89–3.91 (4H, m, $\text{O}(\text{CH}_2)_2\text{O}$), 4.75 (1H, dt, $J = 9, 7$ Hz, CHOSO_2), 7.33 (2H, d, $J = 8$ Hz, aromatic), 7.78 (2H, d, $J = 8$ Hz, aromatic). MS m/z : 223 ($M^+ - \text{CH}_3\text{C}_6\text{H}_4\text{SO}_2$). Anal. Calcd for $\text{C}_{20}\text{H}_{26}\text{O}_5\text{S}$: C, 63.48; H, 6.93. Found: C, 63.40; H, 6.87.

(-)-Diethyl ((1*R*,2*R*,3*R*,6*R*,7*R*)-10-Ethylenedioxy-6-methyltricyclo[5.3.0.0^{2,6}]decan-3-yl)malonate (33) Diethyl malonate (1.2 ml, 7.9 mmol) was added to a suspension of NaH (247 mg, 55% in oil, 5.9 mmol) in dimethoxyethane (DME) (9 ml) at 0 °C. Then a solution of (-)-32 (432 mg, 1.14 mmol) in DME (24 ml) was added, the whole was heated under reflux for 19 h and cooled. The reaction mixture was taken up in 10% HCl, and extracted with CH_2Cl_2 (30 ml $\times$ 3). The combined extracts were washed successively with water and satd. NaHCO_3 , and then dried over MgSO_4 . Evaporation of the solvent and chromatography of the residue (silica gel, benzene-AcOEt 30:1) gave (-)-33 (378 mg, 91%) as a colorless oil. $[\alpha]_D^{20} - 23.8^\circ$ ($c = 1.57$, CHCl_3). IR $\nu_{\text{max}}^{\text{CHCl}_3} \text{cm}^{-1}$: 1740 (C=O), 1725 (C=O). $^1\text{H-NMR}$ (CDCl_3) δ : 1.04 (3H, s, CH_3C), 1.26 (6H, t, $J = 7$ Hz, CH_2CH_3), 1.40–2.33 (11H, m), 3.09 (1H, d, $J = 11$ Hz, $\text{CH}(\text{CO}_2\text{Et})_2$), 2.48 (1H, dd, $J = 6, 11$ Hz, $\text{CHCH}(\text{CO}_2\text{Et})_2$), 3.70–4.00 (4H, m, $\text{O}(\text{CH}_2)_2\text{O}$), 4.18 (2H, q, $J = 7$ Hz, CH_2CH_3), 4.17 (2H, q, $J = 7$ Hz, CH_2CH_3). MS m/z : Calcd for $\text{C}_{20}\text{H}_{30}\text{O}_6$ (M^+): 366.2040. Found: 366.2039.

(-)-(1*R*,2*R*,3*R*,6*R*,7*R*)-3-(1,3-Dihydroxyprop-2-yl)-10-ethylenedioxy-6-methyltricyclo[5.3.0.0^{2,6}]decan-3-ol (34) A solution of (-)-33 (378 mg, 1.03 mmol) in ether (30 ml) was added to a stirred suspension of lithium aluminum hydride (LAH) (0.16 g, 4.2 mmol) in ether (10 ml) at 0 °C, and the whole was stirred at room temperature for 2 h and then recooled to 0 °C. Water (0.32 ml) and 15% NaOH (0.26 ml) were added successively at 0 °C, and the whole was stirred at room temperature for 2 h. After filtration, the filtrate was concentrated to give 34 (248 mg, 85%) as colorless needles of mp 108–108.5 °C (from AcOEt-hexane). $[\alpha]_D^{20} - 30.4^\circ$ ($c = 0.77$, CHCl_3). IR $\nu_{\text{max}}^{\text{KBr}} \text{cm}^{-1}$: 3280 (OH), $^1\text{H-NMR}$ (CDCl_3) δ : 0.96 (3H, s, CH_3C), 1.20–2.08 (12H, m), 2.15–2.38 (1H, m), 2.47 (1H, t, $J = 5$ Hz), 2.70 (1H, t, $J = 6$ Hz), 3.48–4.03 (8H, m). MS m/z : 282 (M^+). Anal. Calcd for $\text{C}_{16}\text{H}_{26}\text{O}_4 \cdot 1/2\text{H}_2\text{O}$: C, 65.94; H, 9.34. Found: C, 65.80; H, 9.41.

(-)-Norbourbonone Ethylene Ketal (36) Methanesulfonyl chloride (0.38 ml, 5 mmol) was added to a solution of (-)-34 (228 mg, 0.81 mmol) in pyridine (1 ml) at 0 °C, and the whole was stirred at room temperature for 36 h. After dilution with CH_2Cl_2 (30 ml), the mixture was washed successively with water, 10% HCl, water, and satd. NaHCO_3 , then dried

over MgSO_4 . Concentration gave dimesylate (**35**) (355 mg, quant.) as a pale yellow oil. A solution of **35** (355 mg, 0.81 mmol) in THF (30 ml) was added to a refluxing suspension of LiAlH_4 (0.18 g, 4.7 mmol) in THF (30 ml), and the whole was heated under reflux for 1.8 h and cooled to 0°C . Water (0.27 ml) and 15% NaOH (0.20 ml) were successively added to the mixture at 0°C , and the whole was stirred at room temperature for 1 h. After removal of precipitate by filtration, the filtrate was concentrated and chromatographed (silica gel, pentane-ether 20:1) to give (–)-**36** (157 mg, 78% from (–)-**34**) as a colorless oil. $[\alpha]_D^{20} -23.1^\circ$ ($c=0.78$, CHCl_3). IR $\nu_{\text{max}}^{\text{CHCl}_3} \text{cm}^{-1}$: 1100 (ether). $^1\text{H-NMR}$ (CDCl_3) δ : 0.85 (3H, d, $J=5$ Hz, $\text{CH}(\text{CH}_3)_2$), 0.91 (3H, d, $J=5$ Hz, $\text{CH}(\text{CH}_3)_2$), 0.98 (3H, s, CH_3C), 1.06–2.04 (12H, m), 2.06–2.30 (1H, m), 3.60–4.00 (4H, m, $\text{O}(\text{CH}_2)_2\text{O}$). MS m/z : Calcd for $\text{C}_{16}\text{H}_{26}\text{O}_2$ (M^+): 250.1930. Found: 250.1917.

(–)-**Norbourbonene** (**4**) A solution of (–)-**36** (146 mg, 0.58 mmol) in acetone (10 ml) containing *p*-toluenesulfonic acid (12 mg) was left standing at room temperature for 23 h. After concentration, the residue was taken up into ether (30 ml), and washed with satd. NaHCO_3 and satd. NaCl, successively, then dried over MgSO_4 . Evaporation of the solvent and column chromatography (silica gel, pentane-ether 10:1) gave (–)-**4** (98 mg, 82%) as colorless needles of mp $24.5\text{--}25.5^\circ\text{C}$ (lit.^{3b} $23\text{--}25^\circ\text{C}$). $[\alpha]_D^{25} -193^\circ$ ($c=0.38$, CHCl_3) (lit.^{3b}) $[\alpha]_D^{25} -213^\circ$ ($c=0.39$, CHCl_3), $[\alpha]_D^{25} -194^\circ$ ($c=0.38$, CHCl_3) (lit.^{4a}) $[\alpha]_D^{25} -181^\circ$ ($c=0.46$, CHCl_3). IR $\nu_{\text{max}}^{\text{neat}} \text{cm}^{-1}$: 1730 ($\text{C}=\text{O}$). $^1\text{H-NMR}$ (CDCl_3) δ : 0.86 (6H, d, $J=5$ Hz, $\text{CH}(\text{CH}_3)_2$), 1.13 (3H, s, CH_3C), 1.00–2.68 (13H, m). MS m/z : Calcd for $\text{C}_{14}\text{H}_{22}\text{O}$ (M^+): 206.1668. Found: 206.1662.

(–)- **β -Bourbonene** (**3**) A solution of BuLi (1.56 M in hexane, 0.47 ml, 0.73 mmol) was added to a stirred suspension of methyltriphenylphosphonium bromide (252 mg, 0.71 mmol) in ether (25 ml) at room temperature and the whole was stirred at room temperature for 2 h. A solution of (–)-**4** (71 mg, 0.34 mmol) in ether (3.5 ml) was added at room temperature and the whole was stirred at room temperature for 20 h. After addition of water, the mixture was extracted with pentane ($20\text{ ml} \times 3$), and the combined organic layer was dried over MgSO_4 . Concentration and column chromatography (silica gel, pentane) gave (–)-**3** (69 mg, 98%). $[\alpha]_D^{20} -97.0^\circ$ (neat) (lit.^{3a}) $[\alpha]_D^{20} -91.12^\circ$ (neat). IR $\nu_{\text{max}}^{\text{neat}} \text{cm}^{-1}$: 1655 ($\text{C}=\text{C}$). $^1\text{H-NMR}$ (CDCl_3) δ : 0.85 (6H, d, $J=5$ Hz, $\text{CH}(\text{CH}_3)_2$), 1.00 (3H, s, CH_3C), 1.04–2.00 (9H, m), 2.08–2.64 (4H, m), 4.71 (2H, brs, $\text{C}=\text{CH}_2$). MS m/z : Calcd for $\text{C}_{14}\text{H}_{24}$ (M^+): 204.1878. Found: 204.1879.

Acknowledgement We are grateful to Dr. Yoko Naya, Suntory Institute for Bioorganic Research, Osaka, Japan for providing spectra of (–)- **β -bourbonene**.

References and Notes

- Part XV: K. Tomioka, H. Masumi, T. Yamashita and K. Koga, *Tetrahedron*, **45**, 643 (1989).
- a) S. L. Fernandes, S. Y. Kamat and S. K. Paknikar, *Tetrahedron Lett.*, **21**, 2249 (1980); b) W. H. Gerwick, W. Fenical, D. V. Engen and J. Clardy, *J. Am. Chem. Soc.*, **102**, 7991 (1980); c) W. H. Germick, W. Fenical and M. U. S. Sultanbawa, *J. Org. Chem.*, **46**, 2233 (1981); d) W. H. Gerwick and W. Fenical, *ibid.*, **48**, 3325 (1983); e) B. N. Ravi and R. J. Wells, *Aust. J. Chem.*, **35**, 129 (1982).
- a) J. Krepinsky, Z. Samek, F. Sorm, D. Lamparsky, P. Ochsner and Y.-R. Navas, *Tetrahedron*, **S8**, 53 (1967); b) C. Gianotti and H. Schwang, *Bull. Soc. Chim. France*, **1968**, 2452.
- Synthesis of bourbonene sesquiterpenes. a) J. D. White and D. N. Gupta, *J. Am. Chem. Soc.*, **90**, 6171 (1968); b) M. Brown, *J. Org. Chem.*, **33**, 162 (1968); c) C. H. Heathcock and R. A. Bodger, *J. Chem. Soc., Chem. Commun.*, **1968**, 1510; d) K. Yoshihara, Y. Ohta, T. Sakai and Y. Hirose, *Tetrahedron Lett.*, **1969**, 2263; e) T. Uyehara, T. Ohnuma, T. Sato and K. Kato, *J. Chem. Soc., Chem. Commun.*, **1981**, 127.
- Synthesis of spatane diterpenes. R. G. Solomon, N. D. Sachinvala, S. R. Raychaudhari and D. B. Miller, *J. Am. Chem. Soc.*, **106**, 2211 (1984).
- M. Tanaka, K. Tomioka and K. Koga, *Tetrahedron Lett.*, **26**, 3035 (1985); *idem, ibid.*, **26**, 6109 (1985).
- A preliminary report of this work has appeared. K. Tomioka, M. Tanaka and K. Koga, *Tetrahedron Lett.*, **23**, 3401 (1982).
- Review: S. M. Baldwin, *Organic Photochemistry*, **5**, 123 (1981).
- Several examples of asymmetric (2+2) photocycloaddition reactions have been reported: J. J. Partridge, N. K. Chadha and M. R. Uskokovic, *J. Am. Chem. Soc.*, **95**, 532 (1973); J. R. Williams and J. F. Callahan, *J. Org. Chem.*, **45**, 4475, 4479 (1980); S. W. Baldwin and M. T. Crimmins, *J. Am. Chem. Soc.*, **104**, 1132 (1982); S. Jarosz and A. Zamojski, *Tetrahedron*, **38**, 1447, 1453 (1982); L. M. Tolbert and M. B. Ali, *J. Am. Chem. Soc.*, **104**, 1742 (1982); K. Bruneel, D. D. Keukeleire and M. Vandewelle, *J. Chem. Soc., Perkin Trans. 1*, **1984**, 1967; R. G. Solomon, N. D. Sachinvala, S. R. Raychaundhuri and D. B. Miller, *J. Am. Chem. Soc.*, **106**, 2211 (1984); G. L. Lange, C. Decicco, S. L. Tan and G. Chamberlain, *Tetrahedron Lett.*, **26**, 4707 (1985); G. L. Lange and M. Lee, *ibid.*, **26**, 6163 (1985); H. Herzog, H. Koch, H.-D. Sharf and J. Runsink, *Tetrahedron*, **42**, 3547 (1986); A. I. Meyers and S. A. Fleming, *J. Am. Chem. Soc.*, **108**, 306 (1986).
- a) K. Tomioka, F. Sato and K. Koga, *Heterocycles*, **17**, 311 (1982); b) K. Tomioka, T. Ishiguro, Y. Iitaka and K. Koga, *Tetrahedron*, **40**, 1303 (1984).
- For the application of **5** to the asymmetric synthesis of natural products, see: a) reviews: K. Tomioka, "Advances in Pharmaceutical Sciences," The Research Foundation for Pharmaceutical Sciences, Vol. I, 1985, p. 121; b) K. Tomioka, *Yakugaku Zasshi*, **104**, 1009 (1984). For leading references, see: c) K. Tomioka, Y.-S. Cho, F. Sato and K. Koga, *J. Org. Chem.*, **53**, 4094 (1988); d) K. Tomioka, M. Sugimori and K. Koga, *Chem. Pharm. Bull.*, **34**, 1501 (1986); e) K. Tomioka, H. Kawasaki, Y. Iitaka and K. Koga, *Tetrahedron Lett.*, **26**, 903 (1985).
- H. Kosugi, S. Sekiguchi, R. Sekita and H. Uda, *Bull. Chem. Soc. Jpn.*, **49**, 520 (1976); K. Ohga and T. Matsuo, *ibid.*, **49**, 1590 (1976); H. D. Scharf, J. Janus and E. Müller, *Tetrahedron*, **35**, 25 (1979).
- K. Ohga and T. Matsuo, *Bull. Chem. Soc. Jpn.*, **43**, 3505 (1970).
- M. Tanaka, K. Tomioka and K. Koga, *Heterocycles*, **23**, 2347 (1985).
- H. W. Wanzlick, G. Gollmer and H. Milz, *Chem. Ber.*, **88**, 69 (1955).
- Melting points were measured using Büchi 510 melting point apparatus and are not corrected. Optical rotations were taken with a JASCO DIP-181 automatic polarimeter. IR spectra were taken with a JASCO Infrared Spectrometer Model DS-402 G and a JASCO IRA-I Grating Infrared Spectrometer. $^1\text{H-NMR}$ spectra were taken with a JEOL FX-100 Spectrometer at 100 MHz, or with a Hitachi R-24B Spectrometer at 60 MHz. Chemical shift values are expressed in ppm relative to internal tetramethylsilane. Abbreviations are as follows: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; br, broad. MS spectra were taken with a JEOL JMS DX-300 Mass Spectrometer.
- G. B. Payne, *J. Org. Chem.*, **23**, 310 (1958).
- Extremely low specific rotation has been reported for this compound. R. M. Ortuño, J. Bigorra and J. Font, *Tetrahedron*, **43**, 2199 (1987) and references cited therein.