Studies on Cytotoxic Constituents in Pericarps of *Mallotus japonicus*. IV

Munehisa ARISAWA,*^a Akio FUJITA,^a Toshimitsu HAYASHI,^a Naokata MORITA,^a Tohru KIKUCHI,^b and Yasuhiro TEZUKA^b

Department of Medicinal Resources, Faculty of Pharmaceutical Sciences,^a and Department of Natural Product Chemistry, Research Institute for Wakan-Yaku (Oriental Medicine),^b Toyama Medical and Pharmaceutical University, 2630, Sugitani, Toyama 930-01, Japan. Received July 8, 1989

Two new phloroglucinol derivatives, isomallotolerin (**1**) and isomallotochromanol (**2**), were isolated from the cytotoxic fraction of the pericarps of *Mallotus japonicus*. The new derivatives were identified as 3-(3-methyl-2-hydroxybut-3-enyl)-5-(3-acetyl-2,4-dihydroxy-5-methyl-6-methoxybenzyl)-phlorisobutyrophenone (**1**) and 6-acetyl-5,7-dihydroxy-8-(3-acetyl-2,4-dihydroxy-5-methyl-6-methoxybenzyl)-2,2-dimethyl-3-hydroxychroman (**2**) from chemical and spectral data. Isomallotolerin and its acetate were found to be cytotoxic to KB cell line.

Keywords *Mallotus japonicus*; Euphorbiaceae; phloroglucinol derivative; cytotoxicity; ¹³C-NMR

Previously, we reported several cytotoxic phloroglucinol derivatives in pericarps of *Mallotus japonicus* MUELL. ARG. (Euphorbiaceae).^{1–3)} In a continuing search for cytotoxic constituents in the CHCl₃-soluble fraction of the pericarps of *M. japonicus*, two new compounds named isomallotolerin (**1**) and isomallotochromanol (**2**) were isolated. We wish to report the structural elucidation and the cytotoxic activities of these new compounds. Separation of the extract of *M. japonicus* by column chromatography on Si gel³⁾ yielded **2** and a mixture of **1** and mallotolerin (**3**). The isolation of **1** was performed by high-performance liquid chromatography (HPLC).

Compound **1**, C₂₆H₃₂O₉, gave a positive FeCl₃ reaction. The ultraviolet (UV) spectrum was similar to that of **3**. The proton nuclear magnetic resonance (¹H-NMR) spectrum closely resembled that of **3**, except for the appearance of the signals of an isopropyl ketone group at δ 1.19 (6H, d, *J*=6.7 Hz) and 4.05 ppm (1H, sept, *J*=6.7 Hz) instead of the signals of a propyl ketone. The carbon-13 nuclear magnetic resonance (¹³C-NMR) spectrum of **1** was also similar to that of **3**, except for the appearance of the carbon signals of an isopropyl group at δ 18.93(q), 19.12(q) and 38.97 ppm(d) instead of the signals of a propyl group. The mass spectrum (MS) of **1** showed a molecular ion peak at *m/z* 488 and prom-

inent peaks at *m/z* 274, 259, 221, 209, 196 and 181, indicating a 3-acetyl-2,4-dihydroxy-5-methyl-6-methoxybenzyl moiety.^{1–3)} Reductive alkaline cleavage of **1** afforded 2,6-dihydroxy-3-methyl-4-methoxyacetophenone.¹⁾ From these chemical and spectral data and from biosynthetic considerations, the structure of **1** is proposed to be 3-(3-methyl-2-hydroxybut-3-enyl)-5-(3-acetyl-2,4-dihydroxy-5-methyl-6-methoxybenzyl)-phlorisobutyrophenone, and it was named isomallotolerin (**1**). A Cotton effect was not detected in a circular dichroism (CD) study on its benzoate^{4,5)} and **1** was converted into the L-menthoxyacetyl derivatives, which showed two peaks on HPLC, the areas of which were equal.⁶⁾ From these results, the allylic alcohol on its side chain is considered to be racemic.

Compound **2**, C₂₄H₂₈O₉, also gave a positive FeCl₃ reaction. The UV, IR and MS were similar to those of mallotochromanol (**4**). The ¹H-NMR spectrum closely resembled that of **4** with a slight difference in the methylene signals at δ 3.07 (1H, dd, *J*=15.1, 8.3 Hz) and 3.18 ppm (1H, dd, *J*=15.1, 9.3 Hz) and a methine proton at δ 4.87 ppm (1H, dd, *J*=9.3, 8.3 Hz). The signal of the methylene protons between the rings of **4** shifted upfield on acetylation, whereas the signal of the methylene protons between the rings of **2** showed a down-

Chart 1

TABLE I. ^{13}C -NMR Chemical Shifts of **1**–**4**

Carbon No.	1 ^{a)}	3 ^{b)}	2 ^{c)}	4 ^{b)}
1	104.32 ^{d)}	104.94 ^{d)}		
2	159.68 ^{e)}	159.71 ^{e)}	69.96	78.49
3	106.31 ^{d)}	106.07 ^{d)}	89.96	68.49
4	159.79 ^{e)}	159.81 ^{e)}	26.99	26.09
5	104.61 ^{d)}	105.32 ^{d)}	154.54	160.72
6	160.12 ^{e)}	160.12 ^{e)}	104.77	105.20
7	211.87	207.55	158.69	160.91
8	38.97	46.04	101.52	105.01
9	18.93	18.13	165.05	155.08
10	19.12	14.02	103.66	99.47
11			22.94	22.02
12			26.34	24.84
13			202.46	203.79
14			32.44	32.89
1'	108.53	108.73	112.10	108.79
2'	157.16	157.39	163.54	157.22
3'	108.67	109.15	108.22	108.79
4'	162.02	162.47	158.97	162.56
5'	109.17	109.48	108.57	109.95
6'	159.99	159.90	163.65	159.95
7'	205.57	205.62	204.68	205.56
8'	33.50	33.77	33.09	33.75
1''	29.20	29.31		
2''	78.11	78.16		
3''	146.81	146.75		
4''	110.02	110.67		
5''	18.10	18.39		
Ar-CH ₂ -Ar	16.83	17.04	16.85	16.55
Ar-Me	8.52	8.82	8.84	8.27
OMe	61.58	61.83	60.01	61.81

a) Measured in $\text{CDCl}_3 + \text{CD}_3\text{OD}$. b) Measured in CDCl_3 . c) Measured in $\text{DMSO}-d_6$. d, e) Assignments may be interchanged.

TABLE II. Cytotoxicities of Phloroglucinol Derivatives Against KB Cells

Compound	ED ₅₀ (μg/ml)	Compound	ED ₅₀ (μg/ml)
1	0.84	1 -Hexaacetate	1.5
2	>20	2 -Tetraacetate	8.5
3	0.95	3 -Hexaacetate	3.5
4	>20	4 -Pentaacetate	8.6

field shift like that found in mallotophenone **5** after conversion to acetates.¹⁾ The ^{13}C -NMR spectrum was also similar to that of **4**, except for the signal due to C-3 which was shifted significantly downfield in comparison with that of **4**. The assignment of the ^1H - and ^{13}C -NMR signals of **2** was done by means of two-dimensional NMR with long range ^1H - ^{13}C shift correlation and by making use of the deuterium induced up-field shifts of alcoholic carbons in the ^{13}C -NMR spectrum.⁷⁾ From these spectral data, the structure of **2** is proposed to be 6-acetyl-5,7-dihydroxy-8-(3-acetyl-2,4-dihydroxy-5-methyl-6-methoxybenzyl)-2,2-dimethyl-3-hydroxychroman, and it was named isomallotochromanol (**2**). The hydroxy group on the chroman ring is considered to be racemic based on the CD study⁸⁾ and separation of diastereomeric derivatives by HPLC, as in the case of compound **1**.

The isolated compounds **1**, **2** and their acetates were tested for cytotoxic activity in the KB system.¹⁾ While both **2** and its tetraacetate were inactive, as was **4**, **1** and its

hexaacetate showed significant cytotoxicity with ED₅₀ of 0.84 and 1.5 μg/ml, respectively. Further cytotoxic constituents of this plant are now under investigation.

Experimental

General Procedures All melting points were determined on a Yanagimoto micro melting point apparatus and are recorded uncorrected. UV and IR spectra were recorded on a Hitachi 220S double beam spectrophotometer and 260-10 infrared spectrometer with polystyrene calibration at 1601 cm^{-1} , respectively. Specific rotations were determined on a JASCO DIP-140 digital polarimeter. ^1H - and ^{13}C -NMR spectra were taken on a Varian XL-200 spectrometer at 200 and 50.3 MHz, respectively, and 2D-NMR spectra were taken on a JEOL JNM-GX 400 spectrometer with tetramethylsilane as an internal standard. The chemical shifts are recorded in δ (ppm) values. Mass spectra were obtained on a JEOL JMS-D-200 mass spectrometer operating at 70 eV. HPLC was performed on a Shimadzu LC-6A liquid chromatograph with an SPD-MIA spectrophotometric detector.

Extraction and Separation The extraction and separation of the dried pericarps of *M. japonicus* have been described previously.¹⁻³⁾ The 1% MeOH/ CHCl_3 eluent (26 g) was rechromatographed on a silica gel column (benzene : AcOEt = 4 : 1) to give **2** (25 mg). The crude **1** and **3** from the 15% AcOEt/hexane eluate of the rechromatography on a silica gel column³⁾ were purified by preparative HPLC (column, Cosmosil 5C₁₈ 20 mm i.d. $\times$ 25 cm; solvent, CHCl_3 : MeOH : 0.005 M phosphate buffer = 1 : 35 : 5; flow rate 5 ml/min; detection, UV 282 nm) to give **1** (13 mg, t_R 110 min) and **3** (30 mg, t_R 115 min), respectively.

Isomallotolerin (1) Yellow needles, mp 216–217°C (CHCl_3). $[\alpha]_D^{23} \pm 0^\circ$ ($c = 0.63$, CHCl_3). Anal. Calcd for $\text{C}_{26}\text{H}_{32}\text{O}_9$: C, 63.91; H, 6.61. Found: C, 63.72; H, 6.55. UV $\lambda_{\text{max}}^{\text{EtOH}}$ nm (log ϵ): 285 (4.04), 323 (3.86). IR $\nu_{\text{max}}^{\text{KBr}}$ cm^{-1} : 3440 (OH), 3280, 1620 (C=O), 1605, 1420, 1300, 1280, 1145. ^1H -NMR (CDCl_3) δ : 1.19 (6H, d, $J = 6.7$ Hz, Me $\times$ 2), 1.86 (3H, s, 5'-Me), 2.12 (3H, s, 5'-Me), 2.69 (1H, dd, $J = 15.0$, 8.9 Hz, 1''-Ha), 2.73 (3H, s, Ac), 3.14 (1H, dd, $J = 15.0$, 1.8 Hz, 1''-Hb), 3.74 (2H, s, Ar-CH₂-Ar), 3.98 (3H, s, OMe), 4.05 (1H, sept, $J = 6.7$ Hz, 8-H), 4.32 (1H, br d, $J = 8.9$ Hz, 2''-H), 4.90 (1H, br s, 4''-Ha), 5.03 (1H, br s, 4''-Hb). MS m/z 488 (M^+), 470, 417, 274, 259, 221, 209, 196, 181. High-resolution MS measurement m/z 488.2079 ($\text{C}_{26}\text{H}_{32}\text{O}_9$ requires 488.2044). ^{13}C -NMR see Table I.

Acetylation of 1 Compound **1** was treated overnight with Ac₂O and pyridine at room temperature, and the reaction mixture was worked up as usual to give a hexaacetate as a colorless oil. ^1H -NMR (CDCl_3) δ : 1.10 (6H, d, $J = 6.8$ Hz, Me $\times$ 2), 1.71 (3H, s, Me), 1.94 (3H, s, OAc), 2.06 (3H, s, Ar-Me), 2.13 (3H, s, OAc), 2.14 (3H, s, OAc), 2.23 (3H, s, OAc), 2.26 (3H, s, OAc), 2.29 (3H, s, OAc), 2.36 (3H, s, Ac), 2.56 (1H, m, 1''-Ha), 2.82 (1H, m, 1''-Hb), 2.88 (1H, m, 8-H), 3.56 (3H, s, OMe), 3.67 (2H, s, Ar-CH₂-Ar), 4.80 (2H, br s, 4''-H₂), 5.28 (1H, dd, $J = 7.3$, 6.6 Hz, 2''-H). MS m/z : 740 (M^+).

Isomallotochromanol (2) Yellow needles, mp 150–152°C (MeOH). $[\alpha]_D^{23} \pm 0^\circ$ ($c = 0.27$, EtOH). Anal. Calcd for $\text{C}_{24}\text{H}_{28}\text{O}_9$: C, 62.59; H, 6.09. Found: C, 62.48; H, 6.09. UV $\lambda_{\text{max}}^{\text{EtOH}}$ nm (log ϵ): 288 (4.40), 345 (3.74). IR $\nu_{\text{max}}^{\text{KBr}}$ cm^{-1} : 3480 (OH), 3430, 2930, 1630 (C=O), 1595, 1440, 1370, 1130, 1095. ^1H -NMR (CDCl_3) δ : 1.25 (3H, s, Me), 1.41 (3H, s, Me), 2.14 (3H, s, Ar-Me), 2.68 (3H, s, Ac), 2.71 (3H, s, Ac), 3.07 (1H, dd, $J = 15.1$, 8.3 Hz, 4-Ha), 3.18 (1H, dd, $J = 15.1$, 9.3 Hz, 4-Hb), 3.98 (3H, s, OMe), 3.70 (2H, d, $J = 1.71$ Hz, Ar-CH₂-Ar), 4.87 (1H, dd, $J = 9.3$, 8.3 Hz, 3-H). MS m/z : 460 (M^+), 265, 209, 196, 181. High-resolution MS measurement m/z 460.1712 ($\text{C}_{24}\text{H}_{28}\text{O}_9$ requires 460.1732). ^{13}C -NMR see Table I.

Acetylation of 2 After acetylation as described for **1**, a tetraacetate was obtained as a colorless oil. ^1H -NMR (CDCl_3) δ : 1.00 (3H, s, Me), 1.02 (3H, s, Me), 2.03 (3H, s, Ar-Me), 2.09 (3H, s, OAc), 2.27 (6H, s, OAc $\times$ 2), 2.30 (3H, s, OAc), 2.36 (3H, s, Ac), 2.39 (3H, s, Ac), 2.92 (1H, dd, $J = 15.7$, 9.5 Hz, 4-Ha), 3.00 (1H, dd, $J = 15.7$, 8.4 Hz, 4-Hb), 3.67 (1H, d, $J = 15.8$ Hz, Ar-HCH-Ar), 3.72 (3H, s, OMe), 3.90 (1H, d, $J = 15.8$ Hz, Ar-HCH-Ar), 4.55 (1H, m, 3-H). MS m/z : 628 (M^+), 570, 528, 486, 249, 238, 209, 196.

Preparation of Benzoates of 1 and 2 Two drops of benzoyl chloride were added to a dry pyridine solution of **1** (5 mg), with ice cooling. The reaction mixture was evaporated *in vacuo* to obtain the crude benzoate, which was purified by preparative TLC with benzene-AcOEt (19 : 1) as the developing solvent. Recrystallization from MeOH gave the benzoate as colorless prisms, mp 110–112°C. UV $\lambda_{\text{max}}^{\text{EtOH}}$ nm: 233. Similar treatment of **2** (5 mg) gave its benzoate as colorless prisms, mp 94–96°C. UV $\lambda_{\text{max}}^{\text{EtOH}}$ nm: 234. These benzoates were examined by CD (EtOH) but showed no Cotton effect.

Analysis of L-Methoxyacetyl Derivatives of 1 and 2 Two drops of L-menthoxyacetyl chloride were added to a dry pyridine solution of **1** (3 mg), at room temperature. After 12 h, water was added to the solution and the solvent was removed *in vacuo*. The residue was dissolved in chloroform (5 ml) and washed with water (3 ml $\times$ 2). After being dried (Na_2SO_4), the solution was evaporated *in vacuo* and the residue was purified by preparative TLC with hexane-ethyl acetate (10:3) as the developing solvent. The diastereomeric mixture was obtained as a colorless viscous oil, which showed two peaks (t_R 18 and 19 min, respectively), the areas of which were equal on HPLC, under the following conditions: column, TSKgel Silica-60 (4.6 mm i.d. $\times$ 250 nm); mobile phase, cyclohexane-ether (9:1); flow rate, 0.5 ml/min; temperature, 32 °C; detection, UV 254 nm. Similar treatment of **2** gave its L-menthoxyacetyl derivative as a colorless viscous oil, which showed two peaks (t_R 12 and 13 min, respectively) the areas of which were equal on HPLC, under the following conditions: column, Cosmosil 5C₁₈ (4.6 mm i.d. $\times$ 150 mm); mobile phase, acetonitrile-H₂O (99:1); flow rate, 1.0 ml/min; temperature, 32 °C; detection, UV 254 nm.

Reductive Alkaline Cleavage Reductive alkaline cleavage was carried out as described previously.¹⁾

Cytotoxicity Test The test employing KB cell line was carried out as

described previously.¹⁾

Acknowledgments The authors thank Mr. M. Morikoshi, analytical center of our University, for measurement of mass spectra.

References and Notes

- 1) M. Arisawa, A. Fujita, R. Suzuki, T. Hayashi, N. Morita, N. Kawano, and S. Koshimura, *J. Nat. Prod.*, **48**, 455 (1985).
- 2) M. Arisawa, A. Fujita, M. Saga, T. Hayashi, N. Morita, N. Kawano, and S. Koshimura, *J. Nat. Prod.*, **49**, 298 (1986).
- 3) A. Fujita, T. Hayashi, M. Arisawa, M. Shimizu, and N. Morita, *J. Nat. Prod.*, **51**, 708 (1988).
- 4) N. Harada, J. Iwabuchi, Y. Yokota, H. Uda, and K. Nakanishi, *J. Am. Chem. Soc.*, **103**, 5590 (1981).
- 5) N. C. Gonnella, K. Nakanishi, V. S. Martin, and K. B. Sharpless, *J. Am. Chem. Soc.*, **104**, 3775 (1982).
- 6) R. A. Halpin, S. F. El-Naggar, K. M. McCombe, K. P. Vyas, D. R. Boyd, and D. M. Jerina, *Tetrahedron Lett.*, **23**, 1655 (1982).
- 7) P. E. Pfeffer, K. M. Valentine, and F. W. Parrish, *J. Am. Chem. Soc.*, **101**, 1265 (1979).
- 8) N. Harada and K. Nakanishi, *Accounts Chem. Res.*, **5**, 257 (1972).